

## **Nucleic Acids, Proteins, and Antibodies**

### ***Statement under 37 C.F.R. § 1.77(b)(4)***

[001] This application refers to a "Sequence Listing" listed below, which is provided as an electronic document on two identical compact discs (CD-R), labeled "Copy 1" and "Copy 2." These compact discs each contain the file "PC007C1\_seqList.txt" (4,826,998 bytes, created on March 6, 2002), which is hereby incorporated in its entirety herein.

[002] The Sequence Listing may be viewed on an IBM-PC machine running the MS-Windows operating system.

### ***Field of the Invention***

[003] The present invention relates to novel cardiovascular system related polynucleotides, the polypeptides encoded by these polynucleotides herein collectively referred to as "cardiovascular system antigens," and antibodies that immunospecifically bind these polypeptides, and the use of such cardiovascular system polynucleotides, antigens, and antibodies for detecting, treating, preventing and/or prognosing disorders of the cardiovascular system, including, but not limited to, the presence of cancer of the cardiovascular system tissues and cancer metastases. More specifically, isolated cardiovascular system nucleic acid molecules are provided

encoding novel cardiovascular system polypeptides. Novel cardiovascular system polypeptides and antibodies that bind to these polypeptides are provided. Also provided are vectors, host cells, and recombinant and synthetic methods for producing human cardiovascular system polynucleotides, polypeptides, and/or antibodies. The invention further relates to diagnostic and therapeutic methods useful for diagnosing, treating, preventing and/or prognosing disorders related to the cardiovascular system, including of cancer of the cardiovascular system tissues, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying agonists and antagonists of polynucleotides and polypeptides of the invention. The invention further relates to methods and/or compositions for inhibiting or promoting the production and/or function of the polypeptides of the invention.

### ***Background of the Invention***

- [004] The Human Cardiovascular, or circulatory, system is responsible for the delivery of oxygen, nutrient molecules, and hormones, in addition to the removal of carbon dioxide via blood. The system is comprised of the heart and a complex system of arteries, arterioles, capillaries, venules, and veins that innervate and affect the entire body. The circulatory system provides the primary mechanism of transport of materials between the organs and tissues of the body.
- [005] The central organ of the circulatory system is the heart, a muscular pump responsible for the propulsion of blood throughout the body. The heart is a two-sided, four chambered structure with muscular walls, contracting and relaxing in a cyclical pattern, or cardiac cycle. The cardiac cycle consists of two parts: systole (e.g., the contraction of the heart muscle) and diastole (e.g., relaxation of the heart muscle). Blood from the body flows through the vena cava into the right atrium (or upper chamber) while oxygenated blood from the lungs flows from the pulmonary vein into the left atrium. Atrial systole occurs as the muscles of both atria contract, forcing blood downward through each atrioventricular (AV) valve into the corresponding ventricle. Diastole occurs as the ventricles are filling with blood. Ventricular systole opens the semilunar, or arterial, valve, forcing the blood out of the ventricles through the pulmonary artery or aorta, depending on the oxygenated state of the blood. The pulmonary artery carries deoxygenated blood to the lungs for gas exchange. The



aorta, the main artery leaving the heart, is responsible for transporting the oxygenated blood from the heart to the network of arteries and capillaries in the body.

[006] Blood vessels that carry blood away from the heart (e.g., arteries) are composed of thick smooth muscle fibers that allows them to expand and contract which enables blood to be carried under high blood pressure. Arterioles are small arteries that connect larger arteries with collections of capillaries (e.g., capillary beds). Capillaries are tiny, extremely thin-walled vessels that act as bridges between arteries and the surrounding tissues. Nutrients, wastes, and hormones are exchanged across the thin walls of capillaries by passing through or between the cells that line the capillary. Some capillaries have small pores between the cells of the capillary walls that allow material exchange as well as the passage of white blood cells.

[007] Blood leaving the capillary beds flows into a progressively larger series of venules that eventually will join to form veins. Because veins carry blood from the capillaries to the heart, most veins carry oxygen-poor blood. The exceptions, pulmonary veins, carry oxygenated blood from the lungs back to the heart via the vena cava. Unlike arteries, pressure in the veins is low; therefore, veins depend on muscular contractions to move the blood through them.

[008] Cardiovascular disease can affect all aspects of the circulatory system, from the heart muscles and valves to the arteries and veins. These diseases can range in severity from mild to severe and acute to chronic. Cardiovascular diseases can be grouped according to the area of the circulatory afflicted, the cause of the affliction, or the result of the affliction.

#### *Disorders and Diseases of the Heart*

[009] The most common heart disease is Heart Failure in which the cardiac output of the heart is insufficient to meet the body's normal requirements for oxygen and nutrients. Any disease can lead to heart failure if (a) it increases the heart muscle's workload (e.g., myocarditis, diabetes, coronary artery disease) which eventually weakens the force of contractions, or (b) affects the heart's electrical conduction system (e.g., hyperkalemia) which results in slow, fast, or irregular heartbeats. The body has several mechanisms to initially compensate for heart failure; however, these mechanisms eventually will malfunction and cause edema, making the heart failure worse.

[0010] Several treatment regimens are available for chronic and acute heart failure including dietary adjustments and exercise regimens. Diuretics are used to reduce fluid retention. Other drugs are used to increase the power of a heartbeat and slows a rapid heart rate (e.g., digoxin) and to dilate the blood vessels (e.g., vasodilators, including angiotensin converting enzyme, nitroglycerin, and hydralazine). Heart transplantation may be also recommended if there is no response to any other regimen.

[0011] Cardiomyopathy is group of disorders characterized by an alteration of the structure or an impairment of the function of the muscular wall of the ventricles. The most common form is dilated congestive cardiomyopathy in which the ventricle cavities enlarge, are unable to produce enough cardiac output, and eventually result in congestive heart failure. Viral cardiomyopathy is a form of dilated congestive cardiomyopathy resulting from a viral infection (e.g., coxsackievirus B). Treatments for dilated congestive cardiomyopathy involve managing angina with a nitrate, beta-blocker, or calcium channel blocker. Anticoagulant drugs are also used to prevent embolus. Hypertrophic cardiomyopathy is an inherited genetic defect resulting in thickening of the ventricle walls and possibly blockage of the blood flow into the ventricle. Treatment regimens for hypertrophic cardiomyopathy are aimed primarily at reducing the heart's resistance to filling with blood between heartbeats by the use of beta-blockers and calcium channel blockers. Surgery can also be used to relieve symptoms. Restrictive cardiomyopathy is a rare disorder in which the walls of the ventricles become stiff without thickening and resist normal filling with blood between heartbeats. There are two basic types of restrictive cardiomyopathy: one in which the heart muscle is gradually replaced by scar tissue, and the other in which the heart muscle is infiltrated by abnormal material (e.g., white blood cells, hemochromatosis, amyloidosis, sarcoidosis, or tumor invading the heart tissue). No current therapy for restrictive cardiomyopathy is satisfactory. The therapies for other forms of cardiomyopathy and heart failure worsen this disorder instead of improving it.

[0012] Primary tumors of the heart are generally rare, and their symptoms imitate other heart diseases. Half of all primary heart tumors are myxomas, with most occurring in the left atrium. Myxomas are noncancerous tumors, irregular in shape and jellylike in consistency. The tumor may block or damage the mitral valve or another valve if

pieces of the tumor break off. Symptoms depend on which vessel is blocked. Less common noncancerous heart tumors include fibromas and rhabdomyomas. Single noncancerous primary tumors are generally removed by surgery. Multiple noncancerous and any cancerous heart tumor can not be removed surgically; only their symptoms are treated.

#### *Disorders and Diseases of the Pericardium*

[0013] The pericardium is a flexible, two-layered sac that surrounds the heart and is responsible for keeping the heart in position, preventing the heart from overfilling with blood, and protecting the heart from chest infections. Although the pericardium is not essential to the heart's performance, inflammation of the pericardium (e.g., pericarditis), can be painful and result in heart damage. Acute pericarditis occurs when inflammation of the pericardium begins suddenly, resulting in fluid and blood products (e.g., fibrin, red blood cells, and white blood cells) pouring into the pericardial space and compressing the heart. Acute pericarditis has many causes ranging from viral infections to diseases such as AIDS, systemic lupus erythematoses, rheumatoid disease, and kidney failure. Treatment includes analgesics or anti-inflammatory drugs to relieve pain, and in the cases of viral or bacterial infections, antibiotics are also administered.

[0014] Chronic pericarditis is inflammation that results in gradual fluid accumulation and thickening of the pericardium. One form, chronic effusive pericarditis, occurs when fluid slowly accumulates in the pericardium. A rare form, chronic constrictive pericarditis, results when fibrous tissue forms around the heart, compressing the heart, and making the heart smaller. The compression increases the venous pressure resulting in the leaking out and accumulation of fluid in the body. Current protocols for treatment involve diuretics to remove excess fluid or surgery to remove the pericardium.

#### *Disorders and Diseases of the Endocardium and Heart Valves*

[0015] Bacterial Endocarditis is an inflammation of the smooth interior lining of the heart (e.g., endocardium) or heart valve occurring most often in people with a heart defect or damaged valves. This disease can appear suddenly and become life-threatening within days (e.g., acute bacterial endocarditis) or can develop gradually over a period

of weeks to several months (e.g., subacute bacterial endocarditis). As a preventive measure, people with heart valve abnormalities, artificial valves, or congenital defects are given antibiotics prior to any dental or surgical procedure. Current treatment for bacterial endocarditis consists of a series of high-dose intravenous antibiotics. Heart surgery may also be required to repair or replace damaged valves and remove accumulations of bacteria and blood clots on the valves (e.g., vegetations).

[0016] Regurgitation describes the leakage back through a valve upon ventricular contraction. The mitral valve opens from the left atrium into the left ventricle. When this valve develops regurgitation, some blood leaks back into the left atrium, increasing the volume and blood pressure, and resulting in lung congestion. The aortic valve opens from the left ventricle into the aorta. Regurgitation of the aortic valve results in a characteristic heart murmur, in mild cases, and enlargement of the ventricle and eventually heart failure in severe cases. The tricuspid valve opens from the right atrium into the right ventricle. Leakage of this valve results in blood pumped forward to the lungs and back into the right atrium and pressure increased in the right atrium and veins, creating resistance to the flow of blood from the body to the heart. Treatment for regurgitation depends on the valve that is leaking. Mitral valve regurgitation is generally treated with surgery, or if atrial fibrillation is also present, with drugs such as beta-blockers, digoxin, and verapamil to slow the heart rate and control the fibrillation. Treatment for aortic valve regurgitation involves reducing the risk for infection of the valve by antibiotics. Currently, there is no treatment for tricuspid valve regurgitation.

[0017] Stenosis is the narrowing of a valve opening that increases resistance to blood flow across the valve. Mitral valve stenosis occurs due to a congenital defect or rheumatic fever and, in the severe cases, results in heart failure and pulmonary edema. Drug therapy (e.g., beta-blockers and digoxin to control fibrillation and diuretics to reduce blood volume and pressure in the lungs) and valve replacement surgery is the current treatment for this stenosis. In addition to the scarring and calcium accumulation in the leaflets of the aortic valve, a congenital defect or rheumatic fever can also cause aortic stenosis. The ventricle wall thickens, requiring an increasing blood supply from the coronary arteries, and resulting in angina, coronary artery disease, and eventually, heart failure. Stenosis can also occur in the tricuspid valve and pulmonary valve; however, it is rarely severe enough to require surgery.

*Diseases and Disorders of the Arteries*

[0018] Arteriosclerosis is a general term for arterial diseases in which the wall of an artery becomes thicker and less elastic, resulting in an obstruction of the blood flow. Arteriolosclerosis, primarily caused by high blood pressure, is a less common form of arteriosclerosis that affects the inner and middle layer of the walls of arterioles. The most common, atherosclerosis, occurs when fatty material accumulates under the inner lining of the arterial wall. Artherosclerosis can affect the arteries of the brain, heart, kidneys, and other vital organs of the body and results in a severe narrowing of the vessel or a rupture in the vessel, triggering the formation of thrombi. The thrombus may further narrow or occlude the artery, or it may detach and cause an embolism. Treatment for artherosclerosis involves preventing by controlling the risk factors (e.g., high blood pressure, high blood cholesterol levels, cigarette smoking, diabetes) associated with the disease.

[0019] The major cause of cardiovascular disease, coronary artery disease, is arteriosclerosis of the coronary arteries encircling the heart. As coronary artery disease progresses, several major complications can occur, including myocardial ischemia, angina, and myocardial infarction. Ischemia is an inadequate supply of oxygenated blood to the heart that results in heart damage. Angina, or angina pectoris, is the chest pain or pressure sensation that occurs when the heart muscle does not receive enough oxygen. Although ischemia is usually accompanied by an episode of angina, it can occur alone (e.g., silent ischemia).

[0020] Angina pectoris generally occurs upon exertion when the heart's needs increase and the blood flow is no longer enough to meet those needs. However, variant angina, or Prinzmetal's angina, occurs at rest from a spasm of the large coronary arteries on the surface of the heart. Unstable angina occurs when the pattern of symptoms changes, usually reflecting a rapid progression of coronary artery disease. Unstable angina corresponds to a high risk for heart attack and usually translates into a medical emergency. Treatment for both angina pectoris and Prinzmetal's angina is designed to prevent or reduce ischemia and minimize symptoms through drug therapy. Four types of drugs are available: beta-blockers to reduce the resting heart rate and demand for oxygen, nitrates to dilate the blood vessel walls, calcium antagonists to prevent vessel constrictions, and antiplatelet drugs to prevent clot

formation. Unstable angina is currently treated with anticoagulants (e.g., heparin), glycoprotein IIb/IIIa inhibitor (e.g., abciximab) beta-blockers, and intravenous nitroglycerin. If drugs are not effective, coronary arteriography or bypass surgery may be required.

#### *Diseases and Disorders of the Veins*

- [0021] The main disorders of the veins involve inflammation, clotting, and defects that lead to distention and varicose veins. Varicose veins are enlarged superficial veins most commonly found in the legs. The veins become weak, elongate, and widen, causing the valve cusps to separate, and the veins rapidly fill with blood when the person stands. Varicose veins commonly ache. Although there is no cure, treatment relieves symptoms, improves appearance, and prevents complication. Surgery may also be able to remove the veins.

#### *Disorders Associated with Blood Pressure*

- [0022] Blood pressure varies naturally over a lifetime. Adjustments in blood pressure are governed by changes in kidney function and in the autonomic nervous system. The most common disorder associated with blood pressure is hypertension or high blood pressure. Hypertension is defined as an average at-rest systolic pressure of 140 mm Hg or more and an average at-rest diastolic pressure of 90 mm Hg or more. Generally, both the systolic and diastolic pressures are elevated. However, in isolated systolic hypertension, only the systolic pressure is elevated. Hypertension probably has more than one cause. Several changes in the heart and blood pressure may combine to elevate blood pressure. Other causes may be kidney disease, hormonal disorders, and drugs. Hypertension increases the risk of developing heart disease, kidney failure, and especially, stroke. Hypertension can not be cured. Treatment involves drug therapy to prevent complications. Current drug regimens include thiazide diuretics to help eliminate salt and water from the kidneys and lower fluid volume, adrenergic blockers (e.g., alpha-blockers, beta-blockers, and alpha-beta blockers) to block the effects of the sympathetic nervous system, angiotensin converting enzyme inhibitors and angiotensin II blockers to dilate arteries, calcium antagonists to dilate blood vessels, and direct vasodilators.

[0023] Low blood pressure can also cause problems. For example, shock results when a low blood volume, an inadequate pumping of the heart, or excessive dilation of the blood vessels causes severe low blood pressure. An inadequate supply of blood reaches the cells, which can be quickly and irreversibly damaged. Low blood volume can occur with excessive loss of body fluids associated with such conditions as pancreatitis, perforation of the intestinal wall, severe diarrhea, excessive use of diuretics, or kidney disease. If untreated, shock is usually fatal. Treatment regimens are aimed at increasing the rate and rhythm of the heartbeat, increasing blood volume, and improve heart muscle contraction.

#### *Disorders Associated with Electrolyte Imbalance*

[0024] Changes in the total amount of sodium are closely linked to changes in the volume of water in the blood. Normal kidneys adjust the amount of sodium excreted in the urine so that sodium levels remain relatively constant. Hyponatremia, or low sodium blood level, occurs when the blood sodium concentration falls below 136 milliequivalents (mEq) per liter of blood. A reduction in the blood sodium concentration results in a reduction in blood volume and blood pressure. The heart rate rises and light-headedness and shock can occur. Treatment involves the restriction of fluid intake and a slow increase in the blood sodium levels with intravenous fluids. Hypernatremia, high sodium blood level, occurs when the blood sodium concentration reaches above 145 mEq per liter of blood. This causes an increase in blood volume, resulting in edema. Treatment involves replacing the body's water concentration.

[0025] Changes in potassium concentration can have serious consequences, such as an abnormal heart rhythm or cardiac arrest. Therefore, potassium levels in the blood must be maintained within a narrow range. If the kidneys are not functioning properly or potassium has been lost through the gastrointestinal tract, potassium levels in the blood can drastically drop. Hypokalemia occurs when the blood potassium level falls below 3.8 mEq per liter of blood. Severe hypokalemia can result in muscle weakness, twitches, paralysis, or abnormal heart rhythms. Administration of potassium supplements is generally sufficient to correct this disorder. Hyperkalemia, or high potassium blood level, results when kidneys don't excrete enough potassium or when a rapid influx of potassium is released from the reservoir in cells. Mild

hyperkalemia causes few, if any, symptoms. However, if potassium concentrations rise above 5.5 mEq per liter of blood, the heart's electrical conducting system is affected, resulting in abnormal heart rhythms and, possibly, cardiac arrest. Potassium can be removed from the body by dialysis, by induction of diarrhea, or by administering a potassium-absorbing resin. An intravenous solution of calcium, glucose, or insulin can also be given to protect the heart from the high levels of potassium and to drive the potassium from the blood into the cells, respectively.

[0026] The discovery of new human cardiovascular system associated polynucleotides, the polypeptides encoded by them, and antibodies that immunospecifically bind these polypeptides, satisfies a need in the art by providing new compositions which are useful in the diagnosis, treatment, prevention and/or prognosis of disorders of cardiovascular system, particularly disorders of the cardiovascular system, including, but not limited to, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease (e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease (e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or as described under "Cardiovascular Disorders" below.

### *Summary of the Invention*

[0027] The present invention relates to novel cardiovascular system related polynucleotides, the polypeptides encoded by these polynucleotides herein collectively referred to as "cardiovascular system antigens," and antibodies that immunospecifically bind these polypeptides, and the use of such cardiovascular system polynucleotides, antigens, and antibodies for detecting, treating, preventing and/or prognosing disorders of the cardiovascular system, including, but not limited to, the presence of cancer and cancer metastases. More specifically, isolated cardiovascular system nucleic acid molecules are provided encoding novel cardiovascular system polypeptides. Novel cardiovascular system polypeptides and antibodies that bind to these polypeptides are provided. Also provided are vectors, host cells, and recombinant and synthetic methods for producing human cardiovascular system polynucleotides, polypeptides, and/or antibodies. The



invention further relates to diagnostic and therapeutic methods useful for diagnosing, treating, preventing and/or prognosing disorders related to the cardiovascular system, including of cancer of the cardiovascular system tissues, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying agonists and antagonists of polynucleotides and polypeptides of the invention. The invention further relates to methods and/or compositions for inhibiting or promoting the production and/or function of the polypeptides of the invention.

### *Detailed Description*

#### Tables

[0028] Table 1A summarizes some of the polynucleotides encompassed by the invention (including cDNA clones related to the sequences (Clone ID NO:Z), contig sequences (contig identifier (Contig ID:) and contig nucleotide sequence identifier (SEQ ID NO:X)) and further summarizes certain characteristics of these polynucleotides and the polypeptides encoded thereby. The first column provides a unique clone identifier, "Clone ID NO:Z", for a cDNA plasmid related to each cardiovascular system associated contig sequence disclosed in Table 1A. The second column provides a unique contig identifier, "Contig ID:" for each of the contig sequences disclosed in Table 1A. The third column provides the sequence identifier, "SEQ ID NO:X", for each of the contig polynucleotide sequences disclosed in Table 1A. The fourth column, "ORF (From-To)", provides the location (i.e., nucleotide position numbers) within the polynucleotide sequence of SEQ ID NO:X that delineate the preferred open reading frame (ORF) shown in the sequence listing and referenced in Table 1A as SEQ ID NO:Y (column 5). Column 6 lists residues comprising predicted epitopes contained in the polypeptides encoded by each of the preferred ORFs (SEQ ID NO:Y). Identification of potential immunogenic regions was performed according to the method of Jameson and Wolf (CABIOS, 4:181-186 (1988)); specifically, the Genetics Computer Group (GCG) implementation of this algorithm, embodied in the program PEPTIDESTRUCTURE (Wisconsin Package v10.0, Genetics Computer Group (GCG), Madison, Wisc.). This method returns a measure of the probability that a given residue is found on the surface of the protein. Regions where the antigenic index score is greater than 0.9 over at least 6 amino acids are indicated in Table 1A as "Predicted Epitopes." In particular embodiments, cardiovascular system associated

polypeptides of the invention comprise, or alternatively consist of, one, two, three, four, five or more of the predicted epitopes described in Table 1A. It will be appreciated that depending on the analytical criteria used to predict antigenic determinants, the exact address of the determinant may vary slightly. Column 7, "Tissue Distribution" shows the expression profile of tissue, cells, and/or cell line libraries which express the polynucleotides of the invention. The first number in column 7 (preceding the colon), represents the tissue/cell source identifier code corresponding to the code and description provided in Table 4. Expression of these polynucleotides was not observed in the other tissues and/or cell libraries tested. For those identifier codes in which the first two letters are not "AR", the second number in column 7 (following the colon), represents the number of times a sequence corresponding to the reference polynucleotide sequence (e.g., SEQ ID NO:X) was identified in the tissue/cell source. Those tissue/cell source identifier codes in which the first two letters are "AR" designate information generated using DNA array technology. Utilizing this technology, cDNAs were amplified by PCR and then transferred, in duplicate, onto the array. Gene expression was assayed through hybridization of first strand cDNA probes to the DNA array. cDNA probes were generated from total RNA extracted from a variety of different tissues and cell lines. Probe synthesis was performed in the presence of  $^{33}\text{P}$  dCTP, using oligo(dT) to prime reverse transcription. After hybridization, high stringency washing conditions were employed to remove non-specific hybrids from the array. The remaining signal, emanating from each gene target, was measured using a Phosphorimager. Gene expression was reported as Phosphor Stimulating Luminescence (PSL) which reflects the level of phosphor signal generated from the probe hybridized to each of the gene targets represented on the array. A local background signal subtraction was performed before the total signal generated from each array was used to normalize gene expression between the different hybridizations. The value presented after "[array code]:" represents the mean of the duplicate values, following background subtraction and probe normalization. One of skill in the art could routinely use this information to identify normal and/or diseased tissue(s) which show a predominant expression pattern of the corresponding polynucleotide of the invention or to identify polynucleotides which show predominant and/or specific tissue and/or cell expression. Column 8, "Cytologic Band," provides the chromosomal location of

10091504.030702

polynucleotides corresponding to SEQ ID NO:X. Chromosomal location was determined by finding exact matches to EST and cDNA sequences contained in the NCBI (National Center for Biotechnology Information) UniGene database. Given a presumptive chromosomal location, disease locus association was determined by comparison with the Morbid Map, derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM™. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). If the putative chromosomal location of the Query overlapped with the chromosomal location of a Morbid Map entry, an OMIM identification number is provided in Table 1A, column 9 labeled "OMIM Disease Reference(s)". A key to the OMIM reference identification numbers is provided in Table 5.

[0029] Table 1B summarizes additional polynucleotides encompassed by the invention (including cDNA clones related to the sequences (Clone ID NO:Z), contig sequences (contig identifier (Contig ID:) contig nucleotide sequence identifiers (SEQ ID NO:X)), and genomic sequences (SEQ ID NO:B). The first column provides a unique clone identifier, "Clone ID NO:Z", for a cDNA clone related to each contig sequence. The second column provides the sequence identifier, "SEQ ID NO:X", for each contig sequence. The third column provides a unique contig identifier, "Contig ID:" for each contig sequence. The fourth column, provides a BAC identifier "BAC ID NO:A" for the BAC clone referenced in the corresponding row of the table. The fifth column provides the nucleotide sequence identifier, "SEQ ID NO:B" for a fragment of the BAC clone identified in column four of the corresponding row of the table. The sixth column, "Exon From-To", provides the location (i.e., nucleotide position numbers) within the polynucleotide sequence of SEQ ID NO:B which delineate certain polynucleotides of the invention that are also exemplary members of polynucleotide sequences that encode polypeptides of the invention (e.g., polypeptides containing amino acid sequences encoded by the polynucleotide sequences delineated in column six, and fragments and variants thereof).

[0030] Table 2 summarizes homology and features of some of the polypeptides of the invention. The first column provides a unique clone identifier, "Clone ID NO:Z", corresponding to a cDNA disclosed in Table 1A. The second column provides the

unique contig identifier, "Contig ID:" corresponding to contigs in Table 1A and allowing for correlation with the information in Table 1A. The third column provides the sequence identifier, "SEQ ID NO:X", for the contig polynucleotide sequences. The fourth column provides the analysis method by which the homology/identity disclosed in the row was determined. Comparisons were made between polypeptides encoded by the polynucleotides of the invention and either a non-redundant protein database (herein referred to as "NR"), or a database of protein families (herein referred to as "PFAM") as further described below. The fifth column provides a description of PFAM/NR hits having significant matches to a polypeptide of the invention. Column six provides the accession number of the PFAM/NR hit disclosed in the fifth column. Column seven, "Score/Percent Identity", provides a quality score or the percent identity, of the hit disclosed in column five. Columns 8 and 9, "NT From" and "NT To" respectively, delineate the polynucleotides in "SEQ ID NO:X" that encode a polypeptide having a significant match to the PFAM/NR database as disclosed in the fifth column. In specific embodiments, polypeptides of the invention comprise, or alternatively consist of, an amino acid sequence encoded by the polynucleotides in SEQ ID NO:X as delineated in columns 8 and 9, or fragments or variants thereof.

[0031] Table 3 provides polynucleotide sequences that may be disclaimed according to certain embodiments of the invention. The first column provides a unique clone identifier, "Clone ID NO:Z", for a cDNA clone related to cardiovascular system associated contig sequences disclosed in Table 1A. The second column provides the sequence identifier, "SEQ ID NO:X", for contig polynucleotide sequences disclosed in Table 1A. The third column provides the unique contig identifier, "Contig ID", for contigs disclosed in Table 1A. The fourth column provides a unique integer 'a' where 'a' is any integer between 1 and the final nucleotide minus 15 of SEQ ID NO:X, represented as "Range of a", and the fifth column provides a unique integer 'b' where 'b' is any integer between 15 and the final nucleotide of SEQ ID NO:X, represented as "Range of b", where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:X, and where b is greater than or equal to a + 14. For each of the polynucleotides shown as SEQ ID NO:X, the uniquely defined integers can be substituted into the general formula of a-b, and used to describe polynucleotides which may be preferably excluded from the invention. In certain embodiments,

preferably excluded from the polynucleotides of the invention (including polynucleotide fragments and variants as described herein and diagnostic and/or therapeutic uses based on these polynucleotides) are at least one, two, three, four, five, ten, or more of the polynucleotide sequence(s) having the accession number(s) disclosed in the sixth column of this Table (including for example, published sequence in connection with a particular BAC clone). In further embodiments, preferably excluded from the invention are the specific polynucleotide sequence(s) contained in the clones corresponding to at least one, two, three, four, five, ten, or more of the available material having the accession numbers identified in the sixth column of this Table (including for example, the actual sequence contained in an identified BAC clone).

**[0032]** Table 4 provides a key to the tissue/cell source identifier code disclosed in Table 1A, column 7. Column 1 provides the key to the tissue/cell source identifier code disclosed in Table 1A, Column 7. Columns 2-5 provide a description of the tissue or cell source. Codes corresponding to diseased tissues are indicated in column 6 with the word "disease". The use of the word "disease" in column 6 is non-limiting. The tissue or cell source may be specific (e.g. a neoplasm), or may be disease-associated (e.g., a tissue sample from a normal portion of a diseased organ). Furthermore, tissues and/or cells lacking the "disease" designation may still be derived from sources directly or indirectly involved in a disease state or disorder, and therefore may have a further utility in that disease state or disorder. In numerous cases where the tissue/cell source is a library, column 7 identifies the vector used to generate the library.

**[0033]** Table 5 provides a key to the OMIM<sup>TM</sup> reference identification numbers disclosed in Table 1A, column 9. OMIM reference identification numbers (Column 1) were derived from Online Mendelian Inheritance in Man (Online Mendelian Inheritance in Man, OMIM<sup>TM</sup>. McKusick-Nathans Institute for Genetic Medicine, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine, (Bethesda, MD) 2000. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>). Column 2 provides diseases associated with the cytologic band disclosed in Table 1A, column 8, as determined from the Morbid Map database.

- [0034] Table 6 summarizes ATCC Deposits, Deposit dates, and ATCC designation numbers of deposits made with the ATCC in connection with the present application.
- [0035] Table 7 shows the cDNA libraries sequenced, tissue source description, vector information and ATCC designation numbers relating to these cDNA libraries.
- [0036] Table 8 provides a physical characterization of clones encompassed by the invention. The first column provides the unique clone identifier, "Clone ID NO:Z", for certain cDNA clones of the invention, as described in Table 1A. The second column provides the size of the cDNA insert contained in the corresponding cDNA clone.

### Definitions

- [0037] The following definitions are provided to facilitate understanding of certain terms used throughout this specification.
- [0038] In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing features of the polynucleotide sequences of the present invention.
- [0039] As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence encoding SEQ ID NO:Y or a fragment or variant thereof, a nucleic acid sequence contained in SEQ ID NO:X (as described in column 3 of Table 1A) or the complement thereof, a cDNA sequence contained in Clone ID NO:Z (as described in column 1 of Table 1A and contained within a library deposited with the ATCC); a nucleotide sequence encoding the polypeptide encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B or a fragment or variant thereof; or a nucleotide coding sequence in SEQ ID NO:B as defined in column 6 of Table 1B or the complement thereof. For example, the polynucleotide can contain the nucleotide

sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having an amino acid sequence encoded by a polynucleotide of the invention as broadly defined (obviously excluding poly-Phenylalanine or poly-Lysine peptide sequences which result from translation of a polyA tail of a sequence corresponding to a cDNA).

[0040] As used herein, a "cardiovascular system antigen" refers collectively to any polynucleotide disclosed herein (e.g., a nucleic acid sequence contained in SEQ ID NO:X or the complement thereof, or cDNA sequence contained in Clone ID NO:Z, or a nucleotide sequence encoding the polypeptide encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B, or a nucleotide coding sequence in SEQ ID NO:B as defined in column 6 of Table 1B or the complement thereof and fragments or variants thereof as described herein) or any polypeptide disclosed herein (e.g., an amino acid sequence contained in SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X, or the complement thereof, an amino acid sequence encoded by the cDNA sequence contained in Clone ID NO:Z, an amino acid sequence encoded by SEQ ID NO:B, or the complement thereof, and fragments or variants thereof as described herein). These cardiovascular system antigens have been determined to be predominantly expressed in cardiovascular system tissues, including normal or diseased tissues (as shown in Table 1A column 7 and Table 4).

[0041] In the present invention, "SEQ ID NO:X" was often generated by overlapping sequences contained in multiple clones (contig analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X is deposited at Human Genome Sciences, Inc. (HGS) in a catalogued and archived library. As shown, for example, in column 1 of Table 1A, each clone is identified by a cDNA Clone ID (identifier generally referred to herein as Clone ID NO:Z). Each Clone ID is unique to an individual clone and the Clone ID is all the information needed to retrieve a given clone from the HGS library. Furthermore, certain clones disclosed in this application have been deposited with the ATCC on October 5, 2000, having the ATCC designation numbers PTA 2574 and PTA 2575; and on January 5, 2001, having the depositor reference numbers TS-1, TS-2, AC-1, and AC-2. In addition to the individual cDNA clone deposits, most of the cDNA libraries from which the clones

were derived were deposited at the American Type Culture Collection (hereinafter "ATCC"). Table 7 provides a list of the deposited cDNA libraries. One can use the Clone ID NO:Z to determine the library source by reference to Tables 6 and 7. Table 7 lists the deposited cDNA libraries by name and links each library to an ATCC Deposit. Library names contain four characters, for example, "HTWE." The name of a cDNA clone (Clone ID NO:Z) isolated from that library begins with the same four characters, for example "HTWEP07". As mentioned below, Table 1A correlates the Clone ID NO:Z names with SEQ ID NO:X. Thus, starting with an SEQ ID NO:X, one can use Tables 1A, 6 and 7 to determine the corresponding Clone ID NO:Z, which library it came from and which ATCC deposit the library is contained in. Furthermore, it is possible to retrieve a given cDNA clone from the source library by techniques known in the art and described elsewhere herein. The ATCC is located at 10801 University Boulevard, Manassas, Virginia 20110-2209, USA. The ATCC deposits were made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure.

[0042] In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5 kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

[0043] A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to sequences contained in SEQ ID NO:X, or the complement thereof (e.g., the complement of any one, two, three, four, or more of the polynucleotide fragments described herein), the polynucleotide sequence delineated in columns 8 and 9 of Table 2 or the complement thereof, and/or cDNA sequences contained in Clone ID NO:Z (e.g., the complement of any one, two, three, four, or more of the polynucleotide fragments, or the cDNA



clone within the pool of cDNA clones deposited with the ATCC, described herein) and/or the polynucleotide sequence delineated in column 6 of Table 1B or the complement thereof. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

[0044] Also contemplated are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency), salt conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH<sub>2</sub>PO<sub>4</sub>; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 µg/ml salmon sperm blocking DNA; followed by washes at 50 degree C with 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

[0045] Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

[0046] Of course, a polynucleotide which hybridizes only to polyA<sup>+</sup> sequences (such as any 3' terminal polyA<sup>+</sup> tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone generated using oligo dT as a primer).

10094504.030702

[0047] The polynucleotide of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

[0048] The polypeptide of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-

links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth. Enzymol. 182:626-646 (1990); Rattan et al., Ann. N.Y. Acad. Sci. 663:48-62 (1992).)

[0049] "SEQ ID NO:X" refers to a polynucleotide sequence described, for example, in Tables 1A or 2, while "SEQ ID NO:Y" refers to a polypeptide sequence described in column 5 of Table 1A. SEQ ID NO:X is identified by an integer specified in column 3 of Table 1A. The polypeptide sequence SEQ ID NO:Y is a translated open reading frame (ORF) encoded by polynucleotide SEQ ID NO:X. "Clone ID NO:Z" refers to a cDNA clone described in column 1 of Table 1A.

[0050] "A polypeptide having biological activity" refers to a polypeptide exhibiting activity similar to, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention).

[0051] Table 1A summarizes some of the polynucleotides encompassed by the invention (including contig sequences (SEQ ID NO:X) and clones (Clone ID NO:Z) and further summarizes certain characteristics of these polynucleotides and the polypeptides encoded thereby.

#### **Polynucleotides and Polypeptides**

TABLE 1A

Clone ID NO: Z	Contig ID:	SEQ ID NO: X	ORF (From-To)	AA SEQ ID NO: Y	Predicted Epitopes	Tissue Distribution Library code: count (see Table IV for Library Codes)	Cytologic Band	OMIM Disease Reference(s):
HAHCL07	952661	11	489 - 136	626	Phe-16 to Cys-24, Thr-49 to Asp-60.	L0471: 3, L0759: 2, L0459: 1, L0622: 1, H0599: 1, H0196: 1, L0662: 1, L0747: 1 and L0604: 1.		
HAHCP26	681258	12	2 - 403	627	Val-18 to Glu-26, Val-43 to Gln-48, Pro-56 to Asp-63.	H0599: 2		
HAHCP41	712086	13	81 - 239	628	Gly-13 to Glu-18.	H0599: 2		
HAHCP49	722659	14	1 - 234	629	Tyr-24 to His-32, Asp-43 to Thr-48, Ala-54 to Lys-59.	H0599: 2		
HAHCP55	865095	15	62 - 250	630		H0599: 2		
HAHCP67	756916	16	2 - 262	631		H0599: 2	3p21.1	150250, 164500, 168468, 182280, 238310, 600163, 601226, 601916

HAHCP91	789998	17	163 - 288	632	Ser-29 to Ala-38.	H0599: 2		
HAHCR57	865096	18	1 - 387	633	Ala-6 to Ser-17, Ser-19 to Trp-24.	H0599: 2		
HAHEE04	922257	19	2 - 346	634		H0599: 2		
HAHEE05	928673	20	3 - 422	635	Asp-1 to Pro-10, Thr-13 to Asn-20, His-28 to Gln-45, Phe-56 to Ser-61, Glu-63 to Pro-70, Ser-98 to Ser-110, Thr-115 to Ser-122, Arg-130 to Gln-140.	H0599: 2 and L0777: 1.		
HAHEO46	718772	21	85 - 222	636	Arg-9 to Phe-14, Val-28 to Gln-35.	H0599: 3		
HAHES10	961594	22	2 - 241	637	Gly-1 to Asn-14, Val-20 to Ser-25, Thr-46 to Gly-54.	H0599: 2		
HAHFX20	925753	23	53 - 139	638	Glu-1 to Ala-7.	H0599: 2		
HAHHE12	969107	24	88 - 804	639		H0373: 5, L0604: 4, H0599: 2 and L0485: 1.		
HAHHO04	925814	25	41 - 313	640	Asp-18 to Asp-24, Pro-47 to Ser-53.	H0599: 2		
HAHHS01	913863	26	44 - 181	641	Ser-18 to Leu-28.	H0599: 2		
HAHIK10	961425	27	1 - 171	642	Glu-31 to Leu-47.	H0599: 2		
HAHIW08	955803	28	16 - 312	643	Tyr-4 to His-14,	H0599: 2		

HAHSC42	695111	29	3 - 440	644	Thr-32 to Ala-38, Pro-46 to Pro-51.			
HCMB95	508774	30	138 - 293	645	Pro-7 to Gly-21.			H0097: 1 and H0373: 1.
HCMS52	522615	31	32 - 88	646	Ile-1 to Ala-12.			H0173: 1 and H0196: 1.
HCMS09	530511	32	119 - 358	647	Glu-12 to Lys-17.			H0196: 2
HCMSQ77	862373	33	1 - 216	648	Phe-10 to His-18.			H0196: 2
HCMS72	526183	34	1 - 174	649	Arg-1 to Pro-7.			H0196: 2
HCMSU37	706337	35	1 - 312	650	Ser-1 to Leu-6, Lys-16 to Arg-23, Gln-26 to Asn-42, Ser-54 to Pro-66.			H0599: 1, H0196: 1 and L0485: 1.
HCMSX59	522598	36	3 - 173	651				H0230: 1 and H0196: 1.
HELA85	507230	37	63 - 266	652	Cys-1 to Gly-7.			S0045: 2
HELAI26	920922	38	79 - 216	653				S0045: 2
HELAL34	527994	39	1 - 210	654	Asn-3 to Gly-29.			S0045: 2
HELA61	507205	40	87 - 305	655	Phe-2 to Thr-12, His-49 to Arg-62.			S0045: 2
HELB08	960106	41	69 - 203	656				S0045: 2
HELB07	527677	42	5 - 166	657	Pro-4 to Cys-10.			S0045: 2
HELBK27	527669	43	35 - 163	658				S0045: 2
HELB45	527666	44	1 - 195	659	Tyr-7 to Tyr-12,			S0045: 2

Variable	Mean	SD	Min	Max
Age	34.5	10.2	21	55
Gender	Male	Female		
Marital Status	Married	Single		
Education	High School	College		
Occupation	Manager	Worker		
Income	\$30,000	\$40,000		
Health Status	Good	Fair		
Stress Level	Low	High		
Life Satisfaction	High	Low		
Work-Life Balance	Good	Poor		
Family Support	Strong	Weak		
Community Involvement	Active	Passive		
Personal Growth	High	Low		
Relationship Quality	Good	Poor		
Financial Stability	Stable	Unstable		
Emotional Well-being	High	Low		
Physical Well-being	Good	Poor		
Social Support	Strong	Weak		
Work Engagement	High	Low		
Job Satisfaction	High	Low		
Life Balance	Good	Poor		
Overall Quality of Life	High	Low		

[illegible]

HELFG79	577248	63	126 - 371	678	Asp-22 to Gly-27, Ala-36 to Leu-43, Phe-58 to Trp-73.	S0045: 3		
HELGA54	576374	64	12 - 440	679	Pro-7 to Trp-15, Leu-17 to Cys-32.	S0045: 2		
HELGC24	576377	65	56 - 247	680	Arg-15 to Arg-24, Ser-39 to Val-47.	S0045: 2		
HELGC32	699375	66	2 - 238	681	Pro-10 to Ser-17, Ala-21 to Thr-26, Pro-60 to Leu-72.	L0455: 2, S0045: 1 and H0050: 1.		
HELGC77	577317	67	72 - 242	682	Ser-2 to Asp-14.	S0045: 2		
HELGG77	825795	68	312 - 527	683	Lys-27 to Asn-45, Pro-62 to His-68.	S0045: 1 and H0413: 1.		
HELGH89	545009	69	338 - 3	684	Arg-41 to Tyr-46, Ile-92 to Asn-97.	AR050: 1, AR054: 1, AR051: 0 S0045: 2		
	553652	606	338 - 3	1221	Arg-41 to Tyr-46, Ile-92 to Asn-97.			
HELGN53	963160	70	180 - 368	685	Arg-1 to Gly-6.	S0045: 1 and H0619: 1.		
HELGP50	576339	71	27 - 197	686	Arg-1 to Gly-7.	S0045: 2		
HELGQ48	851178	72	97 - 222	687		S0045: 2		
HELGT48	879483	73	312 - 647	688	Arg-41 to Tyr-46, Ile-92 to Asn-97.	S0045: 2		
HELGZ48	721742	74	1 - 348	689	Pro-20 to Thr-26, Gln-62 to Thr-68.	S0045: 1 and S0046: 1.		



HELHB12	970863	75	86 - 184	690	Val-75 to Gln-83.				
HELHC49	576292	76	40 - 228	691			S0045: 1 and S0046: 1.		
HELHD46	719129	77	1 - 279	692			S0045: 2		
HELHF07	949067	78	65 - 340	693	Thr-16 to Pro-21.		S0045: 1 and S0046: 1.		
HELHM29	883505	79	1 - 303	694			AR061: 5, AR089: 4 S0045: 2		
HEMBC56	577797	80	148 - 381	695	Gly-33 to Ser-38.		S0045: 2		
HEMBI16	507220	81	23 - 169	696			S0046: 2		
HEMBZ84	527989	82	33 - 125	697			S0046: 2		
HEMCA89	527985	83	9 - 131	698			S0046: 2		
HEMCM25	948738	84	84 - 212	699			AR051: 22, AR054: 20, AR050: 17 S0046: 4		
HEMDG56	715834	85	1 - 120	700			S0046: 2		
HEMDG83	576508	86	231 - 452	701			S0046: 4 and L0754: 1.		
HEMDK92	574278	87	124 - 303	702			S0046: 2, L0748: 1 and L0755: 1.		
HEMDM56	577251	88	94 - 252	703			S0045: 1 and S0046: 1.		
HEMDO24	577297	89	101 - 391	704	Asp-10 to Gln-20, Trp-27 to Ser-38, Glu-54 to Lys-69.		S0046: 2		
HEMEA72	527804	90	2 - 184	705	Lys-16 to Phe-21.		S0046: 1 and H0268: 1.		
HEMEH76	574345	91	2 - 211	706	Pro-6 to Ser-16, Ala-33 to Ser-38.		S0046: 2		

HEMEK19	574209	92	78 - 263	707	Phe-48 to Gly-57.	AR089: 9, AR061: 3 S0046: 2		
HEMEN63	578717	93	1 - 321	708	Gly-1 to Lys-6, Arg-16 to Gly-27, Pro-47 to Gly-67.	S0046: 3		
HEMEU54	947801	94	3 - 371	709	Ala-10 to Arg-17, Pro-74 to Val-79, Gly-85 to Tyr-92, Pro-94 to Lys-100.	AR054: 126, AR050: 108, AR051: 103 S0046: 1 and H0619: 1.		
HEMFL58	576505	95	3 - 221	710	Gln-25 to Leu-30.	S0046: 2		
HEMFN33	702564	96	2 - 169	711	Ala-4 to Ser-10.	S0046: 2		
HEMFX20	840164	97	41 - 160	712	Glu-1 to Lys-7.	S0045: 1 and S0046: 1.		
HEMGL57	971118	98	3 - 182	713	Pro-13 to Arg-19.	S0046: 2		
HEMGL58	578091	99	1 - 195	714		S0046: 1, H0050: 1 and 3 L0743: 1.		
HHBBA47	720472	100	30 - 179	715	Arg-9 to Arg-22.	H0373: 2		
HHBB111	959756	101	1 - 201	716	Cys-11 to Thr-20, Ala-22 to Thr-30.	H0373: 2		
HHBBK65	588062	102	242 - 343	717		H0373: 2		
HHBBL40	588066	103	98 - 274	718	Asn-43 to Gly-51.	H0373: 2		
HHBBL53	588067	104	2 - 199	719	Pro-1 to Leu-15, Lys-18 to Ala-25, Glu-32 to Asp-49, Ser-55 to Thr-66.	H0373: 2		
HHBEA32	927399	105	1 - 159	720	Pro-8 to Gly-17.	H0373: 2		

HHBEE70	697541	106	269 - 424	721			H0373: 2	
HHBEM70	756949	107	45 - 242	722			AR089: 3, AR061: 1 H0373: 3	
HHBEN34	703745	108	278 - 382	723	Ser-11 to Lys-21.		H0373: 2	
HHBFL31	800035	109	41 - 223	724			AR050: 51, AR054: 43, AR051: 35 H0373: 2	
HHBFW44	716283	110	49 - 117	725			H0373: 2	
HHBFW75	958692	111	100 - 423	726	Thr-24 to Lys-30, Gln-33 to Glu-41.		H0373: 2	
HHBGN42	698774	112	124 - 2	727	Arg-12 to Asn-19.		H0373: 2	
HHBGN52	726391	113	2 - 229	728	Pro-1 to Lys-6.		H0373: 3	
HHBGN68	752745	114	3 - 53	729	Glu-10 to Gly-17.		H0373: 2	
HHBGR37	708457	115	139 - 306	730			H0373: 2	
HHBGT39	940578	116	2 - 67	731	Lys-11 to Gly-18.		H0373: 2	
HHBGY59	792027	117	3 - 452	732			H0373: 2	
HHBHE83	780875	118	2 - 76	733	Ala-3 to Arg-8.		H0373: 3	
HHBHK08	958649	119	146 - 301	734	Gly-16 to Pro-24.		H0373: 2	
HHBHK84	858431	120	3 - 350	735	Leu-16 to Gly-26, Pro-67 to Arg-78.		H0373: 3	
HHBHO63	906947	121	179 - 310	736			H0373: 2	
HHBHP27	676601	122	79 - 414	737	Ile-12 to Gln-30, Tyr-57 to Ser-64.		H0373: 2	10p12
HHFBD39	826307	123	1 - 231	738	Cys-51 to Arg-60.		H0050: 2	
HHFBD50	724763	124	301 - 146	739	Asp-36 to Pro-48.		H0619: 1, H0050: 1,	

							L0748: 1 and L0759: 1.		
	858082	607	209 - 3	1222	Gln-1 to Thr-35.				
HHFBF32	502872	125	69 - 179	740	Glu-1 to Glu-8.		H0619: 1, H0050: 1 and H0105: 1.		
HHFBH26	502843	126	46 - 213	741	Cys-34 to Ser-48.		H0050: 3		
HHFBI05	932961	127	3 - 188	742	Gly-2 to Ser-9.		H0050: 3		
HHFBJ81	502954	128	56 - 196	743			H0050: 2		
HHFBL16	509237	129	147 - 362	744	Gln-30 to Gly-49, Pro-52 to Arg-63.		H0050: 2		
HHFBL30	509238	130	1 - 222	745	Arg-18 to Pro-37.		H0050: 2		
HHFBL32	509233	131	82 - 228	746	Ser-1 to Phe-12, Gly-37 to Lys-49.		H0050: 2		
HHFBL36	707930	132	169 - 249	747			H0050: 3		
HHFBL39	509637	133	3 - 185	748	Cys-1 to Arg-8, Pro-13 to Tyr-22.		H0050: 2		
HHFBL60	739670	134	115 - 270	749			H0050: 2		
HHFBP60	503453	135	1 - 183	750	Phe-31 to Tyr-51.		H0050: 2		
HHFBX77	959805	136	1 - 369	751	Thr-19 to His-25, Glu-58 to Lys-66, Ala-74 to Glu-84, Asn-116 to Glu-123.		H0050: 4 and H0619: 1.		
HHFCA64	720849	137	287 - 207	752			H0050: 3 and H0619: 1.		
	926760	608	69 - 260	1223					
HHFCF70	518435	138	2 - 379	753	Glu-4 to Ser-16,		H0050: 2		

HHFCH59	526389	139	174 - 350	754	Pro-41 to Gln-48, Lys-55 to Ser-63, Lys-109 to Leu-115.	H0050: 2		
HHFCI73	518427	140	150 - 1	755	Lys-6 to Leu-12, Arg-24 to Gly-29, Pro-32 to Cys-38, Thr-42 to Asn-50.	H0050: 2		
HHFCK71	781725	141	98 - 214	756		H0050: 2		
HHFCL91	509628	142	3 - 221	757	Asn-32 to Val-41, Pro-52 to Ala-65.	H0050: 2		
HHFCP67	536062	143	110 - 295	758		H0050: 2		
HHFCZ01	509163	144	165 - 287	759		H0050: 2		
HHFDA67	509387	145	16 - 294	760		H0050: 2 and H0645: 1.		
HHFDG32	502957	146	118 - 300	761	Ser-4 to Ser-9, Phe-13 to Tyr-19.	H0050: 2		
HHFDH02	921297	147	1 - 168	762	Asn-1 to Asn-15.	H0050: 2		
HHFDH38	536551	148	3 - 293	763	Thr-3 to Glu-9.	H0050: 2		
HHFDI07	954404	149	47 - 214	764	Pro-24 to Asn-29.	H0050: 2		
HHFDI42	500877	150	3 - 152	765	Arg-11 to Gly-16.	H0050: 2		
HHFDI62	745569	151	169 - 309	766		H0050: 2		
HHFDI66	573283	152	210 - 401	767		H0050: 2		
HHFDJ27	534909	153	166 - 378	768		H0242: 4 and H0050: 1.		

HHFDJ87	575105	154	60 - 293	769	Val-13 to Arg-25, Ile-51 to Pro-56.	H0050: 2		
HHFDM05	932863	155	218 - 427	770	Thr-10 to Glu-27.	H0050: 2		
HHFEJ18	525616	156	30 - 134	771	Gly-29 to Glu-35.	H0619: 1, H0050: 1 and H0242: 1.		
HHFEO24	525611	157	132 - 308	772	Asn-10 to Ser-16.	H0050: 2		
HHFES51	525609	158	1 - 180	773	Asn-1 to Gly-7, Lys-15 to Ser-20.	H0050: 2		
HHFFG41	575009	159	2 - 121	774		H0050: 2		
HHFFO64	523769	160	191 - 316	775		H0050: 2		
HHFFO66	500910	161	77 - 217	776	Ser-31 to Gly-41.	H0050: 3		
HHFFO96	894076	162	186 - 76	777	Ser-1 to Ile-12.	H0619: 1 and H0050: 1.		
HHFFI69	662462	163	230 - 511	778	Ser-72 to Val-81.	L0779: 2, L0731: 2, H0002: 1, H0050: 1, L0769: 1, L0637: 1, L0809: 1 and L0745: 1.		
HHFFX20	525617	164	45 - 116	779		H0050: 2		
HHFFX75	507394	165	115 - 261	780	Gln-28 to Gln-34.	H0050: 1 and H0242: 1.		
HHFFY80	732614	166	124 - 399	781	Asn-1 to Asn-9.	H0050: 2		
HHFFZ04	927869	167	269 - 514	782	Asp-1 to Glu-6.	L0777: 3, H0050: 2 and H0619: 1.	I4q24-q32	107970, 115650, 123270, 182600,

HHFFZ19	509630	168	320 - 466	783	Thr-25 to Gln-31.	H0050: 2 and H0266: 1.	245200, 251600, 270100, 276900, 602091
HHFGA21	573584	169	114 - 299	784		H0050: 2	
HHFGC14	741650	170	38 - 370	785	Gln-1 to Asn-10.	AR089: 9, AR061: 5 H0050: 2	
HHFGC69	573491	171	81 - 275	786		H0050: 2	
HHFGC93	576487	172	90 - 311	787		H0050: 2	
HHFGC95	795968	173	24 - 389	788	Asn-34 to Glu-39.	H0050: 2	
HHFGE01	917137	174	2 - 406	789		H0050: 1 and H0105: 1.	
HHFGJ85	524890	175	183 - 293	790		H0050: 2	
HHFGM50	506635	176	111 - 311	791	Gln-10 to Val-15, Arg-21 to Gly-29.	H0050: 2	
HHFGN31	908508	177	93 - 230	792		AR089: 43, AR061: 5 H0619: 1 and H0050: 1.	
HHFGN67	573509	178	68 - 184	793		H0050: 2	
HHFGP71	573860	179	120 - 263	794	Val-18 to Glu-25.	H0050: 3 and H0619: 1.	
HHFGR35	573510	180	125 - 229	795		H0050: 2 and H0619:	

									1.			
HHFGS09	526331	181	3 - 257	796	Leu-43 to Leu-53.	H0050: 2						
HHFGS40	888332	182	35 - 313	797	Val-1 to His-9, Gly-43 to Leu-49.	AR051: 2, AR054: 2, AR050: 2 H0050: 3						
HHFGS92	871899	183	54 - 332	798		H0050: 2						
HHFGT10	968109	184	172 - 411	799		H0619: 1, H0645: 1 and H0050: 1.						
HHFGY13	573473	185	1 - 234	800	Gly-1 to Gly-9, Gly-24 to Arg-33, Thr-42 to Asp-47, Asn-63 to Lys-68.	H0050: 2						
HHFGY37	711364	186	133 - 414	801	Gly-3 to Trp-25.	H0050: 2						
HHFGY75	573483	187	196 - 450	802		H0050: 2						
HHFGZ54	573477	188	33 - 143	803	Pro-1 to Gly-9.	H0050: 2						
HHFGZ63	661248	189	132 - 281	804	Glu-27 to Lys-38.	H0050: 2						
	744984	609	470 - 228	1224								
HHFGZ69	918322	190	226 - 417	805	Gly-7 to Pro-12.	H0619: 2 and H0050: 1.						
HHFHA44	573498	191	138 - 275	806	Pro-31 to Lys-40.	H0050: 2						
HHFHC02	920510	192	1 - 216	807		H0050: 2						
HHFHC44	716763	193	3 - 227	808		H0050: 2 and L0748: 1.						
HHFHC57	573506	194	3 - 161	809		H0050: 2						
HHFHC72	766128	195	152 - 265	810	Arg-8 to Gly-25.	H0050: 2						



HHFHE28	506630	196	139 - 327	811			H0050: 2		
HHFHE58	526391	197	316 - 525	812			H0050: 2 and L0748: 1.		
HHFHE76	506578	198	281 - 436	813			H0050: 2 and H0619: 1.		
HHFHF91	526396	199	1 - 366	814	Gly-13 to Lys-32, Lys-47 to Glu-54, Thr-90 to Arg-98.		H0050: 2		
HHFHJ46	576949	200	1 - 513	815			S0045: 1, H0050: 1 and L0749: 1.		
HHFHJ72	575030	201	1 - 78	816			H0050: 1 and H0242: 1.		
HHFHM77	934027	202	2 - 187	817			H0619: 1 and H0050: 1.		
HHFHN54	506624	203	1 - 225	818	Ser-12 to Leu-17.		H0050: 2		
HHFHQ86	572923	204	231 - 353	819			H0050: 2		
HHFHU63	572871	205	52 - 300	820	Ser-40 to Lys-45.		H0050: 2		
HHFHX11	967321	206	3 - 83	821			H0050: 2		
HHFHY47	720473	207	151 - 351	822			H0050: 2 and H0242: 1.		
HHFHY66	573294	208	155 - 319	823			H0050: 2		
HHFJC02	918318	209	78 - 296	824	Arg-1 to Gly-10, Pro-44 to Trp-49, Ser-58 to Val-64.		H0619: 2		
HHFJI11	883731	210	2 - 298	825	Leu-2 to Gln-11.		H0619: 2		

HHFJL11	965931	211	56 - 202	826			H0619: 3		
HHFJM56	857990	212	26 - 268	827			H0619: 2		
HHFJN01	913798	213	1 - 312	828	Asn-28 to Gln-33.		H0619: 1 and H0050: 1.		
HHFJN02	918358	214	23 - 226	829			H0619: 1 and H0050: 1.		
HHFJO06	934168	215	271 - 453	830			H0619: 2 and H0050: 1.		
HHFJO12	969624	216	46 - 162	831			H0619: 1 and H0645: 1.		
HHFJR06	934093	217	25 - 243	832	Gly-66 to Asp-73.		H0619: 2		
HHFJX18	907658	218	154 - 342	833	Pro-5 to Lys-15.		H0619: 2		
HHFKB03	922803	219	58 - 318	834	Gly-66 to Gly-71, Ser-79 to Pro-87.		H0619: 2		
HHFKC28	857948	220	44 - 271	835			H0619: 1 and H0050: 1.		
HHFKC43	906903	221	171 - 404	836	Cys-26 to Thr-31.		H0619: 3, L0749: 3 and L0748: 1.		
HHFKE05	920550	222	335 - 472	837	Gln-4 to Trp-12.		H0619: 1 and H0050: 1.		
HHFKH82	857810	223	91 - 336	838			H0619: 1 and H0645: 1.		
HHFKI02	918288	224	83 - 268	839			H0619: 3		
HHFKJ68	934045	225	22 - 321	840	Pro-3 to Thr-9, Cys-34 to Leu-41.		H0619: 3		

HHFKK95	952082	226	340 - 510	841	Pro-49 to Arg-56.	H0619: 3		
HHFKM10	963184	227	55 - 324	842	Ser-15 to Arg-20, Gly-31 to Ala-47, Arg-69 to Gly-77.	H0619: 3		
HHFKU12	887244	228	2 - 229	843	Arg-16 to Gly-21, Gly-33 to Met-39, Glu-71 to His-76.	L0748: 3 and H0619: 2.		
HHFKU44	857923	229	239 - 529	844	Glu-15 to Gly-24.	H0619: 3		
HHFKX11	965874	230	155 - 331	845	Thr-53 to Phe-58.	H0619: 2		
HHFKY11	965876	231	121 - 315	846	Ser-10 to Trp-18, Arg-20 to Cys-27.	H0619: 2		
HHFLA69	933869	232	47 - 211	847		H0619: 1 and H0645: 1.		
HHFLH34	857907	233	2 - 334	848	Glu-45 to Gln-52, Asp-78 to Leu-85.	H0619: 1 and H0645: 1.		
HHFLM06	933868	234	147 - 359	849	Asn-37 to Gln-46, Asp-50 to Gly-65.	H0619: 1 and H0645: 1.		
HHFLP28	857888	235	1 - 204	850	His-8 to Ser-19.	H0619: 2		
HHFLU04	870085	236	127 - 366	851	Asn-1 to Glu-7, Glu-13 to Cys-18.	H0619: 2 and H0050: 1.		
HHFMA58	853959	237	2 - 202	852		H0619: 2		
HHFMB01	914951	238	3 - 179	853	Asp-1 to Glu-7, Leu-32 to Trp-37.	H0619: 2		
HHFME70	926497	239	128 - 265	854	Gln-6 to Arg-22.	H0619: 2		

HHFMMH85	872838	240	2 - 256	855	Pro-5 to Pro-12.	H0619: 2		
HHFMI58	857856	241	1 - 372	856	Thr-1 to Gly-14, Pro-40 to His-48, Thr-50 to Gly-72, His-74 to Glu-87.	H0619: 2 and L0534: 1.		
HHFMO03	922708	242	319 - 426	857	Arg-29 to Lys-36.	H0619: 3, L0766: 3, H0645: 1, H0050: 1 and L0439: 1.		
HHFMO94	957955	243	52 - 204	858	Leu-11 to Cys-17.	H0645: 3 and H0619: 1.		
HHFMQ70	840039	244	97 - 177	859	Asn-16 to Pro-27.	H0619: 2		
HHFMZ40	974299	245	269 - 475	860	Gln-44 to Pro-49, Asn-51 to Ser-57.	H0619: 2		
HHFNA49	914838	246	120 - 263	861		H0619: 2		
HHFND10	963140	247	188 - 451	862	Leu-16 to Leu-32.	H0619: 2		
HHFNF41	857825	248	1 - 357	863		H0619: 2		
HHFNI49	857850	249	1 - 192	864	Ser-17 to Ser-24.	H0619: 2 and H0050: 1.		
HHFOC27	857816	250	216 - 404	865		H0645: 1 and H0050: 1.		
HHFOC66	857955	251	1 - 240	866	Pro-19 to His-27, Pro-41 to Gln-54.	H0619: 1 and H0645: 1.	11q13	102200, 106100, 131100, 131100, 131100,

									133780, 147050, 153700, 161015, 164009, 168461, 168461, 168461, 180721, 180840, 191181, 193235, 209901, 232600, 259700, 259770, 600045, 600319, 600528, 601884
HHFOF07	952062	252	83 - 160	867				H0645: 1 and H0050: 1.	
HHFOF40	930839	253	106 - 327	868				H0645: 3	
HHFOJ05	930856	254	93 - 239	869	Cys-5 to Val-11.			H0645: 1 and H0050: 1.	

HHFON08	957974	255	2 - 154	870			H0645: 2		
HHFON15	857792	256	68 - 259	871			H0645: 2		
HHFON19	910891	257	250 - 825	872		Thr-136 to Cys-141.	AR089: 19, AR061: 17 H0645: 2 and H0050: 1.		
HHFON32	858034	258	111 - 398	873		Lys-39 to Ser-49.	H0645: 3, H0050: 1 and H0105: 1.		
HHFON87	857783	259	1 - 273	874		Pro-8 to Arg-15.	H0645: 2 and L0529: 1.		
HHFOW08	857772	260	351 - 202	875			H0645: 1 and H0050: 1.		
HHFSB02	921364	261	3 - 191	876		Pro-43 to Tyr-49.	H0050: 2 and H0105: 1.		
HHFTC88	500903	262	357 - 190	877			H0050: 1 and H0233: 1.		
HHFUB15	534520	263	81 - 314	878		Ala-32 to Asp-39.	H0242: 4		
HHFUB23	928063	264	101 - 250	879		Leu-3 to Phe-11.	H0242: 2		
HHFUB77	772691	265	76 - 195	880			H0242: 2		
HHFUC24	524840	266	81 - 317	881			H0242: 2		
HHFUC26	960331	267	136 - 381	882		Lys-48 to Ser-53.	H0242: 2 and H0645: 1.		
HHFUC42	525603	268	221 - 325	883			H0242: 2, L0789: 2, L0805: 1, L0776: 1 and L0530: 1.		
HHFUC45	525600	269	1 - 162	884		Thr-1 to Gly-9.	H0242: 2		

HHFUC47	525599	270	194 - 75	885			H0242: 2		
HHFUC83	507459	271	1 - 174	886	Arg-13 to Cys-20.		H0242: 2		
HHFUC92	507018	272	48 - 245	887			H0242: 2 and H0050: 1.		
HHFUK58	525604	273	43 - 168	888	Gly-2 to Cys-8, Pro-17 to Lys-27.		H0242: 2		
HHFUL75	675569	274	1 - 237	889	Thr-4 to Ala-9, His-40 to Ala-51, Glu-53 to Leu-58.		H0050: 1, H0242: 1 and H0373: 1.		
HHFUN56	530953	275	1 - 222	890			H0050: 1 and H0242: 1.		
HMEBA75	530303	276	1 - 159	891	Arg-1 to Gln-10.		S0045: 1 and H0267: 1.		
HMEBY61	947868	277	29 - 193	892			T0049: 1 and H0267: 1.		
HMEDF58	738427	278	284 - 493	893			S0045: 1 and H0266: 1.		
HMEDR48	529173	279	3 - 134	894			H0266: 2		
HMEEF64	918746	280	2 - 229	895	Ser-8 to Arg-13, Gln-19 to Gln-30.		S0045: 1 and H0266: 1.		
HMEEX61	742085	281	44 - 361	896	Lys-61 to Leu-68.		H0050: 1, H0266: 1 and L0362: 1.		
HMEFR17	664438	282	3 - 473	897	Asp-11 to Thr-35.		H0266: 2 and L0742: 2.		
HMEFX12	661951	283	16 - 186	898	Pro-1 to Pro-11, Ala-15 to Pro-21, Pro-29 to Asp-34.		H0266: 2		

HMEGB93	573814	284	3 - 365	899	His-1 to Asn-10, Ile-25 to Asn-38, Gly-60 to Ile-66.	H0266: 2	1p35-p34	118210, 120550, 120570, 120575, 121800, 130500, 133200, 138140, 138971, 168360, 171760, 171760, 172411, 176100, 176100, 178300, 185470, 230000, 230350, 255800, 602771
HMEGF48	573821	285	74 - 256	900		H0266: 2		
HMEGG44	796421	286	15 - 176	901	Cys-12 to Trp-18.	H0266: 2 and L0740: 1.		
HMEGH46	887791	287	1 - 315	902	Asp-13 to Asp-19,	AR054: 29, AR051:		



						Lys-76 to Asn-83.	12, AR061: 6, AR089: 3, AR050: 2 H0196: 1 and H0266: 1.		
HMEGI07	953815	288	9 - 272	903			H0266: 2, S0046: 1, L0776: 1 and L0754: 1.		
HMEIG42	931114	289	35 - 406	904		Ala-8 to Arg-17, Val-43 to Gln-56, Ser-73 to Met-89.	H0266: 2		
HMEIM40	523589	290	61 - 219	905		Lys-5 to Ser-14.	H0266: 2		
HMEIU49	722988	291	99 - 284	906			H0266: 2		
HMEIW23	682863	292	1 - 276	907		Gly-36 to Ser-45, Pro-87 to Ser-92.	H0266: 2		
HMEJD13	657231	293	2 - 271	908		Ala-1 to Pro-11.	S0046: 1 and H0266: 1.		
HMEJ84	781983	294	8 - 136	909		Ile-1 to Gly-13, Asn-18 to Ser-29.	H0266: 2		
HMEJU60	740392	295	33 - 191	910		Glu-21 to Leu-29.	H0266: 2		
HMEKA53	711664	296	76 - 183	911			H0266: 3		
HMEKJ43	715893	297	136 - 381	912		Leu-1 to Leu-13, Gln-55 to Pro-60.	H0266: 2 and L0744: 1.		
HMEKS76	767517	298	3 - 221	913		His-1 to Arg-9, Arg-23 to Ser-32, Gln-39 to Lys-46, Gly-53 to Gly-58, Glu-64 to Asn-73.	H0266: 2		

HMEKX51	727154	299	198 - 314	914			H0266: 2		
HMEKX89	786055	300	1 - 75	915	Ala-7 to Arg-15.		S0045: 1 and H0266: 1.		
HMELC56	745773	301	2 - 319	916	Ala-1 to Lys-6, Lys-75 to His-84.		H0266: 2		
HMEQ62	719681	302	40 - 372	917			H0266: 2		
HMELR10	964629	303	208 - 336	918			H0266: 2		
HMELS59	720341	304	135 - 305	919			H0266: 2		
HMELV19	668665	305	13 - 279	920	Ser-1 to Gly-8, Asn-39 to Ala-45.		H0266: 2		
HULAF89	791261	306	1 - 99	921			H0530: 2		
HULAI37	708923	307	72 - 173	922	Leu-9 to Lys-34.		AR061: 0, AR089: 0 H0530: 6		
HULAX31	868930	308	1 - 219	923			H0530: 2		
HULBU59	636253	309	1 - 234	924			H0530: 2		
HULBY15	659557	310	211 - 387	925	Glu-4 to Gly-10, Leu-47 to Lys-54.		H0530: 3		
HULDF69	754381	311	123 - 257	926	Lys-14 to Lys-20, Lys-25 to Gly-42.		H0530: 2		
HULFA03	918691	312	92 - 379	927	Leu-36 to Phe-55, Lys-61 to Phe-68.		H0530: 2		
HULFB76	767873	313	37 - 150	928			AR089: 1, AR061: 1 H0530: 3		
HUMBE61	838469	314	1 - 414	929			H0531: 2 and H0530: 2.		
HUSAY26	527909	315	263 - 373	930			H0268: 2 and L0666:		

								I.				
HUSCA57	527800	316	142 - 264	931				H0268: 2				
HUSCD26	527799	317	127 - 255	932				H0268: 2				
HUSCH45	507197	318	121 - 273	933				H0268: 2				
HUSCH77	527801	319	1 - 129	934				H0268: 2				
HUSGH88	871776	320	41 - 298	935		Asn-30 to Trp-36, Ser-63 to Ser-69.		L0740: 3, H0413: 2, L0777: 2, T0048: 1, H0412: 1 and L0757: 1.				
HUSGK23	675951	321	102 - 239	936				S0045: 1, H0268: 1 and I5q15 H0412: 1.			177070, 177070, 182500, 218000, 227220, 243500, 600839, 601800	
HUSGL13	575798	322	2 - 169	937		Ala-3 to Ser-10, Arg-24 to Trp-33.		H0412: 2				
HUSHH64	576459	323	1 - 126	938		Trp-25 to Trp-42.		H0437: 3				
HUSHJ55	576381	324	2 - 337	939				H0437: 2	p36		118210, 120550, 120570, 120575, 121800, 130500,	

HUSHL83	868883	325	181 - 348	940	Glu-42 to Val-47.	H0437: 1 and H0268: 1.	133200, 155600, 171760, 171760, 185470, 211420, 230350, 255800, 601990, 602023, 602771
HUSIA38	709472	326	79 - 282	941	Ala-49 to Gln-56.	H0412: 2	
HUSIA43	575768	327	28 - 249	942	Asn-28 to Asp-33.	H0412: 2	
HUSIF23	862494	328	169 - 447	943	Lys-1 to Thr-10, Ala-20 to Val-25.	S0005: 1 and H0412: 1. 22q13.2-q13.31	188826, 250100, 250800, 250800
HUSIS60	727153	329	2 - 235	944	Ala-47 to Trp-54.	H0266: 1 and H0412: 1.	
HUSIW10	963324	330	59 - 322	945		H0412: 2	
HUSIW78	772956	331	3 - 188	946	Asn-6 to Arg-14.	H0412: 2	
HUSKI76	914084	332	3 - 230	947	Pro-65 to Gly-73.	H0412: 2	
HUSXK92	848959	333	221 - 487	948	Trp-1 to Asp-13.	H0413: 2, L0754: 1.	

					Ser-23 to Arg-28, Pro-44 to Arg-60, Leu-79 to Lys-89.	L0747: 1 and L0588: 1.		
HUSXO30	973266	334	203 - 463	949	Met-9 to Ser-16, Pro-24 to Asn-29.	H0412: 1, H0413: 1 and L0749: 1.		
HUSYA63	928021	335	158 - 442	950		H0412: 1 and H0413: 1.		
HUSYB16	868843	336	3 - 308	951	Ser-24 to Thr-29, Asn-58 to Arg-63.	H0413: 2		
HUSYD15	699195	337	106 - 279	952	Trp-16 to Leu-21.	H0266: 1 and H0413: 1.		
HUSYO46	868827	338	190 - 447	953	Lys-1 to Leu-10, Pro-43 to Gln-48, Thr-63 to Gly-69, Asn-76 to Arg-86.	H0413: 2		
HUSYP67	575787	339	2 - 118	954	Arg-1 to Ser-7, Glu-32 to Glu-37.	H0413: 2		
HUSZV72	851170	340	142 - 399	955		S0045: 1, H0413: 1, L0756: 1 and L0588: 1.		
HUSZH03	922852	341	13 - 714	956		L0754: 3 and H0413: 1.		
HUSYX03	922840	342	272 - 397	957	Leu-16 to Gln-21.	H0413: 1, L0438: 1 and L0439: 1.		
HUSYO86	784691	343	189 - 362	958	Glu-50 to Pro-58.	H0413: 1 and L0747: 1.		

HUSYN33	651293	344	189 - 371	959	Cys-34 to Gly-41.	H0413: 1 and L0439: 1.		
HUSYN11	943237	345	133 - 360	960	Pro-21 to Lys-63.	AR089: 11, AR061: 9 H0413: 1		
HUSYM49	723015	346	1 - 159	961		H0413: 1		
HUSYM37	464221	347	2 - 211	962		H0413: 1		
HUSYI13	657288	348	524 - 670	963		L0439: 7, L0759: 4, L0803: 2, L0438: 2, L0740: 2, L0751: 2, L0758: 2, H0413: 1, L0776: 1, L0367: 1, L0790: 1, L0666: 1, L0747: 1 and L0777: 1.		
HUSYG24	677258	349	53 - 277	964		L0439: 2, L0756: 2 and H0413: 1.		
HUSYF74	554723	350	92 - 328	965		L0439: 2, L0752: 2 and H0413: 1.		
HUSYA27	934423	351	459 - 638	966	Lys-47 to Lys-58.	L0717: 1, H0413: 1, L0748: 1, L0749: 1, L0779: 1, L0759: 1, L0608: 1, L0593: 1 and L0595: 1.		
HUSXW61	741856	352	386 - 559	967		L0740: 2 and H0413: 1.		
HUSXM28	703326	353	1 - 372	968	Pro-9 to Gln-22.	L0748: 2 and H0413:		

HUSXI71	760415	354	3 - 83	969			1.	H0413: 1 and L0748: 1.			
HUSXH57	859907	355	3 - 347	970			H0413: 1 and L0747: 1.				
HUSKA86	784887	356	2 - 229	971			L0748: 2 and H0412: 1.	8q22.3-q23.1	216550		
HUSKA65	868860	357	3 - 329	972		Gly-53 to Ile-59.	H0412: 1 and L0367: 1.				
HUSJW03	923035	358	121 - 270	973		Gly-20 to Gly-25.	L0748: 2, L0749: 2 and H0412: 1.				
HUSJN66	886987	359	3 - 455	974		Arg-22 to Pro-31, Pro-39 to Arg-50, Asp-86 to Gly-91, Phe-96 to Ser-105, Gly-114 to Gly-120, Asp-132 to Ser-139.	AR050: 86, AR054: 82, AR051: 63, AR089: 14, AR061: 11 H0412: 1 and L0759: 1.				
HUSIT75	679416	360	227 - 466	975		Arg-45 to Trp-50.	H0412: 1 and L0747: 1.				
HUSIS59	739327	361	1 - 231	976		Pro-10 to Gly-22, Gly-29 to Lys-47.	H0412: 1				
HUSIS54	730734	362	60 - 158	977		Val-22 to Leu-30.	H0412: 1				
HUSIS08	959536	363	93 - 425	978			H0412: 1 and L0593: 1.				
HUSIR04	709228	364	178 - 330	979		Ser-31 to Val-39.	L0748: 2 and H0412: 1.				

HUSIN12	970759	365	1 - 474	980	Leu-1 to Glu-10.	1.	H0412: 1 and L0665: 1.		
HUSIE95	967176	366	2 - 445	981			H0412: 1 and L0747: 1.		
HUSIE18	666523	367	104 - 280	982	Gly-1 to Lys-7, Arg-11 to Asn-23.		H0412: 1 and L0742: 1.		
HUSIE08	908574	368	335 - 661	983	Arg-19 to Ser-26.		L0157: 2, L0758: 2, H0412: 1, L0803: 1, L0744: 1, L0777: 1 and L0753: 1.		
HUSHL86	960355	369	779 - 1003	984	Ser-12 to Gly-17.		L0662: 2, H0437: 1 and L0731: 1.		
HUSHE34	703409	370	276 - 425	985			H0437: 1 and L0777: 1.		
HUSHB71	766060	371	1 - 267	986	Gly-1 to Arg-11, Asn-29 to Arg-35, Phe-37 to Tyr-48, His-74 to Cys-81.		H0437: 1 and L0748: 1.		
HUSHB60	746560	372	2 - 421	987	Glu-10 to Met-17, Thr-23 to Asp-32, Pro-37 to Gln-43, Val-67 to Asp-75.		H0437: 1 and L0596: 1.		
HUSGW06	935574	373	296 - 574	988	Gln-17 to Ser-29.		H0412: 1, L0754: 1 and L0752: 1.		



HUSGV84	813557	374	3 - 113	989	Asp-1 to Ser-8, Ser-22 to Asn-30.	H0412: 1 and L0598: 1.	13q12-q14	109543, 121011, 121011, 129500, 253700, 600631, 601499, 601885, 602221
HUSGU08	959540	375	105 - 281	990	Tyr-25 to Ile-30.	H0412: 1 and L0657: 1.		
HUSGT01	916620	376	3 - 215	991	Glu-61 to Ile-67.	H0412: 1 and L0532: 1.		
HUSGS35	707777	377	97 - 507	992	Thr-11 to Gly-17.	L0777: 3, L0766: 2, L0776: 2, L0759: 2, H0412: 1, L0638: 1, L0646: 1, L0764: 1, L0662: 1, L0659: 1, L0666: 1, L0663: 1, L0740: 1, L0749: 1 and L0588: 1.	19p13.2	108725, 120700, 133171, 143890, 147670, 147670, 147670, 151440, 164953, 231670, 600276, 600957,

[illegible]

Variable	Mean	Standard deviation	Minimum	Maximum
Age	34.5	10.5	20	55
Gender	0.5	0.5	0	1
Marital status	0.5	0.5	0	1
Education	12.5	2.5	9	16
Income	15.5	10.5	5	35
Health status	0.5	0.5	0	1
Smoking status	0.5	0.5	0	1
Alcohol consumption	0.5	0.5	0	1
Exercise frequency	0.5	0.5	0	1
Stress level	0.5	0.5	0	1
Sleep quality	0.5	0.5	0	1
Work satisfaction	0.5	0.5	0	1
Life satisfaction	0.5	0.5	0	1
Depression score	10.5	10.5	0	30
Anxiety score	10.5	10.5	0	30
Overall health score	10.5	10.5	0	30

						Asp-128 to Asp-134, Val-157 to Glu-168, Ser-182 to Lys-192.				223900, 253800, 253800, 268900, 278700, 602088
HUSFE05	932106	390	57 - 263	1005				H0433: 1, L0766: 1 and L0595: 1.		
HUSDA09	461656	391	260 - 487	1006		Gly-10 to Phe-16.		H0403: 1 and L0589: 1.		
HUSAY21	920403	392	255 - 503	1007		Val-27 to Asp-39, Thr-41 to Gly-46, Ser-54 to Thr-63, Asp-77 to Pro-83.		H0268: 1 and L0754: 1.		
HUSAO27	955287	393	160 - 270	1008				H0268: 1		
HUSAM87	529783	394	3 - 137	1009		Arg-1 to Lys-6, Gly-14 to Cys-20.		H0268: 1		
HUSAM35	558191	395	187 - 2	1010				H0268: 1		
HUSAM22	523674	396	3 - 95	1011		Glu-20 to Lys-29.		H0268: 1		
HUSAL04	927719	397	190 - 330	1012				H0268: 1		
HUSAJ57	678932	398	127 - 351	1013		Leu-1 to Thr-6.		H0268: 1		
HUSAJ15	522056	399	105 - 203	1014		Leu-18 to Gln-30.		H0268: 1		
HULAG30	788577	400	2 - 277	1015		Gly-8 to Arg-26.		L0731: 2 and H0530: 1.		
HMELY25	678120	401	201 - 311	1016				L0777: 2, H0266: 1,		

									L0766: 1, L0803: 1 and L0748: 1.			
HMELV14	876087	402	86 - 223	1017	Ser-38 to Thr-46.				H0266: 1, L0746: 1 and L0779: 1.			
HMELR45	717696	403	192 - 52	1018	Lys-14 to Arg-20, Gly-27 to Thr-40.				H0266: 1			
	717797	610	186 - 389	1225	Arg-29 to Leu-37.							
HMELM86	784702	404	23 - 133	1019	Phe-32 to Glu-37.				H0266: 1			
HMELM85	783536	405	72 - 224	1020					H0266: 1			
HMELM03	924168	406	119 - 316	1021					H0266: 1			
HMELI57	734769	407	2 - 292	1022	Met-1 to Gln-15, Ser-22 to Ala-31, Glu-45 to Val-51, Met-58 to Ser-63, Thr-74 to Asn-84, Met-90 to Pro-97.				H0266: 1 and L0591: 1.			
HMELH60	422844	408	338 - 3	1023					H0266: 1			
	878666	611	2 - 166	1226								
HMEKZ06	935467	409	43 - 297	1024	Asp-13 to Asn-25, Ile-41 to Pro-52.				H0266: 1 and L0780: 1.	4		
HMEKW07	953369	410	354 - 623	1025					H0266: 1 and L0776: 1.			
	668659	411	289 - 585	1026					H0266: 1, L0748: 1 and L0759: 1.			
HMEKO03	924172	412	159 - 338	1027					L0766: 2, H0266: 1			

HMEKJ40	711187	413	119 - 340	1028	Ile-13 to Cys-20, Ile-36 to Ser-41.	and L0779: 1.		
HMEKH73	923893	414	65 - 541	1029	Asp-14 to Leu-49, Pro-52 to Pro-73.	H0266: 1 and L0748: 1.		
HMEKC72	760637	415	199 - 393	1030	Cys-1 to Asp-8.	H0266: 1 and L0759: 1.		
HMEJW50	724396	416	61 - 237	1031		H0266: 1 and L0743: 1.		
HMEJ81	777945	417	73 - 309	1032	Arg-1 to Asp-9, Arg-22 to His-33, Ala-37 to Glu-47.	H0266: 1 and L0748: 1.		
HMEJF25	678131	418	99 - 425	1033	Val-30 to Thr-40, Lys-75 to Trp-88, Lys-90 to Ser-95.	H0266: 1 and L0731: 1.		
HMEIV22	674611	419	408 - 503	1034		H0266: 1 and L0748: 1.		
HMEIS07	922703	420	1 - 387	1035	Gly-1 to Arg-7, Pro-13 to Pro-37.	H0266: 1, L0745: 1 and L0603: 1.		
HMEIA06	935966	421	19 - 168	1036	Pro-6 to Gly-11.	H0266: 1		
HMEGK14	796443	422	11 - 334	1037	Gln-18 to Lys-29, Arg-43 to Leu-55, Pro-57 to Pro-67, Pro-81 to Gly-86.	L0749: 2 and H0266: 1.		
HMEGH92	790629	423	18 - 494	1038	Asp-21 to Arg-31.	H0266: 1 and L0766:		

HMEFD72	766185	424		1 - 234	1039	Leu-6 to Thr-11, His-13 to Gly-21, Gln-52 to Arg-62.	1.	H0266: 1 and L0766: 1.			
HMEEL38	733649	425		3 - 200	1040			H0266: 1, L0748: 1, L0756: 1 and L0599: 1.	6p12	180297, 230450, 263200, 601690	
HMEDR76	529897	426		18 - 173	1041			H0266: 1			
HMECQ10	968500	427		2 - 148	1042	Trp-8 to Ser-14, Arg-30 to Arg-39.	1.	H0266: 1 and L0766: 1.			
HMECH43	715568	428		2 - 133	1043			H0266: 1 and L0766: 1.			
HMEBY95	796058	429		172 - 327	1044			H0267: 1 and L0766: 1.			
HMEBG01	921763	430		94 - 240	1045			H0267: 1 and L0744: 1.			
HMEAN12	655220	431		59 - 259	1046			H0266: 1 and L0754: 1.			
HMEAI38	709207	432		325 - 504	1047	Pro-2 to Gly-7, Asp-13 to Ser-20.	1.	H0266: 1 and L0748: 1.			
HMEAH31	698403	433		68 - 190	1048	Gly-32 to Asp-39.		L0581: 2, L0471: 1, H0266: 1 and L0364: 1.			
HMEAE24	880925	434		2 - 169	1049			H0266: 1			
HMEAE01	916744	435		2 - 118	1050			H0266: 1			

HMEAD86	785802	436	3 - 299	1051	Phe-32 to Arg-37, Cys-40 to Thr-45, Glu-63 to Gly-70.	H0266: 1 and L0766: 1.		
HMEAD47	720638	437	2 - 346	1052	Arg-41 to Ser-46, Pro-52 to Gly-57.	H0266: 1 and L0748: 1.		
HMEAA17	921765	438	105 - 332	1053		H0266: 1 and L0439: 1.		
HHFUB83	800580	439	2 - 235	1054	Gly-1 to Ala-7.	H0242: 1		
HHFOU02	918070	440	142 - 432	1055	Ser-7 to Ser-14.	L0803: 2, H0645: 1 and L0747: 1.		
HHFOL43	974002	441	16 - 324	1056	His-1 to His-10, Pro-14 to Asn-19, Leu-48 to Arg-56, Lys-65 to Ile-76, Lys-93 to Arg-103.	H0645: 1		
HHFOK10	961346	442	22 - 279	1057	Ser-2 to Ser-8, Lys-14 to Arg-19.	H0645: 1 and L0766: 1.		
HHFNJ05	930899	443	3 - 320	1058	Pro-13 to Gly-22, Arg-31 to Gly-42, Ser-86 to Cys-92, Pro-94 to Pro-101.	L0774: 2, H0619: 1 and L0769: 1.		
HHFMX34	945385	444	3 - 461	1059	Ala-7 to Gly-13, Ile-71 to Gln-77, Phe-79 to Ala-85, Gly-142 to Pro-148.	AR089: 11, AR061: 8 H0619: 1		

HHFLU06	857884	445	2 - 328	1060		AR061: 5, AR089: 2 H0619: 1		
HHFLT84	857890	446	407 - 225	1061		H0619: 1 and L0745: 1.		
HHFLL08	958103	447	67 - 354	1062	Asn-6 to Asn-27, Ser-44 to Tyr-52.	H0619: 1 and L0599: 1.		
HHFLJ51	857898	448	186 - 443	1063	Phe-12 to His-17, Gly-53 to Tyr-60.	L0617: 1 and H0619: 1.		
HHFLI10	963162	449	64 - 471	1064	Thr-11 to Lys-19, Ser-44 to Asn-49, Cys-102 to Ala-111, Pro-114 to Thr-120.	H0619: 1		
HHFLI07	952081	450	404 - 589	1065	Ser-1 to Cys-24.	H0619: 1		
HHFLH62	857908	451	46 - 240	1066	Thr-26 to Tyr-31, Ile-52 to Cys-65.	H0619: 1 and L0759: 1.		
HHFLE12	969531	452	2 - 205	1067		H0619: 1 and L0758: 1.		
HHFKX28	971102	453	454 - 176	1068	Tyr-41 to Gly-52, Leu-66 to Trp-71, Ser-73 to Cys-85.	L0766: 2 and H0619: 1.		
HHFKB24	887025	454	3 - 173	1069	Asp-1 to Ser-14.	AR054: 27, AR050: 20, AR051: 19 H0619: 1		
HHFJM64	958384	455	170 - 1030	1070	Val-4 to Asn-11, Cys-24 to Tyr-31.	H0619: 1 and L0805: 1.		



HHFIA58	858011	456	99 - 284	1071	Glu-31 to Ala-40.	H0050: 1 and L0070: 1.		
HHFIA13	657405	457	108 - 308	1072	Val-19 to Trp-24, Ser-59 to Trp-67.	L0745: 2 and H0050: 1.		
HHFHY95	795053	458	13 - 243	1073		H0050: 1 and L0599: 1.		
HHFHR63	745215	459	164 - 391	1074	Pro-36 to Asn-45.	H0050: 1, L0748: 1 and L0439: 1.		
HHFHP68	753144	460	232 - 432	1075		H0050: 1 and L0666: 1.		
HHFHN74	765758	461	602 - 799	1076		L0439: 7 and H0050: 1.		
HHFHM22	674841	462	18 - 287	1077	Leu-6 to Gln-13.	H0050: 1 and L0439: 1.		
HHFHJ90	675218	463	200 - 364	1078	Gln-32 to Phe-37.	H0050: 1 and L0748: 1.		
HHFHD38	709082	464	33 - 149	1079		H0050: 1 and L0748: 1.		
HHFGX13	656806	465	566 - 414	1080	Pro-8 to Leu-22, Leu-42 to Glu-49.	H0050: 1		
	657080	612	30 - 131	1227	Pro-1 to Gln-6.			
HHFGX03	924753	466	3 - 338	1081	Trp-5 to Asp-16, Asp-25 to Asp-30, Arg-49 to Pro-54.	H0050: 1 and L0766: 1.		
HHFGR31	953204	467	109 - 405	1082	Leu-22 to Gly-31.	H0050: 1 and L0750: 1.		

HHFGR30	692887	468	1 - 141	1083	Thr-25 to Thr-30, Gly-32 to Asp-37.	1.	H0050: 1 and L0589:		
HHFGP91	800328	469	52 - 579	1084		1.	H0050: 1 and L0794:		
HHFGP69	918393	470	712 - 485	1085	Asn-1 to Gly-9.		AR051: 57, AR050: 56, AR054: 46 H0050: 1		
	918394	613	281 - 126	1228	Phe-4 to Gln-24, Thr-29 to Leu-38, Ser-46 to Arg-52.				
HHFGL77	490379	471	322 - 119	1086	Val-3 to Ser-29, Phe-47 to Gly-52.		H0050: 1		
	570229	614	120 - 275	1229					
HHFGH81	778193	472	46 - 267	1087	Asp-14 to Gln-22, Gln-35 to Phe-41.		H0050: 1 and L0439:		
HHFGH43	573495	473	2 - 202	1088		1.	L0744: 3, H0050: 1, L0743: 1 and L0748: 1.	2p14-pl3.4	203800
HHFFZ50	513773	474	263 - 436	1089			H0050: 1		
	858039	615	103 - 309	1230					
HHFFT05	932675	475	3 - 137	1090	Thr-2 to Ser-19, Glu-21 to Trp-30.		H0050: 1		
HHFFT01	880757	476	179 - 400	1091	Arg-37 to Phe-45.		AR051: 33, AR054: 28, AR050: 14 H0050: 1		

HHFFR95	796677	477	1 - 213	1092		L0439: 2 and H0050: 1.		
HHFFR75	766630	478	483 - 250	1093	Pro-1 to Ala-8.	H0050: 1		
	858041	616	207 - 335	1231				
HHFFR32	699723	479	92 - 313	1094	Pro-6 to Gly-12.	H0050: 1 and L0777: 1.		
HHFFP17	880667	480	215 - 409	1095		H0050: 1 and L0803: 1.		
HHFFO46	530501	481	32 - 286	1096	Ser-1 to Cys-6, Ala-28 to His-34.	H0050: 1		
HHFFM05	932738	482	324 - 470	1097		H0050: 1 and L0748: 1.		
HHFFL66	530503	483	1 - 150	1098	Ile-1 to Ala-6.	H0050: 1		
HHFFK30	858051	484	2 - 193	1099	Ile-8 to Gly-13.	H0050: 1 and L0378: 1.		
HHFFI59	739587	485	74 - 280	1100	Gln-22 to Ser-29.	H0050: 1 and L0748: 1.		
HHFFI08	960254	486	49 - 342	1101	Thr-13 to Ala-18, Ala-39 to Ser-46, Cys-51 to Thr-58, Pro-70 to Tyr-75, Glu-83 to Gln-95.	L0748: 2, L0002: 1 and H0050: 1.		
HHFFH76	767623	487	265 - 546	1102	Ala-1 to Asn-6.	H0050: 1 and L0750: 1.	2q32.1	600258, 602087
HHFFG82	530662	488	3 - 281	1103	Ala-21 to Asn-27.	H0050: 1		

HHFFF92	790572	489	52 - 345	1104		L0745: 3, L0750: 2 and H0050: 1.		
HHFFF07	954258	490	409 - 576	1105		H0050: 1 and L0749: 1.		
HHFEB86	785653	491	161 - 367	1106		H0050: 1 and L0748: 1.		
HHFDN80	781634	492	93 - 287	1107	Ile-7 to Asn-26, Gly-35 to Gly-42, Ala-44 to Gly-49.	H0050: 1 and L0591: 1.	6p22	248611
HHFDI82	499010	493	370 - 492	1108	Ile-18 to Asp-28, Ile-32 to Asn-37.	H0050: 1		
	511062	617	518 - 306	1232				
HHFDH26	685188	494	1 - 441	1109	Ser-36 to Lys-53, Pro-63 to Pro-79, Val-82 to Gly-104.	H0050: 1 and L0754: 1.		
HHFDC10	968647	495	2 - 136	1110	Ser-34 to Leu-40.	L0748: 2, H0050: 1 and L0599: 1.		
HHFDA13	667804	496	221 - 337	1111	Met-6 to Ile-18.	H0050: 1 and L0748: 1.		
HHFCT63	572784	497	184 - 32	1112		H0050: 1 and L0748: 1.		
HHFPC39	429442	498	81 - 194	1113	Asn-24 to Asn-36.	H0050: 1 and L0748: 1.		
HHFCO13	500899	499	322 - 221	1114		H0050: 1 and L0754: 1.		

HHFCN59	739657	500	386 - 138	1115	His-39 to Pro-45.	H0050: 1 and L0750: 1.		
HHFCN13	667805	501	221 - 6	1116	Phe-38 to Lys-44.	L0749: 3 and H0050: 1.		
HHFCM51	509631	502	23 - 151	1117	Asn-28 to Leu-33.	H0050: 1		
HHFCH52	911570	503	307 - 50	1118		H0050: 1		
HHFCF58	575183	504	166 - 333	1119	Ser-2 to Gly-12.	L0748: 3, H0050: 1 and L0750: 1.		
	575184	618	500 - 159	1233				
HHFCE73	764763	505	5 - 214	1120		L0748: 3 and H0050: 1.		
HHFCE40	712866	506	71 - 199	1121		H0050: 1 and L0439: 1.		
HHFCD43	714353	507	52 - 150	1122		H0050: 1, L0803: 1 and L0596: 1.		
HHFCC60	739669	508	197 - 436	1123		H0050: 1 and L0439: 1.		
HHFCC45	858066	509	23 - 217	1124		H0050: 1 and L0746: 1.		
HHFCC20	600231	510	605 - 411	1125		H0050: 1		
	825658	619	3 - 344	1234	Gln-41 to Gly-53, Pro-55 to Glu-64, Arg-108 to Gly-113.			

HHFBW92	575156	511	393 - 551	1126			H0050: 1 and L0748: 1.		
HHFBU63	745661	512	212 - 382	1127	Lys-3 to Arg-21, Pro-40 to Lys-47.		H0050: 1 and L0748: 1.		
HHFBU07	954478	513	172 - 405	1128			H0050: 1 and L0804: 1.		
HHFBT24	508067	514	2 - 418	1129			H0050: 1		
HHFBQ94	796838	515	108 - 278	1130			L0750: 2, H0050: 1 and L0604: 1.		
HHFBP29	710894	516	1 - 279	1131	Val-17 to Thr-23, Ser-57 to Arg-64.		L0748: 5, H0050: 1, L0805: 1 and L0749: 1.		
HHFBN17	589798	517	180 - 1	1132	Ser-30 to Tyr-35, Leu-52 to Glu-60.		H0050: 1		
	858072	620	358 - 528	1235					
HHFBM11	968002	518	224 - 3	1133	Val-36 to Ser-54.		L0758: 4, H0050: 1, L0769: 1 and L0438: 1.		
HHFBD83	781525	519	602 - 411	1134			L0748: 2, H0050: 1 and L0745: 1.		
HHFBD42	712899	520	463 - 77	1135			L0754: 3, L0749: 2, H0050: 1, L0662: 1 and L0596: 1.		
HHFBB14	522375	521	3 - 191	1136	Arg-26 to Glu-37.		H0050: 1 and L0605: 1.		

HHFBA11	967991	522	181 - 321	1137	Asp-1 to Thr-9.	H0050: 1		
HHFAB62	824590	523	608 - 255	1138	Arg-4 to Asp-9.	L0439: 4 and H0019: 1.		
HHBGN74	765214	524	145 - 471	1139	Phe-12 to Pro-22, Glu-29 to Asn-38, Glu-45 to Tyr-55.	H0373: 1 and L0604: 1.		
HHBGJ53	909912	525	1 - 282	1140	Ser-1 to Ser-7, Ser-25 to Arg-31.	AR089: 8, AR061: 5 L0740: 2 and H0373: 1.		
HHBGG10	963849	526	215 - 400	1141	Lys-1 to Thr-7, Gln-12 to Glu-25.	H0373: 1 and L0752: 1.		
HHBGC75	767042	527	16 - 219	1142	Lys-48 to Lys-63.	L0777: 3 and H0373: 1.		
HHBFT06	934826	528	358 - 489	1143	Asp-6 to Ile-12, Met-21 to Ala-26.	H0373: 1 and L0142: 1.		
HHBFM77	771816	529	186 - 266	1144		H0373: 1 and L0753: 1.		
HHBEV93	792041	530	2 - 139	1145		L0742: 3 and H0373: 1.		
HHBES89	786667	531	216 - 398	1146	His-12 to Ser-17.	L0439: 4 and H0373: 1.		
HHBEM49	722337	532	288 - 455	1147	Lys-1 to Phe-6.	H0373: 1, L0749: 1 and L0601: 1.		

HHBEG80	951688	533	284 - 442	1148	Ser-11 to Cys-26, Lys-30 to Lys-53.	AR089: 21, AR061: 7 H0373: 1 and L0731: 1.		
HHBEG72	761150	534	157 - 402	1149	Arg-29 to Ser-34.	H0373: 1 and L0748: 1.		
HEMHA53	728297	535	450 - 605	1150		S0046: 1 and L0749: 1.		
HEMGX57	872083	536	382 - 606	1151		S0046: 1 and L0766: 1.		
HEMGT27	851065	537	123 - 293	1152	Cys-36 to Trp-42.	S0046: 1 and L0592: 1.		
HEMGL56	767669	538	3 - 332	1153	Gly-89 to Ala-95.	AR089: 25, AR061: 7 S0046: 1	150250, 164500, 277730, 600971, 601226	
HEMFN30	692818	539	344 - 805	1154	Gln-1 to Asn-8, Lys-17 to Arg-24, Ser-89 to Arg-103.	L0598: 2, L0731: 2 and S0046: 1.		
HEMFF16	576539	540	3 - 140	1155	Asn-1 to Pro-11.	S0046: 1		
HEMEF34	596812	541	477 - 695	1156	Gly-37 to Val-43.	S0046: 1		
	851083	621	3 - 182	1236	Gly-24 to Val-30.			
HEMEA03	921922	542	124 - 258	1157		S0046: 1 and L0749: 1.		
HEMDX96	935963	543	650 - 255	1158		L0751: 2 and S0046: 1.		
HEMCV44	574321	544	235 - 345	1159	Tyr-6 to Ser-11.	S0046: 1		



HEMCK53	728424	545	69 - 251	1160	Gly-11 to Asp-24, Ser-41 to Ile-46.	L0748: 2 and S0046: 1.		
HEMCI59	739551	546	1 - 354	1161	His-1 to Met-11, Asp-40 to Glu-48, Tyr-72 to Arg-77, Asn-111 to His-117.	AR089: 4, AR061: 3 S0046: 1 and L0740: 1.		
HEMCI41	712614	547	300 - 578	1162		L0805: 2, L0608: 2, S0046: 1, L0775: 1, L0375: 1, L0651: 1, L0756: 1, L0759: 1 and L0592: 1.		
HEMCC38	707453	548	201 - 341	1163		S0046: 1 and L0748: 1.		
HEMBU26	684928	549	1 - 81	1164	Thr-11 to Lys-17.	S0046: 1 and L0754: 1.		
HEMBT61	939957	550	1 - 351	1165		AR061: 8, AR089: 4 L0547: 2, S0046: 1, L0471: 1, L0772: 1, L0529: 1 and L0780: 1.		
HEMAL61	851106	551	130 - 270	1166	Pro-4 to Lys-13, Leu-21 to Pro-26, Lys-32 to Gly-41.	S0046: 1		
HEMAA63	745498	552	3 - 149	1167		S0046: 1 and L0748: 1.		

HELHJ74	765696	553	181 - 513	1168	Arg-4 to Lys-26, Thr-31 to Gly-36, Glu-83 to Lys-88.	L0745: 3 and S0045: 1.		
HELHD20	668881	554	222 - 434	1169	Lys-1 to Trp-10, Thr-18 to Val-27.	S0045: 1		
HELHC59	769404	555	285 - 518	1170		S0045: 1 and L0740: 1.		
HELGY42	713019	556	47 - 220	1171		L0742: 2 and S0045: 1.		
HELGY02	948302	557	3 - 410	1172	His-1 to Gly-8, Leu-20 to Pro-30, Lys-44 to Glu-58, Pro-60 to Gln-67.	AR089: 0, AR061: 0 S0045: 1		
HELGW31	610003	558	576 - 1337	1173	Asp-56 to Ser-62, Gly-195 to Ser-202.	AR061: 335, AR089: 290, AR051: 10, AR050: 2, AR054: 2 S0045: 1		
	957568	622	1035 - 394	1237				
	964303	623	21 - 449	1238				
HELGV36	597120	559	233 - 373	1174	Arg-4 to Ala-10.	S0045: 1, L0748: 1 and L0754: 1.		
HELQ55	732223	560	1 - 315	1175		S0045: 1 and L0750: 1.		
HELK56	925698	561	129 - 788	1176		AR061: 3, AR089: 1 S0045: 1		
HELGG21	671071	562	2 - 253	1177		S0045: 1 and L0748: 1.		

[illegible]

HELGD47	851143	563	304 - 453	1178	Arg-1 to Cys-6, Pro-43 to Met-50.	S0045: 1 and L0717: 1.		
HELFAQ55	732224	564	116 - 265	1179	Thr-28 to Gln-34, His-38 to Thr-44.	S0045: 1 and L0439: 1.		
HELFN75	658681	565	264 - 635	1180		AR051: 6, AR054: 4, AR050: 1, AR089: 1, AR061: 0 S0045: 1		
HELFI35	506277	566	133 - 321	1181	Gln-6 to Lys-15.	S0045: 1		
HELFA38	851146	567	85 - 318	1182	Trp-55 to Gly-60.	S0045: 1		
	954009	624	482 - 210	1239				
HELEZ81	571340	568	3 - 329	1183	Pro-82 to Gln-87.	S0045: 1		
HELET68	800029	569	17 - 208	1184	Thr-1 to Glu-9.	S0045: 1 and L0748: 1.		
HELEO42	579016	570	3 - 467	1185	His-1 to Ser-20, Leu-35 to Glu-43, Ser-73 to Glu-79.	AR051: 19, AR054: 11, AR050: 9 S0045: 1		
HELEH76	506674	571	1 - 411	1186		S0045: 1, L0766: 1 and L0754: 1.	102770, 188540, 600234, 601414, 601691, 601691, 601691, 601691, 601691,	

HELEE83	577206	572	1 - 123	1187	Gly-1 to Ala-7.	S0045: 1	601718, 602094
HELDT63	744864	573	635 - 516	1188	Arg-12 to Lys-19, Gly-35 to Gly-40.	AR051: 21, AR050: 18, AR054: 3 S0045: 1	
	898960	625	3 - 107	1240	Leu-28 to Ser-35.		
HELDL15	660557	574	170 - 406	1189	Lys-40 to Pro-48, Cys-66 to Cys-74.	L0750: 2, S0045: 1, L0803: 1 and L0783: 1.	
HELDL08	959919	575	267 - 377	1190		S0045: 1 and L0589: 1.	
HELDK22	567310	576	1 - 309	1191		S0045: 1	
HELDH71	740198	577	239 - 634	1192	Leu-6 to Asn-14, Thr-22 to Lys-47.	S0045: 1 and L0748: 1.	
HELDG91	790371	578	555 - 704	1193	Ala-24 to Pro-37.	S0045: 1, L0743: 1 and L0740: 1.	
HELWCW51	531073	579	110 - 262	1194	Asn-12 to Lys-20.	S0045: 1	
HELCI30	691024	580	12 - 200	1195		S0045: 1 and L0439: 1.	
HELCCG36	655045	581	96 - 218	1196		S0045: 1, L0754: 1 and L0731: 1.	
HELBUI1	967661	582	10 - 204	1197		L0748: 2, S0045: 1 and L0608: 1.	
HELBBC83	781412	583	132 - 425	1198		L0439: 2 and S0045: 1.	

HELAZ48	864515	584	2 - 226	1199			L0439: 2, L0740: 2, S0045: 1, L0750: 1 and L0752: 1.		
HELAW26	684925	585	279 - 542	1200			L0748: 3 and S0045: 1.		
HELAQ36	707420	586	69 - 176	1201			S0045: 1 and L0748: 1.		
HELAM32	699661	587	1 - 228	1202			S0045: 1 and L0748: 1.		
HELAH32	699665	588	1 - 249	1203			L0757: 2 and S0045: 1.		
HCMYS80	526182	589	1 - 153	1204			H0196: 1		
HCMSEQ63	745584	590	142 - 303	1205		Pro-1 to Pro-7, Pro-10 to Asn-15.	H0196: 1 and L0777: 1.		
HCMK41	940260	591	109 - 402	1206		Ser-1 to Trp-6, Leu-24 to Ser-29.	AR061: 3, AR089: 1 H0196: 1		
HAHSB27	501010	592	196 - 417	1207		Trp-5 to His-16, Gln-45 to Phe-50.	H0097: 1		
HAHFS80	954432	593	202 - 510	1208		Pro-12 to Thr-19.	L0623: 1, H0599: 1, L0471: 1 and L0485: 1.		
HAHFE11	965293	594	88 - 282	1209			H0599: 1 and L0758: 1.		
HAHEP68	738501	595	84 - 431	1210		Ser-1 to Val-14, Ser-26 to Glu-39, Ala-48 to Pro-53, Trp-62 to Thr-68, Lys-74 to Asn-84, Thr-89 to Pro-95,	H0599: 1, L0439: 1 and L0747: 1.		

HAHCU22	84831	596	35 - 211	1211	Val-99 to Thr-108. Thr-18 to Gly-26.	H0599: 1 and L0471: 1.			
HAHCR15	810326	597	3 - 455	1212		H0599: 1, L0471: 1 and L0602: 1.			
HAHCL94	794044	598	2 - 316	1213	Ala-1 to Ala-6, Ser-10 to Gly-17.	H0599: 1 and L0748: 1.	5p13.1-5cen	108962, 600837	
HAHBC03	923542	599	232 - 519	1214	Gln-48 to Trp-71, His-87 to Lys-94.	H0002: 1 and L0758: 1.	1p36-p34	118210, 120550, 120570, 120575, 121800, 130500, 133200, 138140, 155600, 168360, 171760, 171760, 176100, 176100, 178300, 185470, 211420, 230000,	

									230350, 255800, 601990, 602023, 602771
HAHAD95	865104	600	106 - 279	1215	Gly-7 to Gly-14.			H0002: 1	
HAFBG30	693363	601	284 - 472	1216				T0049: 1 and L0731: 1.	
HAFAY37	928705	602	332 - 529	1217				T0049: 1, L0598: 1, L0766: 1 and L0748: 1.	
HAFJ63	845452	603	273 - 497	1218				T0049: 1, L0748: 1 and L0754: 1.	
HAECA04	932993	604	40 - 195	1219	Glu-1 to Arg-9, Leu-17 to Ala-25.			T0048: 1 and L0060: 1.	
HAEM82	781539	605	272 - 571	1220	Tyr-13 to Met-23, Ser-40 to Pro-49.			T0048: 1 and L0595: 1.	

10057504.030706

[0052] The first column in Table 1A provides a unique "Clone ID NO:Z" for a cDNA clone related to each contig sequence disclosed in Table 1A. This clone ID references the cDNA clone which contains at least the 5' most sequence of the assembled contig, and at least a portion of SEQ ID NO:X was determined by directly sequencing the referenced clone. The reference clone may have more sequence than described in the sequence listing or the clone may have less. In the vast majority of cases, however, the clone is believed to encode a full-length polypeptide. In the case where a clone is not full-length, a full-length cDNA can be obtained by methods known in the art and/or as described elsewhere herein.

[0053] The second column in Table 1A provides a unique "Contig ID" identification for each contig sequence. The third column provides the "SEQ ID NO:X" identifier for each of the cardiovascular system associated contig polynucleotide sequences disclosed in Table 1A. The fourth column, "ORF (From-To)", provides the location (i.e., nucleotide position numbers) within the polynucleotide sequence "SEQ ID NO:X" that delineate the preferred open reading frame (ORF) shown in the sequence listing and referenced in Table 1A, column 5, as SEQ ID NO:Y. Where the nucleotide position number "To" is lower than the nucleotide position number "From", the preferred ORF is the reverse complement of the referenced polynucleotide sequence.

[0054] The fifth column in Table 1A provides the corresponding SEQ ID NO:Y for the polypeptide sequence encoded by the preferred ORF delineated in column 4. In one embodiment, the invention provides an amino acid sequence comprising, or alternatively consisting of, a polypeptide encoded by the portion of SEQ ID NO:X delineated by "ORF (From-To)". Also provided are polynucleotides encoding such amino acid sequences and the complementary strand thereto.

[0055] Column 6 in Table 1A lists residues comprising epitopes contained in the polypeptides encoded by the preferred ORF (SEQ ID NO:Y), as predicted using the algorithm of Jameson and Wolf, (1988) Comp. Appl. Biosci. 4:181-186. The Jameson-Wolf antigenic analysis was performed using the computer program PROTEAN (Version 3.11 for the Power MacIntosh, DNASTAR, Inc., 1228 South Park Street Madison, WI). In specific embodiments, polypeptides of the invention comprise, or alternatively consist of, at least one, two, three, four, five or more of the



predicted epitopes as described in Table 1A. It will be appreciated that depending on the analytical criteria used to predict antigenic determinants, the exact address of the determinant may vary slightly.

[0056] Column 7 in Table 1A provides an expression profile and library code: count for each of the contig sequences (SEQ ID NO:X) disclosed in Table 1A, which can routinely be combined with the information provided in Table 4 and used to determine the normal or diseased tissues, cells, and/or cell line libraries which predominantly express the polynucleotides of the invention. The first number in column 7 (preceding the colon), represents the tissue/cell source identifier code corresponding to the code and description provided in Table 4. For those identifier codes in which the first two letters are not "AR", the second number in column 7 (following the colon) represents the number of times a sequence corresponding to the reference polynucleotide sequence was identified in the tissue/cell source. Those tissue/cell source identifier codes in which the first two letters are "AR" designate information generated using DNA array technology. Utilizing this technology, cDNAs were amplified by PCR and then transferred, in duplicate, onto the array. Gene expression was assayed through hybridization of first strand cDNA probes to the DNA array. cDNA probes were generated from total RNA extracted from a variety of different tissues and cell lines. Probe synthesis was performed in the presence of <sup>33</sup>P dCTP, using oligo(dT) to prime reverse transcription. After hybridization, high stringency washing conditions were employed to remove non-specific hybrids from the array. The remaining signal, emanating from each gene target, was measured using a Phosphorimager. Gene expression was reported as Phosphor Stimulating Luminescence (PSL) which reflects the level of phosphor signal generated from the probe hybridized to each of the gene targets represented on the array. A local background signal subtraction was performed before the total signal generated from each array was used to normalize gene expression between the different hybridizations. The value presented after "[array code]:" represents the mean of the duplicate values, following background subtraction and probe normalization. One of skill in the art could routinely use this information to identify normal and/or diseased tissue(s) which show a predominant expression pattern of the corresponding polynucleotide of the invention or to identify polynucleotides which show predominant and/or specific tissue and/or cell expression. The sequences disclosed

herein have been determined to be predominantly expressed in cardiovascular system tissues, including normal and diseased cardiovascular system tissues (See Table 1A, column 7 and Table 4).

[0057] Column 8 in Table 1A provides a chromosomal map location for certain polynucleotides of the invention. Chromosomal location was determined by finding exact matches to EST and cDNA sequences contained in the NCBI (National Center for Biotechnology Information) UniGene database. Each sequence in the UniGene database is assigned to a "cluster"; all of the ESTs, cDNAs, and STSs in a cluster are believed to be derived from a single gene. Chromosomal mapping data is often available for one or more sequence(s) in a UniGene cluster; this data (if consistent) is then applied to the cluster as a whole. Thus, it is possible to infer the chromosomal location of a new polynucleotide sequence by determining its identity with a mapped UniGene cluster.

[0058] A modified version of the computer program BLASTN (Altshul et al., J. Mol. Biol. 215:403-410 (1990), and Gish et al., Nat. Genet. 3:266-272 (1993)) was used to search the UniGene database for EST or cDNA sequences that contain exact or near-exact matches to a polynucleotide sequence of the invention (the 'Query'). A sequence from the UniGene database (the 'Subject') was said to be an exact match if it contained a segment of 50 nucleotides in length such that 48 of those nucleotides were in the same order as found in the Query sequence. If all of the matches that met this criteria were in the same UniGene cluster, and mapping data was available for this cluster, it is indicated in Table 1A under the heading "Cytologic Band". Where a cluster had been further localized to a distinct cytologic band, that band is disclosed; where no banding information was available, but the gene had been localized to a single chromosome, the chromosome is disclosed.

[0059] Once a presumptive chromosomal location was determined for a polynucleotide of the invention, an associated disease locus was identified by comparison with a database of diseases, which have been experimentally associated with genetic loci. The database used was the Morbid Map, derived from OMIM™ (*supra*). If the putative chromosomal location of a polynucleotide of the invention (Query sequence) was associated with a disease in the Morbid Map database, an OMIM reference identification number was noted in column 9, Table 1A, labeled "OMIM Disease

Reference(s)". Table 5 is a key to the OMIM reference identification numbers (column 1), and provides a description of the associated disease in Column 2.

10091504.030702

TABLE 1B

Clone ID NO:Z	SEQ ID NO:X	CONTIG ID:	BAC ID: A	SEQ ID NO:B	EXON From-To
HAHCL07	11	952661	AC022706	1241	1-1814
HAHCL07	11	952661	AC022916	1242	1-1814
HAHCL07	11	952661	AC022706	1243	1-107
HAHCL07	11	952661	AC022916	1244	1-107
HAHCP41	13	712086	AP002755	1245	1-326
HAHCP41	13	712086	AP000774	1246	1-326
HAHCP41	13	712086	AC020742	1247	1-326
HAHCP41	13	712086	AC001234	1248	1-326
HAHCP41	13	712086	AP002755	1249	1-538
HAHCP41	13	712086	AP000774	1250	1-538
HAHCP41	13	712086	AC020742	1251	1-538
HAHCP41	13	712086	AC001234	1252	1-538
HAHCP49	14	722659	AC046164	1253	1-244
HAHCP55	15	865095	AC011610	1254	1-125 1640-1708 1938-2132 3921-4067 4332-4975 6267-6343 6434-6748 6940-7010 9423-9456 9548-9900 12131-12199 13540-13627 13844-13921 14168-14201 14997-15076 15386-16024 16177-16523 16776-16849 17304-17711 17901-18042 18921-19274
HAHCP55	15	865095	AC066599	1255	1-69 524-931 1121-1262 2140-2493

20200405T600T

HAHCP55	15	865095	AC022382	1256	1-69 524-931 1121-1262 2142-2495
HAHCP55	15	865095	AC024163	1257	1-69 524-931 1121-1262 2140-2493
HAHCP55	15	865095	AC007850	1258	1-125 1640-1708 1938-2132 3921-4067 4332-4975 6267-6343 6434-6748 6940-7010 9423-9456 9548-9900 12130-12198 13539-13626 13843-13920 14167-14200 14996-15075 15385-16023 16176-16522 16774-16847 17302-17709 17899-18040 18918-19206
HAHCP55	15	865095	AC008034	1259	1-125 1640-1708 1938-2132 3921-4067 4332-4975 6267-6343 6434-6748 6940-7010 9637-9816 12130-12198 13539-13626 13843-13920

					14167-14200 14996-15075 15385-16023 16176-16522 16774-16847 17302-17709 17899-18040 18918-19271 20211-20888 21472-21834 22101-22510 24271-24974 25056-25409 25860-26636 26691-26822 27128-28128 30303-30454 30745-31348
HAHCP55	15	865095	AC066599	1260	1-347
HAHCP55	15	865095	AC022382	1261	1-347
HAHCP55	15	865095	AC011610	1262	1-564
HAHCP55	15	865095	AC066599	1263	1-678
HAHCP55	15	865095	AC022382	1264	1-678
HAHCP55	15	865095	AC024163	1265	1-347
HAHCP55	15	865095	AC007850	1266	1-564
HAHCP55	15	865095	AC008034	1267	1-564
HAHCP55	15	865095	AC024163	1268	1-678
HAHCP67	16	756916	AC044892	1269	1-160 1117-1442 1844-2103 2869-3022 3971-4116 5198-5472 6138-6590 7022-7205 7607-7703 7910-8685 9075-9227 9362-9724 9937-10537
HAHCP67	16	756916	AC006252	1270	1-160

					1116-1441 1843-2102 2868-3021 3978-4123 5226-5479 6145-6597 7029-7211 7613-7709 7916-8691 9081-9233 9368-9730 9943-10543
HAHCP67	16	756916	AC044892	1271	1-995 1441-1664
HAHCP67	16	756916	AC006252	1272	1-995 1441-1664
HAHCP91	17	789998	AC005833	1273	1-288
HAHCP91	17	789998	AC005833	1274	1-216
HAHCP91	17	789998	AC005833	1275	1-328 378-1059 1980-2100
HAHCR57	18	865096	AC073487	1276	1-587 2146-3842 3989-4201 5499-5529 6178-6728 7177-7679 8088-9978 10716-10977 10981-11359 11459-11652 11709-12255 12352-12457 12485-12662 12775-12853 12946-13797 13854-13972 14616-14764 15230-15403 15550-15610 15784-15879

					16027-16157 16273-16359 16716-16906 17040-17252 18205-18727 18898-19487 20516-20784 21205-21286 21386-21515 25809-26225
HAHCR57	18	865096	AC073487	1277	1-259
HAHEE04	19	922257	AL158169	1278	1-217 532-704 791-908 1623-1788 1831-1985 2107-2447 4415-4873 5085-5172 5281-5360 5579-5646 5738-5988 6535-8317
HAHEE04	19	922257	AL356104	1279	1-459 671-758 867-946
HAHEE04	19	922257	AL158169	1280	1-84
HAHEE04	19	922257	AL356104	1281	1-68 160-410 957-2739
HAHEO46	21	718772	AC008640	1282	1-988
HAHEO46	21	718772	AC011376	1283	1-988
HAHEO46	21	718772	AC008640	1284	1-308
HAHEO46	21	718772	AC011376	1285	1-308
HAHES10	22	961594	AC007618	1286	1-417
HAHES10	22	961594	AC007618	1287	1-545
HAHES10	22	961594	AC007618	1288	1-557
HAHFX20	23	925753	AC022965	1289	1-265
HAHFX20	23	925753	AL390725	1290	1-265
HAHFX20	23	925753	AC022965	1291	1-489
HAHFX20	23	925753	AL390725	1292	1-489



HAHFX20	23	925753	AC022965	1293	1-518
HAHFX20	23	925753	AL390725	1294	1-518
HAHHO04	25	925814	AC009417	1295	1-166 344-432 941-1480
HAHHO04	25	925814	AC073650	1296	1-31 529-793 1012-1296 1451-1539 2048-2587
HAHHO04	25	925814	AC009417	1297	1-289
HAHHO04	25	925814	AC073650	1298	1-289
HAHHS01	26	913863	AC011189	1299	1-290
HAHHS01	26	913863	AC027040	1300	1-290
HAHHS01	26	913863	AC002993	1301	1-290
HAHHS01	26	913863	AC011189	1302	1-562
HAHIK10	27	961425	AL354822	1303	1-172
HAHIK10	27	961425	AL159982	1304	1-168
HAHIK10	27	961425	AC025100	1305	1-170
HAHIK10	27	961425	AC023572	1306	1-173
HAHIK10	27	961425	AL354817	1307	1-173
HAHIK10	27	961425	AL354697	1308	1-173
HAHIK10	27	961425	AC000382	1309	1-170
HAHIK10	27	961425	AC025100	1310	1-293
HAHIK10	27	961425	AC023572	1311	1-293
HAHIK10	27	961425	AL354822	1312	1-293
HAHIK10	27	961425	AL354817	1313	1-455
HAHIK10	27	961425	AL354697	1314	1-455
HAHIK10	27	961425	AL159982	1315	1-292
HAHIW08	28	955803	AC002413	1316	1-647
HAHIW08	28	955803	AL139396	1317	1-647
HAHIW08	28	955803	AC002413	1318	1-352
HAHSC42	29	695111	AC073389	1319	1-119 1375-1607 2148-2689 2792-3076 3843-4509 4904-5519
HAHSC42	29	695111	AC073389	1320	1-457
HAHSC42	29	695111	AC073389	1321	1-327

HCMSC52	31	522615	AC026228	1322	1-111
HCMSQ77	33	862373	AC006463	1323	1-303
HCMSX59	36	522598	AC012134	1324	1-362
HCMSX59	36	522598	AC073904	1325	1-361
HCMSX59	36	522598	AP001811	1326	1-361
HCMSX59	36	522598	AP001811	1327	1-558
HCMSX59	36	522598	AC012134	1328	1-558
HCMSX59	36	522598	AC073904	1329	1-558
HELAF85	37	507230	AC020988	1330	1-320
HELAF85	37	507230	AC020988	1331	1-886 900-3110
HELAI26	38	920922	AL157829	1332	1-327 383-585 5577-5852 7628-7977 15024-15185 23099-23432
HELAI26	38	920922	AC015464	1333	1-276
HELAI26	38	920922	AL133284	1334	1-276
HELAI26	38	920922	AC015464	1335	1-350
HELAV61	40	507205	AL390880	1336	1-349
HELAV61	40	507205	AL390880	1337	1-264
HELAV61	40	507205	AL390880	1338	1-486
HELBD70	42	527677	AC026185	1339	1-848
HELBD70	42	527677	AC018494	1340	1-848
HELBD70	42	527677	AC026185	1341	1-543
HELBD70	42	527677	AC026185	1342	1-271
HELBD70	42	527677	AC018494	1343	1-543
HELBD70	42	527677	AC018494	1344	1-271
HELBN45	44	527666	AL031295	1345	1-293
HELBN45	44	527666	AL031295	1346	1-289
HELDG77	46	750478	AL159170	1347	1-541
HELDG77	46	750478	AL390024	1348	1-541
HELDG77	46	750478	AL159170	1349	1-483 551-674
HELEF52	51	506677	AC012273	1350	1-317
HELEF52	51	506677	AC012273	1351	1-333
HELER30	53	574038	AC006057	1352	1-1857
HELER30	53	574038	AC006057	1353	1-75
HELEU91	57	851160	AP002386	1354	1-337

HELEU91	57	851160	AP001922	1355	1-337
HELEU91	57	851160	AP000649	1356	1-352
HELEU91	57	851160	AP001142	1357	1-351
HELEU91	57	851160	AC011256	1358	1-352
HELEU91	57	851160	AP001142	1359	1-492
HELEU91	57	851160	AP000649	1360	1-492
HELEU91	57	851160	AC011256	1361	1-492
HELEU91	57	851160	AP001142	1362	1-156
HELEU91	57	851160	AC011256	1363	1-156
HELEW62	58	574025	AC020656	1364	1-398 2110-2216 2736-3244 5756-5912 6792-6965 7092-7180 7274-7632 7650-8158 8486-8607 8718-9027 9103-9257 10633-10932 11428-11621 11900-12183 12559-14453 14491-15109 15316-15541 16291-16817 16893-17205 17694-23419
HELEW62	58	574025	AC019161	1365	1-480 808-929 1040-1349 1425-1579 2955-3254 3750-3943
HELEW62	58	574025	AC020656	1366	1-115
HELEW62	58	574025	AC019161	1367	1-283
HELEW62	58	574025	AC019161	1368	1-355
HELFF40	59	574058	AC000119	1369	1-452 1221-1486 2326-2549

2009040516001

					2833-2982 3534-3620 9221-10633 11535-11911 12219-12718
HELFF40	59	574058	AC000119	1370	1-644
HELFH33	60	576530	AC025152	1371	1-267
HELFH33	60	576530	AC008941	1372	1-246
HELFH33	60	576530	AC010073	1373	1-253
HELFJ03	61	921943	AL360083	1374	1-824
HELFJ03	61	921943	AL360083	1375	1-285
HELFJ03	61	921943	AL360083	1376	1-133
HELFQ79	63	577248	AC025152	1377	1-359
HELFQ79	63	577248	AC021062	1378	1-349
HELFQ79	63	577248	AC018720	1379	1-548
HELFQ79	63	577248	AC034250	1380	1-356
HELFQ79	63	577248	AC020865	1381	1-670
HELGA54	64	576374	AC023611	1382	1-180
HELGA54	64	576374	AC016131	1383	1-399
HELGC24	65	576377	AC016927	1384	1-93 436-573 1504-1585 1922-1971 2635-2726 3541-3731 6137-6496 6517-6700 7141-7604 7621-8670 8705-9096 9766-10170 10425-10501 10987-11087 11341-11941 12244-15601 15897-16018
HELGC24	65	576377	AC024903	1385	1-244 278-784 2960-3209 6353-6483 6826-6963

					7892-7976 8310-8359 9023-9114 9928-10118 12523-12882 12903-13086 13527-13990 14007-15056 15091-15482 16152-16556 16811-16887 17373-17473 17727-18327 18630-21983 22279-22400
HELGC24	65	576377	AC016927	1386	1-442
HELGC24	65	576377	AC024903	1387	1-442
HELGC32	66	699375	AC010646	1388	1-147 327-996 1637-1802 1860-2209 2317-3026
HELGC32	66	699375	AC010646	1389	1-146
HELGC32	66	699375	AC010646	1390	1-425 665-988 1074-1240 1630-1763 2370-2873 3013-3759 3800-4750 5484-5900 6149-6404 7522-7653 7678-7822 9244-9344 9545-9845 10064-10093
HELGG77	68	825795	AC073531	1391	1-935
HELGQ48	72	851178	AL356982	1392	1-337
HELGQ48	72	851178	AL356982	1393	1-485
HELGQ48	72	851178	AL356982	1394	1-1124

10091504.030702

HELGT48	73	879483	AC012043	1395	1-647
HELGT48	73	879483	AC012043	1396	1-346
HELGZ48	74	721742	AC010357	1397	1-411
HELHB12	75	970863	AC026088	1398	1-417
HELHB12	75	970863	AC026088	1399	1-500
HELHB12	75	970863	AC026088	1400	1-170
HELHF07	78	949067	AC073669	1401	1-597
HELHF07	78	949067	AC023605	1402	1-583
HELHF07	78	949067	AC074220	1403	1-362
HELHF07	78	949067	AC074220	1404	1-105
HEMBC56	80	577797	AC016582	1405	1-384
HEMBC56	80	577797	AC008395	1406	1-384
HEMBC56	80	577797	AC016582	1407	1-584
HEMBC56	80	577797	AC016582	1408	1-441
HEMBC56	80	577797	AC008395	1409	1-584
HEMBC56	80	577797	AC008395	1410	1-441
HEMBI16	81	507220	AL354935	1411	1-218
HEMBI16	81	507220	AL161792	1412	1-218
HEMBI16	81	507220	AL354935	1413	1-357
HEMBI16	81	507220	AL161792	1414	1-358
HEMBZ84	82	527989	AC006425	1415	1-187 1153-1682 2086-2186 3494-5612 5887-6014 7903-8173 8656-8807
HEMBZ84	82	527989	AC006425	1416	1-480
HEMCM25	84	948738	AL390057	1417	1-484
HEMCM25	84	948738	AC016105	1418	1-484
HEMCM25	84	948738	AL390057	1419	1-179
HEMCM25	84	948738	AC016105	1420	1-215 219-1162
HEMDG56	85	715834	AL022311	1421	1-391 950-1644 3306-3805 3846-4050 5772-5953 6844-7026 7365-7583

					7689-7802 8673-8951 8981-9224 11685-11826 12396-12476 12580-12702 13304-13554 14057-15041
HEMDG56	85	715834	AL022311	1422	1-314
HEMDG56	85	715834	AL022311	1423	1-226
HEMDG83	86	576508	AC068870	1424	1-599
HEMDG83	86	576508	AC068870	1425	1-312
HEMDG83	86	576508	AC068870	1426	1-394
HEMDK92	87	574278	AC027225	1427	1-297
HEMDK92	87	574278	AF127577	1428	1-348
HEMEN63	93	578717	AC008964	1429	1-309
HEMEN63	93	578717	AC010633	1430	1-309
HEMFX20	97	840164	AC011468	1431	1-1022 1026-1219 1226-1529 2871-4241 5024-5251
HEMFX20	97	840164	AC011468	1432	1-465
HEMFX20	97	840164	AC011468	1433	1-492
HHBBK65	102	588062	AC069164	1434	1-436
HHBBL40	103	588066	AL136979	1435	1-200 1109-1226 1290-1574 2005-2748 2835-3402 3495-4093 4122-4398 4639-4787 6077-6182 8236-8462 10743-10887 11474-12059 12155-12358 14440-16057
HHBBL40	103	588066	AL136979	1436	1-225
HHBEM70	107	756949	AC025036	1437	1-148

HHBEM70	107	756949	AC011953	1438	1-126
HHBEM70	107	756949	AC027300	1439	1-158
HHBEN34	108	703745	AC068489	1440	1-383
HHBEN34	108	703745	AC068489	1441	1-127
HHBEN34	108	703745	AC068489	1442	1-546
HHBFL31	109	800035	AC069066	1443	1-295
HHBFL31	109	800035	AC011186	1444	1-295
HHBFL31	109	800035	AC069066	1445	1-195
HHBFL31	109	800035	AC011186	1446	1-186
HHBFW44	110	716283	AC026596	1447	1-263
HHBFW44	110	716283	AC026596	1448	1-545
HHBGN52	113	726391	AC007377	1449	1-648
HHBGY59	117	792027	AC023356	1450	1-230 1461-1708 3063-3429 3456-3928 5267-5599
HHBGY59	117	792027	AC023356	1451	1-1398 1771-2539 2583-3164
HHBHK84	120	858431	AL031719	1452	1-339
HHBHK84	120	858431	AC046159	1453	1-339
HHFBD39	123	826307	AC009595	1454	1-222
HHFBD39	123	826307	AC009595	1455	1-268
HHFBF32	125	502872	AC012482	1456	1-784
HHFBF32	125	502872	AC011200	1457	1-783
HHFBF32	125	502872	AC012482	1458	1-348
HHFBF32	125	502872	AC011200	1459	1-348
HHFBJ81	128	502954	AL133459	1460	1-344
HHFBJ81	128	502954	AC068460	1461	1-344
HHFBJ81	128	502954	AL133459	1462	1-333
HHFBJ81	128	502954	AC068460	1463	1-333
HHFBL16	129	509237	AC010336	1464	1-384
HHFBL16	129	509237	AC024700	1465	1-383
HHFBL16	129	509237	AC010336	1466	1-290
HHFBL30	130	509238	AL161790	1467	1-342
HHFBL30	130	509238	AC010741	1468	1-342
HHFBL30	130	509238	AL161790	1469	1-106
HHFBL30	130	509238	AC010741	1470	1-175
HHFBL30	130	509238	AC010741	1471	1-101



200504051600

HHFBL36	132	707930	AC010261	1472	1-406
HHFBL39	133	509637	AC005343	1473	1-96 784-1143 1734-1849 2133-2401 3742-5557
HHFBP60	135	503453	AC002126	1474	1-277
HHFBP60	135	503453	AC002126	1475	1-105 1581-1762 2134-2537 4582-4734 5112-5210 6383-6490 6553-6642 7270-7390 7534-7570 7840-7991 9058-9375 10822-10952 12795-13209 13221-13367 13467-13582 13723-13899 14027-14125 14701-15275
HHFBP60	135	503453	AC002126	1476	1-529
HHFCH59	139	526389	AC009474	1477	1-427
HHFCH59	139	526389	AC009474	1478	1-153
HHFCI73	140	518427	AC022919	1479	1-372
HHFCI73	140	518427	AC022919	1480	1-491
HHFCK71	141	781725	AC073990	1481	1-275 554-732 1216-1578 1703-4646
HHFCK71	141	781725	AC005829	1482	1-275 553-731 1218-1580 1989-4645
HHFCK71	141	781725	AC073990	1483	1-573
HHFCK71	141	781725	AC073990	1484	1-1043
HHFCK71	141	781725	AC005829	1485	1-573

2000-40516001

HHFCK71	141	781725	AC005829	1486	1-922
HHFCL91	142	509628	AC013355	1487	1-688
HHFCL91	142	509628	AC018926	1488	1-384 1804-2494
HHFCL91	142	509628	AC013355	1489	1-1408
HHFCP67	143	536062	AL353578	1490	1-295
HHFCP67	143	536062	AL353578	1491	1-473
HHFCZ01	144	509163	AC018939	1492	1-318
HHFCZ01	144	509163	AC016380	1493	1-317
HHFCZ01	144	509163	AL390755	1494	1-35 2258-2648 5778-5868 7978-8065 8456-8861 9396-10035 13564-13756 14340-14510 16437-16828 17616-17707 17794-17902 19502-19601 21470-21624 21859-21971 22649-22831 22959-23049 23854-24009 24167-24269 24379-24467 24748-24987 25534-25676 26138-26270 26365-26449 26893-26995 28121-28175 30159-30348 31129-31265 32058-32441 32580-33058 33663-33737 34588-34663 35890-36034

					37329-37432 37884-38392 38447-38665 39520-39704 40417-40767 41283-41815 44971-45126 45491-45610 47830-48146 48375-48564 50441-51935
HHFCZ01	144	509163	AC016380	1495	1-189 2067-3561
HHFDA67	145	509387	AC017000	1496	1-311
HHFDG32	146	502957	AC021453	1497	1-1671
HHFDG32	146	502957	AC021453	1498	1-117
HHFDG32	146	502957	AC021453	1499	1-677
HHFDH38	148	536551	AL136980	1500	1-331
HHFDH38	148	536551	AL136980	1501	1-661
HHFDI42	150	500877	AL359473	1502	1-442
HHFDI62	151	745569	AL354888	1503	1-1050
HHFDI62	151	745569	AL033527	1504	1-627
HHFDI62	151	745569	AL354888	1505	1-151
HHFDI62	151	745569	AL033527	1506	1-151
HHFDI62	151	745569	AL033527	1507	1-198
HHFDI66	152	573283	AC013723	1508	1-389
HHFDI66	152	573283	AC009973	1509	1-389
HHFDJ87	154	575105	AL158141	1510	1-341
HHFDJ87	154	575105	AL139111	1511	1-341
HHFDJ87	154	575105	AL158141	1512	1-532
HHFDJ87	154	575105	AL158141	1513	1-220
HHFDJ87	154	575105	AL139111	1514	1-532
HHFDJ87	154	575105	AL139111	1515	1-220
HHFEJ18	156	525616	AC027487	1516	1-1524
HHFEJ18	156	525616	AC068403	1517	1-1116
HHFEJ18	156	525616	AC006453	1518	1-2330
HHFEJ18	156	525616	AC005629	1519	1-2332
HHFEJ18	156	525616	AC068403	1520	1-1431
HHFEJ18	156	525616	AC027487	1521	1-129
HHFEJ18	156	525616	AC068403	1522	1-940

HHFEJ18	156	525616	AC006453	1523	1-174
HHFEJ18	156	525616	AC006453	1524	1-129
HHFEJ18	156	525616	AC005629	1525	1-182
HHFEJ18	156	525616	AC005629	1526	1-129
HHFES51	158	525609	AC022281	1527	1-171
HHFES51	158	525609	AC022281	1528	1-337
HHFES51	158	525609	AC022281	1529	1-322
HHFFG41	159	575009	AP002338	1530	1-412
HHFFG41	159	575009	AC073195	1531	1-412
HHFFG41	159	575009	AP002338	1532	1-95
HHFFG41	159	575009	AP002338	1533	1-173
HHFFG41	159	575009	AC073195	1534	1-209
HHFFO66	161	500910	AC068341	1535	1-49 72-448 1609-2003
HHFFO66	161	500910	AC068341	1536	1-338
HHFFT69	163	662462	AL359453	1537	1-1968
HHFFT69	163	662462	AL359096	1538	1-1968 2155-5508
HHFFT69	163	662462	AC053522	1539	1-1968 2155-5508
HHFFT69	163	662462	AC015973	1540	1-1968 2155-5508
HHFFT69	163	662462	AL359453	1541	1-1594
HHFFT69	163	662462	AL359096	1542	1-332
HHFFT69	163	662462	AL359096	1543	1-2319
HHFFT69	163	662462	AC053522	1544	1-2319
HHFFT69	163	662462	AC053522	1545	1-332
HHFFT69	163	662462	AC015973	1546	1-2318
HHFFT69	163	662462	AC015973	1547	1-332
HHFFX20	164	525617	AC021421	1548	1-112
HHFFX20	164	525617	AL133312	1549	1-644 777-888 1007-1671
HHFFX20	164	525617	AL135752	1550	1-644 777-888 1007-1671
HHFFX20	164	525617	AC021421	1551	1-203
HHFFX20	164	525617	AL133312	1552	1-355
HHFFX20	164	525617	AL133312	1553	1-290

20/06/04 10:51:00

HHFFX20	164	525617	AL135752	1554	1-290
HHFFX20	164	525617	AL135752	1555	1-355
HHFFX75	165	507394	Z93024	1556	1-113 552-1029 1177-1329 3299-3343 3350-3543 4372-4456 5937-6110 7289-7459 9374-9496 10624-10699 12785-13230 13572-13759 14164-14313 16011-16073 16995-17273 17279-17695 17765-18278 18343-19051 19109-19192 20461-20639 21437-21601 21866-21938 24242-24654 24773-26048
HHFFX75	165	507394	Z93024	1557	1-278
HHFFZ04	167	927869	AL132641	1558	1-4063 4990-5958
HHFFZ04	167	927869	AL132641	1559	1-2545
HHFGE01	174	917137	AC020723	1560	1-196 920-1040 1113-1520 1666-1933 2807-3264 3459-3729 3808-4062 4374-4487 4725-5161
HHFGE01	174	917137	AC020723	1561	1-2595
HHFGN31	177	908508	AC026271	1562	1-479

HHFGN31	177	908508	AC026964	1563	1-571
HHFGN31	177	908508	AC005324	1564	1-570
HHFGN31	177	908508	AC026271	1565	1-302
HHFGN31	177	908508	AC026271	1566	1-319
HHFGN31	177	908508	AC026964	1567	1-299
HHFGN31	177	908508	AC026964	1568	1-313
HHFGN31	177	908508	AC026964	1568	1769-2018
HHFGN31	177	908508	AC005324	1569	1-285
HHFGN31	177	908508	AC005324	1570	1-132
HHFGP71	179	573860	AL109916	1571	1-435
HHFGR35	180	573510	AC010687	1572	1-498
					694-956
					3087-3395
					3463-3725
					4192-4660
HHFGR35	180	573510	AC005369	1573	1-498
					694-956
					1721-1826
					3090-3398
					3466-3725
					4195-4664
					5068-5524
					7874-7984
					8636-8763
					9038-9626
					9670-10114
					10341-10441
					10646-10780
					10867-11561
					11674-12184
					12626-12746
					12931-13086
					13095-14152
HHFGR35	180	573510	AC010687	1574	1-112
HHFGR35	180	573510	AC005369	1575	1-112
HHFGR35	180	573510	AC005369	1576	1-260
HHFGS09	181	526331	AL356984	1577	1-237
					440-564
					1957-2091
					2278-3345
					3402-3454

					3523-4073
HHFGS09	181	526331	AL008733	1578	1-237 440-564 1957-2091 2278-3345 3402-3454 3523-4073
HHFGS09	181	526331	AL356984	1579	1-364
HHFGS09	181	526331	AL356984	1580	1-289 463-1174 2090-2428 3930-4781
HHFGS09	181	526331	AL008733	1581	1-289 463-1174 2090-2428 3930-4781
HHFGS09	181	526331	AL008733	1582	1-364
HHFGS40	182	888332	AC026173	1583	1-304
HHFGS40	182	888332	AC055829	1584	1-304
HHFGS40	182	888332	AC023176	1585	1-304
HHFGS40	182	888332	AC026173	1586	1-412
HHFGS40	182	888332	AC055829	1587	1-253
HHFGS40	182	888332	AC026173	1588	1-253
HHFGS40	182	888332	AC055829	1589	1-412
HHFGS40	182	888332	AC023176	1590	1-240
HHFGS40	182	888332	AC023176	1591	1-412
HHFGS92	183	871899	AC026055	1592	1-370
HHFGS92	183	871899	AC020565	1593	1-371
HHFGS92	183	871899	AC026055	1594	1-383
HHFGS92	183	871899	AC026055	1595	1-488
HHFGS92	183	871899	AC020565	1596	1-387
HHFGS92	183	871899	AC020565	1597	1-488
HHFGT10	184	968109	AC006312	1598	1-395 407-1052 2090-2221 2608-2705 3245-3347 3542-3934 4847-5734 5975-6244

					6329-6718 7054-7846 8300-9356 9438-10410 10940-11538
HHFGT10	184	968109	AC006312	1599	1-276 499-1675 2091-2333 4232-4282 4455-5232 6242-6970 7182-7481 7977-8098 8744-8899 9163-9247 10920-10984 12504-12581 13430-13629 14388-14696 15345-15477 16954-17086 17114-17792
HHFHA44	191	573498	AL049870	1600	1-262
HHFHA44	191	573498	AL049870	1601	1-300
HHFHA44	191	573498	AL049870	1602	1-399
HHFHC02	192	920510	AC005940	1603	1-463
HHFHC02	192	920510	AC005940	1604	1-1036
HHFHC57	194	573506	AC004084	1605	1-105 839-1021 2069-2302 2470-2855 3818-4265 4371-4610 4761-4810 5364-5802 5930-6517 7073-7807 8063-8618 8636-8875 9438-9537 10568-10774



10051504-030703

					10897-11025
					11718-12323
					13749-13849
					13978-14188
					14474-14554
					16489-16624
					16924-17019
					17239-17458
					17908-18185
					19014-19266
					19356-19451
					19620-19873
					19893-20920
					21092-21247
					21512-21579
					21621-21754
					22001-22831
					22992-23518
					23710-24370
					24426-24596
					25213-25493
					25661-26192
					26588-27433
					27598-27742
					28073-28199
					28359-28651
					28777-29249
					29379-29502
					29646-29794
					29833-30033
					30085-30630
					30702-32661
					33104-33374
					33383-33661
					33808-33871
					33978-37449
					37587-37754
					38296-38433
					38597-39343
					40047-40395
					40462-40743

					40938-41039 41187-41634 42504-42687 42703-42850 42929-43475
HHFHC57	194	573506	AC004084	1606	1-239
HHFHC57	194	573506	AC004084	1607	1-283
HHFHC72	195	766128	AC025422	1608	1-199
HHFHC72	195	766128	AC002110	1609	1-101
HHFHC72	195	766128	AC034119	1610	1-250
HHFHC72	195	766128	AC013817	1611	1-250
HHFHC72	195	766128	AC011175	1612	1-250
HHFHJ72	201	575030	AC073330	1613	1-470
HHFHJ72	201	575030	AC073330	1614	1-294
HHFHJ72	201	575030	AC073330	1615	1-144
HHFHQ86	204	572923	AL109936	1616	1-445
HHFHQ86	204	572923	AP001456	1617	1-406
HHFHX11	206	967321	AL022476	1618	1-414
HHFHX11	206	967321	Z82199	1619	1-414
HHFHX11	206	967321	AL022476	1620	1-329
HHFHX11	206	967321	Z82199	1621	1-329
HHFHX11	206	967321	AL022476	1622	1-364
HHFHX11	206	967321	Z82199	1623	1-364
HHFJL11	211	965931	AC026672	1624	1-414
HHFJL11	211	965931	AC026672	1625	1-309
HHFJM56	212	857990	AC005998	1626	1-278
HHFJM56	212	857990	AC005586	1627	1-280
HHFJM56	212	857990	AC005998	1628	1-404
HHFJM56	212	857990	AC005586	1629	1-411
HHFJN02	214	918358	AC073932	1630	1-423
HHFJN02	214	918358	AC007064	1631	1-440
HHFJN02	214	918358	AC009590	1632	1-423
HHFJN02	214	918358	AC010632	1633	1-453
HHFJN02	214	918358	AC007064	1634	1-361 466-3593
HHFJR06	217	934093	AC010538	1635	1-539
HHFJR06	217	934093	AC005360	1636	1-305 333-452 561-1005 1635-1717

HHFJR06	217	934093	AC010538	1637	1-120
HHFJR06	217	934093	AC010538	1638	1-195
HHFJR06	217	934093	AC005360	1639	1-412
HHFJX18	218	907658	AL354671	1640	1-433
HHFJX18	218	907658	AL354671	1641	1-247 432-558 1491-1606 1796-1903 2051-2210
HHFKC28	220	857948	AC013715	1642	1-395
HHFKC28	220	857948	AC013715	1643	1-328
HHFKC43	221	906903	AC007776	1644	1-475
HHFKC43	221	906903	AC007776	1645	1-105 3385-3585 4362-4516 4950-6638
HHFKC43	221	906903	AC007776	1646	1-482
HHFKH82	223	857810	AC000097	1647	1-74 524-3302 3495-3649 4115-5226 5235-5285 5492-5923 6531-7255 7397-7497 8597-10758
HHFKH82	223	857810	AC005664	1648	1-334
HHFKH82	223	857810	AC000075	1649	1-74 524-3302 3495-3649 4127-4430 4436-5227 5236-5286 5493-5924 6532-7256 7398-7498 8598-10759
HHFKH82	223	857810	AC006547	1650	1-74 524-3302 3495-3649 4115-5226

20200405T001

					5235-5285 5492-5923 6531-7255 7397-7497 8597-10758
HHFKH82	223	857810	AC000097	1651	1-1172
HHFKH82	223	857810	AC000075	1652	1-1175
HHFKH82	223	857810	AC006547	1653	1-1172
HHFKJ68	225	934045	AL161669	1654	1-101 266-417 3879-4182 5465-5877 7378-7616 9205-9314 9529-9652 10000-10103 10254-10332 12045-12162 12684-12807 12895-13047 13660-13779 14848-15425
HHFKK95	226	952082	AL357149	1655	1-795
HHFKU44	229	857923	AP000487	1656	1-63 252-919 1070-2004 2421-2533
HHFKU44	229	857923	AP000405	1657	1-132 801-1028 2210-2693 4235-4529 4710-4823 5012-5680 5831-6762 7179-7291 7766-8298 9901-10176 10267-11292 11651-12864 13121-13176
HHFKU44	229	857923	AP000487	1658	1-295

HHFKU44	229	857923	AP000405	1659	1-441
HHFKX11	230	965874	AC068889	1660	1-153 638-1551 1774-2060 2495-3040 3119-3469 3654-3773
HHFKX11	230	965874	AC021103	1661	1-914
HHFKX11	230	965874	AC073611	1662	1-913
HHFKX11	230	965874	AC023509	1663	1-121 607-1520 1743-2029 2043-3015 3094-3444 3629-3748
HHFKX11	230	965874	AC073611	1664	1-172
HHFKX11	230	965874	AC023509	1665	1-88
HHFKX11	230	965874	AC068889	1666	1-88
HHFKX11	230	965874	AC021103	1667	1-246
HHFKY11	231	965876	AC008676	1668	1-70 646-766 1274-1988 2066-2374 2827-3314 4686-5053 5561-5756 6591-6879 7931-8474 9066-9908 9916-10434
HHFLA69	232	933869	AL133347	1669	1-227
HHFLA69	232	933869	AC009814	1670	1-212
HHFLA69	232	933869	AC009814	1671	1-845
HHFLA69	232	933869	AL133347	1672	1-845
HHFLA69	232	933869	AC009814	1673	1-560
HHFLH34	233	857907	AC068866	1674	1-100
HHFLH34	233	857907	AL359878	1675	1-138
HHFLM06	234	933868	U51560	1676	1-432
HHFLM06	234	933868	Z80896	1677	1-432
HHFLM06	234	933868	U51560	1678	1-1118

HHFLM06	234	933868	U51560	1679	1-155
HHFLM06	234	933868	Z80896	1680	1-155
HHFLM06	234	933868	Z80896	1681	1-1118
HHFLU04	236	870085	AL121747	1682	1-118 250-451 1031-1281 1334-1617 5503-5588 6626-6746 11377-11537 12155-12776 13270-14743 15440-15693 18886-19237 19438-19846
HHFLU04	236	870085	AL034548	1683	1-118 250-451 1031-1281 1334-1617 5503-5588 6626-6746 11377-11537 12155-12776 13270-14743 15440-15693 18886-19237 19438-19846
HHFMB01	238	914951	AC062037	1684	1-396
HHFMB01	238	914951	AC062037	1685	1-130
HHFMB01	238	914951	AC062037	1686	1-316 545-1725 2109-2527 3231-3295
HHFME70	239	926497	AL035686	1687	1-383
HHFMH85	240	872838	AJ002553	1688	1-2930 3044-3177 4005-4035
HHFMH85	240	872838	AP000449	1689	1-124 200-376 462-650 689-4651

207020 49516001

					4762-4898 5724-5901 7362-7660 7953-8121
HHFMH85	240	872838	AC051660	1690	1-64 334-757 899-1050 1198-1599 1738-1891 2098-2392 2922-3049 3142-3244 3319-3389 3482-3598 3691-3857 4138-4515 4601-4789 4828-8792 8903-9039 9865-10042 11509-11797 12091-12259
HHFMH85	240	872838	AP000449	1691	1-1584
HHFMH85	240	872838	AC051660	1692	1-1585
HHFMO03	242	922708	AL139329	1693	1-145 1052-1713 1848-1873 3110-6111
HHFMO03	242	922708	AL139329	1694	1-1003
HHFMO03	242	922708	AL139329	1695	1-922
HHFMO94	243	957955	AC006172	1696	1-520
HHFMO94	243	957955	AC006172	1697	1-449
HHFMQ70	244	840039	AL390318	1698	1-335
HHFMQ70	244	840039	AC010161	1699	1-335
HHFMQ70	244	840039	AL390318	1700	1-764
HHFMQ70	244	840039	AC010161	1701	1-764
HHFMZ40	245	974299	AC025552	1702	1-837 1592-1876 1971-2505 2544-3080
HHFMZ40	245	974299	AC023085	1703	1-837

					1592-1876 1971-2510 2562-3083
HHFMZ40	245	974299	AC025552	1704	1-101
HHFMZ40	245	974299	AC023085	1705	1-101
HHFND10	247	963140	AL158844	1706	1-208 228-830
HHFND10	247	963140	AL158837	1707	1-208 228-677 706-803
HHFND10	247	963140	AC073965	1708	1-208 228-830
HHFND10	247	963140	AL158844	1709	1-126
HHFND10	247	963140	AC073965	1710	1-558
HHFND10	247	963140	AL158844	1711	1-560
HHFND10	247	963140	AC073965	1712	1-166
HHFNI49	249	857850	AC009364	1713	1-312
HHFNI49	249	857850	AC018643	1714	1-312
HHFOC27	250	857816	AL049835	1715	1-440
HHFOC27	250	857816	AL049835	1716	1-135
HHFOC27	250	857816	AL049835	1717	1-114
HHFOC66	251	857955	AC019220	1718	1-292 487-546
HHFOC66	251	857955	AC019220	1719	1-88
HHFOF07	252	952062	AL132987	1720	1-271 645-1063 1112-1370 1810-2112 2483-2590 2916-3117 3613-3945 4310-4376 4642-4756 4874-5203 5480-5850 6289-6892
HHFOF40	253	930839	AC012557	1721	1-527
HHFOF40	253	930839	AC063925	1722	1-527
HHFOF40	253	930839	AC012557	1723	1-705 857-979



					3465-3577 4502-4623 5624-6761
HHFOF40	253	930839	AC063925	1724	1-703
HHFOF40	253	930839	AC012557	1725	1-177 797-1153 1766-2651 2949-3031 8066-8473
HHFOF40	253	930839	AC063925	1726	1-206
HHFOJ05	254	930856	AC010172	1727	1-262
HHFOJ05	254	930856	AC000015	1728	1-262
HHFOJ05	254	930856	AC010172	1729	1-718
HHFOJ05	254	930856	AC010172	1730	1-273
HHFOJ05	254	930856	AC000015	1731	1-273
HHFOJ05	254	930856	AC000015	1732	1-718
HHFON08	255	957974	AL031282	1733	1-125 204-676 2747-6745 7623-8192 8629-8798 8860-9174 10053-10226 10735-11362 11633-11748 12345-12774 13014-13179 13685-13811 14285-14420 14599-15069 16162-16271 18792-18921 19616-20033 20905-21606
HHFON08	255	957974	AL031282	1734	1-1063 1094-1253 1441-1617 1658-1817 1840-1989 2152-2649 2805-2996

10061504-030702

					3259-3521 3693-5265 5800-5870 6414-6575 6657-7261 7558-7800 8352-8943 9884-10022 11068-11767 11920-12427 12724-13300 13754-14098 14157-14314 14362-14444 14556-14768 15311-15585 16256-16553 16790-17078 17330-17431 17894-18012 18663-18696 19219-19499 20187-20735 20926-21129
HHFON19	257	910891	AC003072	1735	1-395 1224-1311 2736-2904 3775-5829 6051-6406 6418-7537
HHFON19	257	910891	AC003072	1736	1-314
HHFON19	257	910891	AC003072	1737	1-384
HHFON32	258	858034	AC024555	1738	1-65 1322-1814 1879-2084 2488-3174 3624-4306 4653-4875 5942-7911
HHFON32	258	858034	AL356796	1739	1-493
HHFON32	258	858034	AL356796	1740	1-206

20/06/2014 10:05:00

					610-1296 1746-2428 2774-2996 4063-5628
HHFOW08	260	857772	AC013420	1741	1-459
HHFOW08	260	857772	AC067745	1742	1-458
HHFOW08	260	857772	AC067745	1743	1-353
HHFSB02	261	921364	AC022003	1744	1-503 1443-2058 2808-3134 4719-4770 5295-5710 5942-6141 7267-7809
HHFSB02	261	921364	AC024059	1745	1-503 1443-2057 2807-3133 4718-4769 5294-5709 5942-6141 7267-7809
HHFSB02	261	921364	AC024058	1746	1-616 1366-1692 3277-3328 3853-4268 4501-4700 5825-6055
HHFSB02	261	921364	AC024058	1747	1-81 187-272 932-1226 4756-4859 6139-6343 6965-7071 7760-7918 8877-8993
HHFSB02	261	921364	AC024058	1748	1-395
HHFTC88	262	500903	AL354887	1749	1-344
HHFTC88	262	500903	AL354886	1750	1-344
HHFTC88	262	500903	AL354878	1751	1-344
HHFTC88	262	500903	AL354886	1752	1-913
HHFTC88	262	500903	AL354887	1753	1-634

2020ED-4051600T

					819-1731
HHFTC88	262	500903	AL354878	1754	1-634 819-1731
HHFUB23	264	928063	AC004985	1755	1-473
HHFUB23	264	928063	AC004985	1756	1-465
HHFUB77	265	772691	AC055119	1757	1-320
HHFUB77	265	772691	AL162592	1758	1-300
HHFUB77	265	772691	AC027170	1759	1-292
HHFUB77	265	772691	AC002369	1760	1-586 2559-2651 3329-3426 3756-5088
HHFUB77	265	772691	AC027170	1761	1-226
HHFUB77	265	772691	AC002369	1762	1-228
HHFUC24	266	524840	AC074086	1763	1-483
HHFUC24	266	524840	AC074086	1764	1-451
HHFUC26	267	960331	AC007686	1765	1-500
HHFUC26	267	960331	AC007686	1766	1-281
HHFUC26	267	960331	AC007686	1767	1-98
HHFUC42	268	525603	AC004836	1768	1-209 5804-6076 6127-7676 7693-8144 9341-9450 9803-10218 10270-11045 12112-12455 15673-15937 16059-16524 26709-26743 28687-29151 29659-29985 30110-30352
HHFUC42	268	525603	AC004836	1769	1-584
HHFUC47	270	525599	AC069489	1770	1-547
HHFUC47	270	525599	AC008421	1771	1-599
HHFUC47	270	525599	AC010232	1772	1-599
HHFUC47	270	525599	AP002016	1773	1-597
HHFUC47	270	525599	AC069489	1774	1-741
HHFUC47	270	525599	AC008421	1775	1-740

HHFUC47	270	525599	AC010232	1776	1-741
HHFUC47	270	525599	AP002016	1777	1-741
HHFUC47	270	525599	AP002016	1778	1-102
HHFUC92	272	507018	AL390734	1779	1-136 625-782 902-1290 2223-2453 2703-3266 4648-4818 5418-5819
HHFUC92	272	507018	AL390734	1780	1-398
HHFUK58	273	525604	AC024585	1781	1-458
HHFUK58	273	525604	AC008474	1782	1-149 303-522 1014-1051 1640-2300 2318-2347 2671-2899 3010-3446 3656-4506 4589-6257
HHFUK58	273	525604	AC024585	1783	1-285
HHFUK58	273	525604	AC008474	1784	1-5710
HHFUN56	275	530953	AC008167	1785	1-825 2446-2521
HHFUN56	275	530953	AC008167	1786	1-449
HHFUN56	275	530953	AC008167	1787	1-304
HMEBA75	276	530303	AC010463	1788	1-56 380-483 579-900 1742-1988
HMEBA75	276	530303	AC010463	1789	1-27 934-1247 2557-2685 2850-2960
HMEBY61	277	947868	AC069364	1790	1-668 846-1170 2965-3152 4672-4870 8016-8176 10842-11293

10091504.030702

					12213-12434 12438-13156 13229-13453 13476-13904 13964-14507 14913-15044
HMEBY61	277	947868	AC009886	1791	1-668 846-1170 2965-3152 4672-4870 8016-8176 10843-11294 12214-12435 12439-13157 13230-13454 13477-13905 13965-14508 14913-15046
HMEBY61	277	947868	AC069364	1792	1-505
HMEBY61	277	947868	AC009886	1793	1-541
HMEEX61	281	742085	AP000779	1794	1-501
HMEEX61	281	742085	AP000848	1795	1-502
HMEEX61	281	742085	AP000612	1796	1-501
HMEEX61	281	742085	AC011875	1797	1-501
HMEEX61	281	742085	AP000779	1798	1-985
HMEEX61	281	742085	AP000848	1799	1-985
HMEEX61	281	742085	AP000612	1800	1-985
HMEEX61	281	742085	AC011875	1801	1-816 908-1405 1931-3042
HMEGF48	285	573821	AL139183	1802	1-409
HMEGF48	285	573821	AL139183	1803	1-278
HMEGG44	286	796421	AC068938	1804	1-3356
HMEGG44	286	796421	AC073407	1805	1-3357
HMEGG44	286	796421	AC027054	1806	1-3223
HMEGG44	286	796421	AC073407	1807	1-2027 2038-3487
HMEGG44	286	796421	AC068938	1808	1-2027 2038-3487
HMEGG44	286	796421	AC027054	1809	1-2027 2038-3487

20200940576001

HMEGG44	286	796421	AC073407	1810	1-323
HMEGI07	288	953815	AC020925	1811	1-141 2805-2932 3116-3196 3786-4021 4066-4397 4614-4799 5738-9513
HMEGI07	288	953815	AC008572	1812	1-39 1415-1510 1913-2053 4717-4851 5028-5109 5699-5934 5979-6388 6528-6713 7652-11427
HMEIU49	291	722988	AC009549	1813	1-374
HMEJD13	293	657231	AC015885	1814	1-267
HMEJD13	293	657231	AC041002	1815	1-267
HMEJD13	293	657231	AC041002	1816	1-267
HMEJD13	293	657231	AC041002	1817	1-267
HMEJD13	293	657231	AC074339	1818	1-266
HMEJD13	293	657231	AC027821	1819	1-50 213-300 859-1126 1195-1903 2104-2487 2628-2737 2791-3020 3064-3182 3190-4185 4433-5145 5857-5953 6331-7971 7973-8232 8452-8515
HMEJD13	293	657231	AC041002	1820	1-267
HMEJD13	293	657231	AC036181	1821	1-84 141-370 414-532

10094504-03000

					540-1535 1776-2488 3200-3296 3674-5593 5809-6696
HMEJD13	293	657231	AC027821	1822	1-267
HMEJD13	293	657231	AC024631	1823	1-267
HMEJD13	293	657231	AC036181	1824	1-524 687-774 1333-1600 1669-2377 2578-2962 3103-3212 3266-3495 3539-3657 3665-4659 4905-5617 6332-6428 6806-8442 8444-8703 8918-9805
HMEJD13	293	657231	AC023133	1825	1-524 688-775 1333-1600 1669-2321 2574-2957 3098-3207 3261-3490 3534-3652 3660-4655 4896-5608 6320-6416 6792-8432 8434-8693 8908-9796
HMEJD13	293	657231	AC067923	1826	1-104 267-354 913-1180 1249-1956 2157-2540 2681-2790



20200740515001

					2844-3073 3117-3235 3243-4238 4484-5196 5908-6004 6382-8301 8517-9404
HMEJD13	293	657231	AC005702	1827	1-524 687-774 1334-1602 1671-2379 2640-2936 3104-3210 3268-3497 3541-3954 4023-4621 4863-5575 5800-6163 6288-6384 6762-8411 8418-8685 8901-9348 9368-9831
HMEJD13	293	657231	AC021317	1828	1-88 802-898 1283-1346 1808-1910 2027-3195 3411-3979
HMEJD13	293	657231	AC003976	1829	1-524 688-775 1334-1601 1670-2378 2579-2962 3103-3212 3266-3495 3539-3657 3665-4659 4905-5617 6332-6428 6806-8442

20070501 105700

					8444-8703 8918-9804
HMEJD13	293	657231	AC036181	1830	1-358
HMEJD13	293	657231	AC027821	1831	1-425 519-886
HMEJD13	293	657231	AC024631	1832	1-425 519-886
HMEJD13	293	657231	AC036181	1833	1-371
HMEJD13	293	657231	AC036181	1834	1-371
HMEJD13	293	657231	AC023133	1835	1-243
HMEJD13	293	657231	AC036181	1836	1-243
HMEJD13	293	657231	AC023133	1837	1-371
HMEJD13	293	657231	AC067923	1838	1-371
HMEJD13	293	657231	AC005702	1839	1-242
HMEJD13	293	657231	AC005702	1840	1-400
HMEJD13	293	657231	AC021317	1841	1-125
HMEJD13	293	657231	AC021317	1842	1-87
HMEJD13	293	657231	AC003976	1843	1-371
HMEJD13	293	657231	AC003976	1844	1-243
HMEJJ84	294	781983	AC024428	1845	1-292
HMEJU60	295	740392	AC024306	1846	1-320
HMEJU60	295	740392	AC024306	1847	1-116
HMEJU60	295	740392	AC024306	1848	1-111
HMEKS76	298	767517	AC006960	1849	1-288 1456-2100 4234-4427 4782-4885 6074-6172 9477-9608 10804-10832 11470-11880 12966-13201 13885-14036 14317-14467 15577-15714 16355-16531 17539-17670 18267-18314 18407-18465 19024-19141

10091504.030702

					19425-19452 22505-22657 26100-26399 29294-29632 32949-33012 34781-34897 34915-35365 35388-35739 36534-36674 37470-37581 37656-37790 37917-38013 38911-39273 41140-41375 43201-43560 45322-45733 46172-47513
HMEKS76	298	767517	AC008266	1850	1-288 1456-2100 4234-4427 4782-4885 6080-6178 9481-9612 10808-10836 11474-11884 12970-13069
HMEKX51	299	727154	AC016747	1851	1-317
HMEKX51	299	727154	AC074249	1852	1-317
HMEKX51	299	727154	AC016747	1853	1-303
HMEKX51	299	727154	AC074249	1854	1-303
HMELC56	301	745773	AL160286	1855	1-316
HMELC56	301	745773	AL160286	1856	1-106
HMELR10	303	964629	AC026364	1857	1-331
HMELR10	303	964629	AC026364	1858	1-536
HMELV19	305	668665	AL355433	1859	1-200 245-325
HMELV19	305	668665	AL355433	1860	1-127
HULAX31	308	868930	AC008738	1861	1-451
HULAX31	308	868930	AC008744	1862	1-451
HULAX31	308	868930	AC008738	1863	1-173
HULAX31	308	868930	AC008744	1864	1-173

HUSAY26	315	527909	AC008758	1865	1-2593
HUSAY26	315	527909	AC008758	1866	1-87
HUSCA57	316	527800	AC026785	1867	1-259
HUSCA57	316	527800	AC026504	1868	1-259
HUSCD26	317	527799	AC011919	1869	1-362
HUSCD26	317	527799	AC013271	1870	1-362
HUSCD26	317	527799	AC020990	1871	1-362
HUSCD26	317	527799	AC011919	1872	1-114
HUSCD26	317	527799	AC011919	1873	1-761
HUSCD26	317	527799	AC013271	1874	1-761
HUSCD26	317	527799	AC020990	1875	1-761
HUSCD26	317	527799	AC013271	1876	1-114
HUSCD26	317	527799	AC020990	1877	1-114
HUSGH88	320	871776	AC069193	1878	1-730 1039-5690
HUSGH88	320	871776	AL049629	1879	1-730 1039-5689
HUSGH88	320	871776	AC069193	1880	1-2271
HUSGH88	320	871776	AL049629	1881	1-2271 8698-8799 9671-9808 10133-10904 13707-14024
HUSHH64	323	576459	AC008623	1882	1-646
HUSHH64	323	576459	AC008623	1883	1-126
HUSHH64	323	576459	AC008623	1884	1-45 1042-1363 2498-2968 2989-3046 3584-3866 3894-4206
HUSHJ55	324	576381	AL358252	1885	1-331 577-603 1059-4966
HUSHJ55	324	576381	Z98044	1886	1-337 577-603 1060-7238
HUSHL83	325	868883	AC012363	1887	1-3333
HUSHL83	325	868883	AC012363	1888	1-181
HUSIA43	327	575768	AC015544	1889	1-931

20200505 10516001

HUSIA43	327	575768	AC015544	1890	1-117
HUSIA43	327	575768	AC015544	1891	1-339
HUSIF23	328	862494	AL035681	1892	1-67 1943-2237 3321-3500 3934-4132 4463-5199 6360-7092 7146-7569 7910-8139 8867-9089 10043-10216 11334-11457 11871-12002 12058-12143 13195-13465 13653-13857 15161-15304 17040-17157 17748-18435 18710-18835 20818-21106 21357-23130
HUSIF23	328	862494	AL035681	1893	1-90 752-1239
HUSIS60	329	727153	AP001488	1894	1-141 722-819 5028-5139
HUSIS60	329	727153	AP001488	1895	1-364
HUSJW78	331	772956	AC006425	1896	1-394
HUSYA63	335	928021	AC024423	1897	1-156 626-1487
HUSYA63	335	928021	AC025019	1898	1-156 626-1487
HUSYA63	335	928021	AC012202	1899	1-156 626-1487
HUSYB16	336	868843	AC008975	1900	1-425
HUSYB16	336	868843	AC008975	1901	1-542
HUSYO46	338	868827	AL158830	1902	1-656 1895-2173 4714-4962

10091504-030702

HUSYO46	338	868827	AL138834	1903	1-656 1895-2173 4714-4962
HUSYO46	338	868827	AL158830	1904	1-151
HUSYO46	338	868827	AL138834	1905	1-151
HUSZV72	340	851170	AC007962	1906	1-912
HUSZV72	340	851170	AC007962	1907	1-2776
HUSYX03	342	922840	AC023659	1908	1-126 217-777 2352-2507 3465-3709 3896-3995 4541-4906 5086-7803
HUSYX03	342	922840	AC009303	1909	1-126 217-777 2350-2506 3464-3708 3895-3994 4540-4905 5085-7803
HUSYN33	344	651293	AC008435	1910	1-828 1842-3253
HUSYN33	344	651293	AC008435	1911	1-843
HUSYN33	344	651293	AC008435	1912	1-1640
HUSYI13	348	657288	AC074043	1913	1-1316
HUSYI13	348	657288	AC009651	1914	1-1773
HUSYI13	348	657288	AC074043	1915	1-493
HUSYI13	348	657288	AC009651	1916	1-424
HUSYI13	348	657288	AC009651	1917	1-493
HUSYG24	349	677258	AC068373	1918	1-500 537-1234
HUSYG24	349	677258	AC068373	1919	1-1086
HUSYF74	350	554723	AL136531	1920	1-511
HUSYF74	350	554723	AL136531	1921	1-1457
HUSYF74	350	554723	AL136531	1922	1-651 825-1223 1944-2538 2613-2793 3366-3657 4544-5302

					7419-7761 8345-8448 9618-9689 10389-10461 15614-15734
HUSYA27	351	934423	AL139413	1923	1-1045
HUSYA27	351	934423	AL138789	1924	1-116 1044-1320 1755-2624 4375-4548 4612-5585 6092-6277 6751-8806 8868-9912
HUSYA27	351	934423	AL138789	1925	1-445
HUSXW61	352	741856	AC006425	1926	1-643
HUSXW61	352	741856	AC006425	1927	1-248 2024-2526 3079-3186 3565-4006 4406-4849 4944-6043 6188-7728
HUSXH57	355	859907	AC011890	1928	1-943 1079-1636 2154-2473 3555-4008 4292-4439 6963-7154 8254-8537 8592-8985
HUSXH57	355	859907	AC011890	1929	1-478
HUSKA86	356	784887	AC069351	1930	1-1358 1586-1688 2017-2092
HUSKA65	357	868860	AC021811	1931	1-1031
HUSKA65	357	868860	AC021811	1932	1-210
HUSKA65	357	868860	AC021811	1933	1-470
HUSJW03	358	923035	AC072049	1934	1-778
HUSJW03	358	923035	AC072049	1935	1-325
HUSJW03	358	923035	AC072049	1936	1-404

2020040516001

HUSIR04	364	709228	AC016274	1937	1-686
HUSIN12	365	970759	AL356489	1938	1-497
HUSIN12	365	970759	AL139008	1939	1-497
HUSIN12	365	970759	AL356489	1940	1-284
HUSIN12	365	970759	AL356489	1941	1-551
HUSIN12	365	970759	AL139008	1942	1-284
HUSIE95	366	967176	AC008622	1943	1-1265 2254-2644 2731-2835 2905-3445 3608-3796 5301-5425 5825-5873 5966-6142 6830-7216 7946-8063 8386-8503 8800-8850 9112-9273 11226-11369 13533-13731 14588-14790 15163-15458 16207-16462 17318-17398 18661-18734 19282-19345
HUSIE95	366	967176	AC025983	1944	1-1265 2254-2644 2731-2835 2905-3445 3608-3796 5296-5420 5820-5868 5961-6137 6825-7210 7940-8057 8379-8496 8793-8843 9105-9266 11219-11362



2009-04-05 16:00

					13525-13723 14580-14782 15155-15450 16199-16454 17308-17388 18651-18724 19271-19334
HUSIE95	366	967176	AC008635	1945	1-1265 2254-2644 2731-2835 2905-3445 3608-3796 5301-5425 5825-5873 5966-6142 6830-7216 7946-8063 8386-8503 8800-8850 9112-9273 11226-11369 13532-13730 14587-14789 15162-15449 16206-16461 17317-17397
HUSIE18	367	666523	AC012202	1946	1-390
HUSIE18	367	666523	AC025019	1947	1-390
HUSIE18	367	666523	AC012202	1948	1-291
HUSIE18	367	666523	AC025019	1949	1-291
HUSIE18	367	666523	AC012202	1950	1-648
HUSIE18	367	666523	AC025019	1951	1-648
HUSHL86	369	960355	AC024060	1952	1-34 629-1217 1422-1504 1960-2505 2765-4239 4518-6834 7022-7142 7275-8801 9628-9839

20200405T001

					10556-11423 11545-11672 12454-12814 15287-15399 15533-15745 16789-17216 18802-18998
HUSHL86	369	960355	AC020751	1953	1-441 2592-2921 3789-4244 4432-4539 5444-5647 6379-6482 6779-7519 8350-8894 9797-10391 10434-11002 11220-11907 11916-12371 13221-14147 21571-22104 23458-23491 24086-24674 24879-24961 25417-25962 26222-30311 30499-32297 33124-33335 34052-34919 35041-35168 35950-36309 38826-38946 39104-39176
HUSHL86	369	960355	AC068314	1954	1-441 2592-2921 3789-4244 4432-4539 5444-5647 6379-6482 6779-7519 8350-8894

20200740916001

					9797-10391 10434-11002 11220-11907 11916-12371 13221-14147 21571-22104 23458-23491 24086-24674 24879-24961 25417-25962 26222-30311 30499-32297 33124-33275
HUSHL86	369	960355	AC018812	1955	1-441 2592-2921 3789-4244 4431-4538 5439-5642 6374-6477 6774-7514 8345-8889 9792-10386 10429-10997 11215-11901 11910-12365 13215-14141 21566-22099 23453-23486 24081-24669 24874-24956 25411-25956 26216-30308 30496-32294 33121-33332 34049-34915 35037-35164 35946-36304 38777-38889 39023-39235 40280-40707 42292-42488

2025040515001

HUSHL86	369	960355	AC024060	1956	1-534
HUSHL86	369	960355	AC068314	1957	1-191
HUSHE34	370	703409	AC021801	1958	1-519
HUSHE34	370	703409	AC009627	1959	1-559
HUSHE34	370	703409	AC009627	1960	1-505 651-1073
HUSHB60	372	746560	AL136109	1961	1-3030 3212-3814 9089-9408 11724-12242 12308-12959 23153-23254 24314-26013
HUSHB60	372	746560	AC023147	1962	1-3041 3223-3825 9096-9415 11730-12248 12314-12965 23152-23259 24319-26018
HUSHB60	372	746560	AC022219	1963	1-3030 3212-3814 9084-9403 11717-12235 12301-12952 23145-23246 24307-26006
HUSHB60	372	746560	AL136109	1964	1-344
HUSHB60	372	746560	AL136109	1965	1-174
HUSHB60	372	746560	AC023147	1966	1-122
HUSHB60	372	746560	AC022219	1967	1-191
HUSHB60	372	746560	AC022219	1968	1-344
HUSGW06	373	935574	AL138902	1969	1-723
HUSGW06	373	935574	AL031431	1970	1-723
HUSGW06	373	935574	AL031431	1971	1-65 2139-2595 4317-4509
HUSGW06	373	935574	AL138902	1972	1-457
HUSGW06	373	935574	AL031431	1973	1-102
HUSGT01	376	916620	AC009549	1974	1-562
HUSGT01	376	916620	AC015805	1975	1-562

202504051500

HUSGS35	377	707777	AL121991	1976	1-76 193-325 1939-2401 2845-2988 3059-3370
HUSGS35	377	707777	AC022307	1977	1-76 193-325 1940-2402 2846-2989 3059-3370
HUSGS35	377	707777	AL121991	1978	1-463
HUSGS35	377	707777	AC022307	1979	1-463
HUSGS35	377	707777	AC022307	1980	1-825
HUSGQ62	378	745727	AC005021	1981	1-1276 2698-3169 3411-4696 4739-5868 6038-6892 7133-8106 9050-9660 10094-10596 11421-12539 12909-13168 13283-15173
HUSGQ62	378	745727	AC005021	1982	1-1198
HUSGQ62	378	745727	AC005021	1983	1-682
HUSGH09	381	625647	AC022760	1984	1-858
HUSGH09	381	625647	AC022760	1985	1-406
HUSGH09	381	625647	AC022760	1986	1-1117 1273-3564
HUSGF79	382	775309	AL135924	1987	1-1865 2137-3262
HUSGF79	382	775309	AL135924	1988	1-242
HUSGF79	382	775309	AL135924	1989	1-752
HUSGF10	384	964844	AC068343	1990	1-692
HUSGF10	384	964844	AC016701	1991	1-568
HUSGF10	384	964844	AC068343	1992	1-88
HUSGF10	384	964844	AC016701	1993	1-88
HUSGB36	386	572924	AC004099	1994	1-469
HUSGB36	386	572924	AC004099	1995	1-98 239-606

2004-03-09

					1233-1898 2052-2151 2509-2746 3158-3664 4346-4810
HUSGB36	386	572924	AC004099	1996	1-459
HUSGB01	387	916804	AL109824	1997	1-2092 2686-3225 5694-5765 6067-6501 9860-9894 10118-10417 11352-11782 12468-12905 13228-13325 13732-14551 15046-15490 17951-18365 21892-22312 27084-27681
HUSGB01	387	916804	AL118520	1998	1-2092 2686-3225 5694-5765 6067-6501 6800-6936 9860-9894 10118-10417 11365-11782 12468-12905 13228-13325 13732-14095 14134-14551 15046-15490 17951-18365 21892-22312 27084-27681
HUSGB01	387	916804	AL109824	1999	1-3383 3917-3977
HUSGB01	387	916804	AL109824	2000	1-498
HUSGB01	387	916804	AL118520	2001	1-3383 3917-3977

HUSGB01	387	916804	AL118520	2002	1-498
HUSFH89	388	786970	AL161787	2003	1-518
HUSFH89	388	786970	AC007847	2004	1-518
HUSFH89	388	786970	AL161787	2005	1-378
HUSFH89	388	786970	AC007847	2006	1-430
HUSFH89	388	786970	AL161787	2007	1-430
HUSFH89	388	786970	AC007847	2008	1-383
HUSFE05	390	932106	AC009754	2009	1-809 1261-2223
HUSFE05	390	932106	AC009754	2010	1-165
HUSDA09	391	461656	AC010310	2011	1-481 1208-1257 2679-2913 3250-3292 3997-4495
HUSDA09	391	461656	AC010358	2012	1-481 1208-1257 2678-2912 3249-3291 3996-4494
HUSDA09	391	461656	AC010308	2013	1-481
HUSDA09	391	461656	AC010473	2014	1-481 1208-1257 2678-2912 3249-3291 3996-4409
HUSDA09	391	461656	AC010310	2015	1-764 1731-1852 2379-2680 6120-6247 6688-6920
HUSDA09	391	461656	AC010358	2016	1-763 1730-1851 2382-2676 4434-4521 6687-6932 10024-10108 10183-10309 11384-11479 12563-12667 13757-14058

20040516001

					14653-14751 15469-15566 17367-17421 19200-19434 19799-19996 20868-21361 22650-23107 23645-23886 25147-25272 26153-26451 27411-27623 28475-28614 29767-29903 30328-30783 32311-32724 32838-33033 34513-36008
HUSDA09	391	461656	AC010473	2017	1-763 1730-1851 2386-2680 4598-4711 6687-6932 10021-10312 11383-11478 12562-12666 13756-14057 14655-14753 15471-15568 17375-17429 19207-19441 19806-20003 20875-21368 22656-23110 23649-23887 25151-25276 26157-26455 27002-27124 27415-27627 28479-28618 29771-29907 30332-30787



20200740516001

					32010-32222 32315-32728 32842-33037 34517-36012 36375-36754 36988-37175 37498-37988
HUSAM87	394	529783	AC027372	2018	1-207
HUSAM22	396	523674	AL353597	2019	1-254
HUSAM22	396	523674	AL353597	2020	1-402
HUSAM22	396	523674	AL353597	2021	1-358
HUSAL04	397	927719	AL162492	2022	1-399
HUSAL04	397	927719	AC004547	2023	1-326
HUSAL04	397	927719	AL162492	2024	1-108
HUSAL04	397	927719	AL162492	2025	1-435
HUSAL04	397	927719	AC004547	2026	1-434
HUSAL04	397	927719	AC004547	2027	1-108
HUSAJ57	398	678932	AC026432	2028	1-399
HUSAJ57	398	678932	AC027332	2029	1-399
HUSAJ57	398	678932	AC026432	2030	1-655
HUSAJ57	398	678932	AC027332	2031	1-655
HMELM86	404	784702	AL138904	2032	1-296
HMELM86	404	784702	AL354990	2033	1-296
HMELI57	407	734769	AL354988	2034	1-165 1557-1835 1925-2828 3023-4548
HMELI57	407	734769	AL136382	2035	1-548 2077-2183 2834-3263 4272-5297 5304-5714 5740-6077 6112-6209 6275-6444 7836-8113 8204-9110 9370-11724 11925-12566
HMELI57	407	734769	AL354988	2036	1-338

HMEI157	407	734769	AL136382	2037	1-297
HMEI157	407	734769	AL136382	2038	1-788
HMEKQ19	411	668659	AC034295	2039	1-1147
HMEKO03	412	924172	AC017079	2040	1-532 831-2091 3985-4071
HMEKO03	412	924172	AC017079	2041	1-87
HMEKJ40	413	711187	AC076959	2042	1-328
HMEKJ40	413	711187	AC026342	2043	1-328
HMEKJ40	413	711187	AC022336	2044	1-328
HMEKC72	415	760637	AC073177	2045	1-699
HMEKC72	415	760637	AC016905	2046	1-699
HMEKC72	415	760637	AC073177	2047	1-440
HMEKC72	415	760637	AC073177	2048	1-361
HMEKC72	415	760637	AC016905	2049	1-440
HMEJW50	416	724396	AC021412	2050	1-301
HMEJW50	416	724396	AC021420	2051	1-314
HMEJW50	416	724396	AC018603	2052	1-314
HMEJW50	416	724396	AC021420	2053	1-417 425-556 573-681 1097-1351 2554-2660 2815-3022
HMEJW50	416	724396	AC018603	2054	1-395
HMEJF25	418	678131	AC018805	2055	1-438
HMEJF25	418	678131	AC061707	2056	1-438 580-1572
HMEJF25	418	678131	AC018805	2057	1-153
HMEJF25	418	678131	AC061707	2058	1-174
HMEJF25	418	678131	AC018805	2059	1-132
HMEJF25	418	678131	AC061707	2060	1-153
HMEIV22	419	674611	AL109923	2061	1-839
HMEIV22	419	674611	AL109923	2062	1-531
HMEIV22	419	674611	AL109923	2063	1-108
HMEGK14	422	796443	AC025552	2064	1-143 194-465 672-978 1683-3496 5420-5721

10091504-030702

					6695-7437 7667-7813 8044-8212 8296-8435 9411-9724 9728-10374 11393-12347 12818-13273 14310-14466 15836-16511
HMEGK14	422	796443	AC025552	2065	1-224
HMEGH92	423	790629	AL139412	2066	1-79 1102-1268 1574-1674 2124-2233
HMEGH92	423	790629	AL390756	2067	1-79 1110-1276 1582-1682 2131-2240
HMEGH92	423	790629	AL390756	2068	1-107
HMEGH92	423	790629	AL139412	2069	1-95
HMEFD72	424	766185	AC023886	2070	1-625
HMEFD72	424	766185	AC023886	2071	1-436
HMEEL38	425	733649	AC011934	2072	1-777 1512-1768 2556-3544 3903-4196 4251-4362 5124-6394 6496-6623 6795-7436 7955-7995 8050-9669
HMEEL38	425	733649	AC025287	2073	1-41 96-868
HMEEL38	425	733649	AC011934	2074	1-166
HMEDR76	426	529897	AC002357	2075	1-172
HMEDR76	426	529897	AC002357	2076	1-248
HMECH43	428	715568	AL161779	2077	1-299
HMECH43	428	715568	AC004390	2078	1-299
HMECH43	428	715568	AL161779	2079	1-722

20200405001

					1211-3695
HMECH43	428	715568	AL161779	2080	1-450
HMECH43	428	715568	AC004390	2081	1-450
HMECH43	428	715568	AC004390	2082	1-722 1211-3695
HMEBY95	429	796058	AC026406	2083	1-457 622-1235
HMEBY95	429	796058	AC012609	2084	1-457 622-1235
HMEBY95	429	796058	AC026406	2085	1-427
HMEBY95	429	796058	AC012609	2086	1-276
HMEBY95	429	796058	AC012609	2087	1-427
HMEBG01	430	921763	AC009822	2088	1-3423 3514-3705 3918-7347
HMEBG01	430	921763	AC019299	2089	1-2183 2274-2465 2678-6107
HMEBG01	430	921763	AC025918	2090	1-3424 3515-3706 3921-7351
HMEAN12	431	655220	AC018850	2091	1-2639
HMEAN12	431	655220	AC012057	2092	1-2640
HMEAN12	431	655220	AC012342	2093	1-2447
HMEAN12	431	655220	AC018850	2094	1-585
HMEAN12	431	655220	AC012057	2095	1-585
HMEAN12	431	655220	AC012342	2096	1-585
HMEAH31	433	698403	AL162262	2097	1-603
HMEAH31	433	698403	AC026577	2098	1-603
HMEAE24	434	880925	AC009095	2099	1-229 833-1956 2114-2687
HMEAE24	434	880925	AC010530	2100	1-229 833-1956 2114-2687
HMEAE24	434	880925	AC020827	2101	1-161
HMEAE24	434	880925	AC009095	2102	1-84
HMEAE24	434	880925	AC009095	2103	1-881
HMEAE24	434	880925	AC010530	2104	1-881
HMEAE24	434	880925	AC010530	2105	1-84

2020074051600T

HMEAE24	434	880925	AC020827	2106	1-435
HMEAE01	435	916744	AC069380	2107	1-238
HMEAE01	435	916744	AL162583	2108	1-130
HHFUB83	439	800580	AC015651	2109	1-2097 5308-5495 5696-5742 5890-6249 7370-7525 7850-8236 8359-8463 8597-8770 8919-9028 9213-9353 9517-9639 9765-9874 9944-11023 11124-11219 11315-11613 11708-12241 12431-12666 12744-12802 12976-13087 13374-13914 14728-15500
HHFUB83	439	800580	AC015651	2110	1-35 159-252 410-783 786-830 953-1035 1452-1553 1651-2071 2161-2264 2352-2454 2494-2758 2847-3006 3135-3272 3477-4138 4907-5738 5972-6059 6132-6367 6650-6834

20040504 0300

					6915-7010 7091-7658 7662-9457 10122-10222 11415-11534 12386-12418 13253-13584 13635-13867 14881-15326 15851-16013 16529-16816 17430-17529 18140-18269 18634-18734 19189-19369 20434-21105 21912-22008
HHFOK10	442	961346	AC006995	2111	1-114 459-1252 1604-1917 2187-2469 3571-3755 4170-4602 4741-4841 5040-5082 5857-5958 6253-6340 7112-8041 8231-8368 8774-10894
HHFOK10	442	961346	AC005098	2112	1-114 553-988 1601-1926 2188-2470 3568-3753 4168-4600 5038-5080 5881-5976 6271-6358 7128-8058 8786-10907

HHFOK10	442	961346	AC004166	2113	1-77 541-951 1564-1889 2151-2235 3531-3716 4131-4563 4702-4802 5001-5043 5844-5939 6234-6321 7115-7476 7548-8020 8745-10867
HHFOK10	442	961346	AC006995	2114	1-89
HHFOK10	442	961346	AC005098	2115	1-509
HHFOK10	442	961346	AC004166	2116	1-509
HHFNJ05	443	930899	AC015980	2117	1-444
HHFNJ05	443	930899	AL121657	2118	1-125 139-1354
HHFNJ05	443	930899	AC015980	2119	1-318
HHFNJ05	443	930899	AL121657	2120	1-183 1297-1615
HHFMX34	444	945385	AC068499	2121	1-255 649-898 1564-1640 1742-1872 2784-3528
HHFMX34	444	945385	AC068499	2122	1-181
HHFMX34	444	945385	AC068499	2123	1-861 1666-1889 1923-2050 2743-2845 3525-3649 4171-4355 4660-4760 5305-6319 7098-7233
HHFLU06	445	857884	AL096870	2124	1-68 218-379 706-840 1000-1180

2020E0405T600T

					1505-2004 2014-2301 3897-3942 4074-4162 4353-4422 4764-4865 4941-5356 5850-5932 6040-6181 6664-6917 7152-7337 7431-7624 8016-8175 8346-8525 9445-9926 10349-10496 10802-10912 10949-11881
HHFLU06	445	857884	AL096870	2125	1-262
HHFLJ51	448	857898	Z80897	2126	1-83 445-925 1105-1728 3829-3949 4113-4310 4851-4962 5196-5510 5739-6151 6188-6735 6764-7120 7155-9664
HHFLJ51	448	857898	Z80897	2127	1-435
HHFLJ51	448	857898	Z80897	2128	1-428
HHFLI10	449	963162	AC025145	2129	1-631
HHFLI10	449	963162	AC025145	2130	1-102 1188-1294 1395-1671 2450-2912 2957-4002 5146-5499 5831-6235 6978-7038



					7222-7295 9054-10739
HHFLH62	451	857908	AC009260	2131	1-655
HHFLH62	451	857908	AC069240	2132	1-655 693-1132 1637-1983 2129-2540
HHFLH62	451	857908	AL080279	2133	1-655 693-1132 1637-1983 2129-2540
HHFLH62	451	857908	AC009260	2134	1-364
HHFLH62	451	857908	AC069240	2135	1-69 154-406 641-789 1249-1488 1646-2009
HHFLH62	451	857908	AL080279	2136	1-277
HHFLH62	451	857908	AC009260	2137	1-440
HHFLH62	451	857908	AC069240	2138	1-277
HHFLH62	451	857908	AL080279	2139	1-69 154-406 641-789 1249-1488 1646-2009
HHFLE12	452	969531	AC016769	2140	1-159 582-1086
HHFLE12	452	969531	AC016769	2141	1-31 286-1049
HHFIA58	456	858011	AC011782	2142	1-172 285-514 731-1209 2624-2993 3007-3112 4226-6800
HHFIA58	456	858011	AC011782	2143	1-454
HHFIA58	456	858011	AC011782	2144	1-486
HHFIA13	457	657405	AP001093	2145	1-875
HHFIA13	457	657405	AP001890	2146	1-419
HHFIA13	457	657405	AC011720	2147	1-875
HHFIA13	457	657405	AC009551	2148	1-875

202004051500T

HHFIA13	457	657405	AP001093	2149	1-303
HHFHY95	458	795053	AC005839	2150	1-1417
HHFHY95	458	795053	AC005839	2151	1-104
HHFHY95	458	795053	AC005839	2152	1-453
HHFHM22	462	674841	AC006487	2153	1-477
HHFHM22	462	674841	AC015656	2154	1-428
HHFHM22	462	674841	AC060233	2155	1-428
HHFHM22	462	674841	AC006487	2156	1-342
HHFHM22	462	674841	AC060233	2157	1-342
HHFHM22	462	674841	AC060233	2158	1-913
HHFHJ90	463	675218	AC025605	2159	1-357 443-919
HHFHJ90	463	675218	AC025605	2160	1-653
HHFHD38	464	709082	AL161651	2161	1-312
HHFHD38	464	709082	AL355542	2162	1-312
HHFHD38	464	709082	AL158037	2163	1-312
HHFHD38	464	709082	AL158037	2164	1-897 971-2743 2892-4334 4530-5979
HHFHD38	464	709082	AL161651	2165	1-897 971-2743 2891-4333 4529-5979
HHFHD38	464	709082	AL355542	2166	1-898 972-2744 2892-4334 4530-5980
HHFGX03	466	924753	AL359815	2167	1-962 1420-1815 2091-2328 3095-3210 3693-3824
HHFGX03	466	924753	AL359815	2168	1-391
HHFGR30	468	692887	AC025857	2169	1-425
HHFGR30	468	692887	AF285442	2170	1-129 1834-2165 3005-3269 3419-3802 4272-4470

					4516-6168 8490-8670 9026-9540 9759-10042 10490-10799 11249-11673 12087-12508 12947-13289 13383-14422 15702-16013 16205-17581
HHFGR30	468	692887	AC025857	2171	1-143 499-1013 1237-1515 1963-2272
HHFGR30	468	692887	AC025857	2172	1-422
HHFGR30	468	692887	AF285442	2173	1-240
HHFFR95	477	796677	AC027518	2174	1-429
HHFFR95	477	796677	AC027518	2175	1-592
HHFFP17	480	880667	AL357497	2176	1-451
HHFFP17	480	880667	AL024498	2177	1-451
HHFFP17	480	880667	AL357497	2178	1-266
HHFFP17	480	880667	AL357497	2179	1-113
HHFFP17	480	880667	AL024498	2180	1-266
HHFFP17	480	880667	AL024498	2181	1-113
HHFFO46	481	530501	AC073075	2182	1-324
HHFFO46	481	530501	AL391056	2183	1-324
HHFFO46	481	530501	AC073075	2184	1-652
HHFFO46	481	530501	AC073075	2185	1-686
HHFFO46	481	530501	AL391056	2186	1-686
HHFFO46	481	530501	AL391056	2187	1-652
HHFFK30	484	858051	AC074244	2188	1-549
HHFFK30	484	858051	AL365495	2189	1-549
HHFFF92	489	790572	AC008267	2190	1-1586 2346-3145
HHFFF92	489	790572	AC008267	2191	1-118
HHFFF92	489	790572	AC008267	2192	1-458
HHFDA13	496	667804	AC073237	2193	1-436
HHFDA13	496	667804	AC073347	2194	1-436
HHFDA13	496	667804	AC073237	2195	1-167

HHFDA13	496	667804	AC073347	2196	1-167
HHFDA13	496	667804	AC073237	2197	1-387
HHFDA13	496	667804	AC073347	2198	1-98
HHFCT63	497	572784	AC021037	2199	1-272
HHFCT63	497	572784	AC021037	2200	-1-42 67-197
HHFCT63	497	572784	AC021037	2201	1-368
HHFCP39	498	429442	AL365197	2202	1-1022
HHFCP39	498	429442	AL360295	2203	1-3499
HHFCP39	498	429442	AL035705	2204	1-3498
HHFCP39	498	429442	AL360295	2205	1-529
HHFCP39	498	429442	AL035705	2206	1-529
HHFCN59	500	739657	AL136171	2207	1-558
HHFCN59	500	739657	AL136171	2208	1-708
HHFCN13	501	667805	AC011324	2209	1-572 671-925 1087-1466 1816-2216 2498-2991 7984-8314 14041-14139 17879-18271 18532-18703 18858-19140 19597-20052 20677-20765 22029-22547 22751-23694 25142-25457 27691-28945 31825-32122 32489-32604
HHFCN13	501	667805	AC022080	2210	1-380 730-1130
HHFCN13	501	667805	AC068293	2211	1-381 731-1131
HHFCN13	501	667805	AC022080	2212	1-494
HHFCN13	501	667805	AC068293	2213	1-254
HHFCN13	501	667805	AC022080	2214	1-255
HHFCN13	501	667805	AC068293	2215	1-494

10091504-030702

HHFCM51	502	509631	AL139193	2216	1-192
HHFCH52	503	911570	AC009469	2217	1-345
HHFCH52	503	911570	AC009469	2218	1-138
HHFCE73	505	764763	AC011236	2219	1-1187
HHFCE73	505	764763	AC011236	2220	1-388
HHFCE73	505	764763	AC011236	2221	1-712
HHFCE40	506	712866	AL138706	2222	1-50 166-221 1953-2025 3258-3677 4973-4998 5552-5583 6350-6836 6918-7452 7594-9776 10184-10341 11085-11577 11979-12219 12391-12435
HHFCD43	507	714353	AC069165	2223	1-6088 6581-7107 7238-7612
HHFCD43	507	714353	AC069165	2224	1-79 176-448 1406-1701 2708-2957 3004-3130 3532-4127 4248-4536 5587-5675 8104-9589 9689-9809 10850-10949 11111-11687 13731-13968
HHFCD43	507	714353	AC069165	2225	1-296
HHFCC60	508	739669	AC024035	2226	1-472
HHFCC60	508	739669	AC013693	2227	1-472
HHFCC60	508	739669	AC024035	2228	1-502
HHFCC60	508	739669	AC013693	2229	1-338
HHFCC60	508	739669	AC013693	2230	1-502

20200405T6001

HHFBU63	512	745661	AL161784	2231	1-702
HHFBU63	512	745661	AC011969	2232	1-702
HHFBU63	512	745661	AC006558	2233	1-704
HHFBU63	512	745661	AL161784	2234	1-348
HHFBU63	512	745661	AC011969	2235	1-348
HHFBU63	512	745661	AC011969	2236	1-300 891-1330 1365-1850 2053-2512
HHFBU63	512	745661	AC006558	2237	1-302 895-1337 2067-2529
HHFBU07	513	954478	AC011271	2238	1-565
HHFBU07	513	954478	AC008741	2239	1-565
HHFBQ94	515	796838	AC073389	2240	1-616 1011-1677 2444-2728 2831-3372 3913-4145 5401-5519
HHFBQ94	515	796838	AC073389	2241	1-457
HHFBQ94	515	796838	AC073389	2242	1-327
HHFBP29	516	710894	AP002420	2243	1-1113 1314-1622 1654-1913 2085-2325 2828-3393 4312-4385
HHFBP29	516	710894	AC009052	2244	1-1113 1314-1622 1654-1913 2085-2325 2830-3396 4315-4391 5871-6041 8072-8343 8483-8561 11009-11161
HHFBP29	516	710894	AC004382	2245	1-1113 1320-1622 1654-1913

					2085-2325 2830-3396 4312-4391 5877-6047 8493-8571 11015-11171 11411-11604 11737-12079 12352-13598
HHFBP29	516	710894	AC004382	2246	1-555
HHFBM11	518	968002	AC008026	2247	1-517
HHFBM11	518	968002	AC010964	2248	1-1737
HHFBM11	518	968002	AC007996	2249	1-1796
HHFBM11	518	968002	AC007996	2250	1-869
HHFBM11	518	968002	AC007996	2251	1-59 98-1081
HHFBD83	519	781525	AC005343	2252	1-96 784-1143 1734-1849 2133-2401 3742-5557
HHFBA11	522	967991	AL356915	2253	1-375
HHFBA11	522	967991	AL356915	2254	1-410
HHFBA11	522	967991	AL356915	2255	1-159
HHBGN74	524	765214	AC010344	2256	1-627 1004-4346
HHBGN74	524	765214	AC010344	2257	1-377
HHBGJ53	525	909912	AC024740	2258	1-160 497-598 882-1177 2484-2930 3412-3474 4974-5279 5401-5511 7287-7408 7618-7741 8122-8503 9322-9392 9508-9613 9792-9880 10554-10663

2020E0740576007

					11328-11539 11871-12149
HHBGJ53	525	909912	AC026803	2259	1-522 979-1066 2563-2870 2993-3103 5037-5169 5211-5334 5715-6053 6915-6985 7101-7206 8147-8256 8922-9137 9467-9745
HHBGJ53	525	909912	AC024740	2260	1-37 713-1145
HHBGJ53	525	909912	AC026803	2261	1-291
HHBGG10	526	963849	AC011374	2262	1-477
HHBGG10	526	963849	AC011337	2263	1-477
HHBFT06	528	934826	AC009506	2264	1-1319
HHBFT06	528	934826	AC009506	2265	1-403
HHBFM77	529	771816	AC044809	2266	1-1731
HHBEG80	533	951688	AL355885	2267	1-75 188-497 613-693 766-3364
HHBEG80	533	951688	AL355885	2268	1-455
HHBEG72	534	761150	AL365197	2269	1-1022
HHBEG72	534	761150	AL360295	2270	1-3499
HHBEG72	534	761150	AL035705	2271	1-3498
HHBEG72	534	761150	AL360295	2272	1-529
HHBEG72	534	761150	AL035705	2273	1-529
HEMGX57	536	872083	AC026532	2274	1-1402
HEMGX57	536	872083	AL133375	2275	1-1402
HEMGX57	536	872083	AL355385	2276	1-1402
HEMGX57	536	872083	AC026532	2277	1-63 351-440 653-831 1091-1225 1550-1914 1987-2093



20200240516001

					2169-2663 3247-3384
HEMGX57	536	872083	AC026532	2278	1-593
HEMGX57	536	872083	AL133375	2279	1-232 1484-1579 1716-1889 2574-2832 3771-3878 4166-4255 4468-4646 4906-5040 5365-5729 5802-5908 5984-6478 7062-7199
HEMGX57	536	872083	AL355385	2280	1-593
HEMGX57	536	872083	AL133375	2281	1-586
HEMGX57	536	872083	AL355385	2282	1-96 233-406 1042-1347 2288-2395 2683-2772 2985-3163 3423-3557 3882-4246 4319-4425 4501-4995 5579-5716
HEMFN30	539	692818	AL138828	2283	1-1045
HEMFN30	539	692818	AC022654	2284	1-1045
HEMFN30	539	692818	AL138828	2285	1-304 767-1440 2390-2488 4487-4781 8192-8273 12771-12988
HEMFN30	539	692818	AL138828	2286	1-273
HEMFN30	539	692818	AC022654	2287	1-94 5044-5347 5810-6483 7433-7531

					9535-9829 13240-13324 17819-18036
HEMFN30	539	692818	AC022654	2288	1-273
HEMFF16	540	576539	AC027272	2289	1-653
HEMFF16	540	576539	AC016586	2290	1-397
HEMFF16	540	576539	AC027272	2291	1-169
HEMFF16	540	576539	AC016586	2292	1-81 1174-1224 1309-1436 1836-1904 2233-2616 3041-3257 4501-4940 5112-5176 7004-7423 7957-8002 8214-8644 9821-10063 10981-11092 11741-11869
HEMFF16	540	576539	AC027272	2293	1-112
HEMFF16	540	576539	AC016586	2294	1-169
HEMEA03	542	921922	AL162730	2295	1-419
HEMEA03	542	921922	AL162730	2296	1-202
HEMEA03	542	921922	AL162730	2297	1-355
HEMCV44	544	574321	AC067756	2298	1-424
HEMCV44	544	574321	AC021837	2299	1-424
HEMCV44	544	574321	AC011308	2300	1-424
HEMCV44	544	574321	AC067756	2301	1-475
HEMCV44	544	574321	AC021837	2302	1-475
HEMCV44	544	574321	AC067756	2303	1-156
HEMCV44	544	574321	AC011308	2304	1-156
HEMCV44	544	574321	AC021837	2305	1-156
HEMCV44	544	574321	AC011308	2306	1-475
HEMCI41	547	712614	AC068637	2307	1-993
HEMCI41	547	712614	AC008120	2308	1-993
HEMCI41	547	712614	AC068637	2309	1-1424
HEMCI41	547	712614	AC008120	2310	1-1424
HEMCC38	548	707453	AC007616	2311	1-283

20091504.030702

					1585-3443 3686-4902 5558-5704 6041-6150
HEMCC38	548	707453	AC007613	2312	-1-283 1585-3443 3686-4902 5558-5704 6036-6151 8423-8526 9199-10562 12224-12577 13496-13996 14166-17946
HEMAL61	551	851106	AL356124	2313	1-532 1491-2021 6384-6477 7549-7576
HELHJ74	553	765696	AC007279	2314	1-108 609-761 3108-3562 4639-4820 5667-5787 8416-8496 11327-11405 13714-14015 15679-16042 18174-18630 25423-25699 25789-25900 25962-26240 26738-27049 33446-33649 34051-34334 37491-37775 38208-38363 39524-39655 39680-40045 40177-40743 43779-43858 44056-44205

					44914-45142 49029-49117 50868-51004 51950-51979 52641-53034 53289-53447 53454-53986
HELHJ74	553	765696	AC007279	2315	1-1654
HELHJ74	553	765696	AC007279	2316	1-159 555-909 2593-2625 4471-5067 6631-6770 10086-10210 11310-11468 13666-13784 14092-14289 15370-15442 16104-16150
HELHC59	555	769404	AC073610	2317	1-2422 2490-2790 3039-3077 3371-3544 3585-3774 3860-3927 4112-4357
HELHC59	555	769404	AC073610	2318	1-696
HELGY42	556	713019	AC012202	2319	1-648
HELGY42	556	713019	AC025019	2320	1-648
HELGY42	556	713019	AC012202	2321	1-390
HELGY42	556	713019	AC025019	2322	1-390
HELGK56	561	925698	AC023789	2323	1-732
HELGG21	562	671071	AC023225	2324	1-472
HELGG21	562	671071	AC023225	2325	1-869 991-1462
HELGG21	562	671071	AL138846	2326	1-304 423-553 1434-2199 3430-3470 6425-6910 7169-7453

10091504.030702

					7726-8594 8716-9187 9274-9340 10220-10323 10333-10484 10770-10868 12169-12276 12626-12744 12959-13087 13628-13739 14496-14592 15842-17239
HELGG21	562	671071	AC023225	2327	1-63 949-1052 1062-1217 1500-1598 2892-2999 3350-3468 3685-3811 4351-4478 5217-5305 6563-7960
HELGG21	562	671071	AC023225	2328	1-426
HELGG21	562	671071	AL138846	2329	1-442
HELGG21	562	671071	AL138846	2330	1-638
HELGD47	563	851143	AP001320	2331	1-964
HELGD47	563	851143	AP000848	2332	1-459
HELGD47	563	851143	AP000929	2333	1-964
HELGD47	563	851143	AP001320	2334	1-454
HELGD47	563	851143	AP000848	2335	1-416
HELGD47	563	851143	AP000929	2336	1-453
HELFQ55	564	732224	AF267167	2337	1-869 883-1439 1680-2082
HELFQ55	564	732224	AF165145	2338	1-869 883-1439 1680-2082
HELFQ55	564	732224	AF267167	2339	1-592
HELFQ55	564	732224	AF267167	2340	1-288
HELFQ55	564	732224	AF165145	2341	1-592
HELFQ55	564	732224	AF165145	2342	1-288

202009.40516007

HELFJ35	566	506277	AC026558	2343	1-370
HELFJ35	566	506277	AC023973	2344	1-370
HELFJ35	566	506277	AC026558	2345	1-530
HELFJ35	566	506277	AC023973	2346	1-530
HELFJ35	566	506277	AC026558	2347	1-871 1399-1797
HELFJ35	566	506277	AC023973	2348	1-873 1403-1801
HELEZ81	568	571340	AL049712	2349	1-813 1243-2039 2304-3038 3047-3450 3473-3638 3800-4012 4383-8646 8760-9748 9758-10031 10041-10310 12504-12632 12709-13273
HELEZ81	568	571340	AL049712	2350	1-133
HELET68	569	800029	AC008267	2351	1-433
HELET68	569	800029	AC008267	2352	1-513
HELEO42	570	579016	AC010674	2353	1-110 3938-4022 5339-5881
HELEO42	570	579016	AC010674	2354	1-114 3201-3287 5782-6912
HELEO42	570	579016	AC010674	2355	1-1038 2072-2350 2444-2783 3079-3144 4658-5350 6459-6639 7394-7624 10010-10103 10404-10463 10557-10641 10928-11380 11817-11998

10091504-030702

					12307-12468 14217-14314 15654-15714
HELEH76	571	506674	AL360298	2356	1-944
HELEH76	571	506674	AL358072	2357	1-972
HELEH76	571	506674	AC053494	2358	1-972
HELEH76	571	506674	AC022671	2359	1-972
HELEH76	571	506674	AC053494	2360	1-470 766-1174
HELDL08	575	959919	AC073595	2361	1-453 819-1737
HELDL08	575	959919	AC027288	2362	1-454 820-1738
HELDL08	575	959919	AC027288	2363	1-584
HELDG91	578	790371	AC026562	2364	1-774
HELDG91	578	790371	AC069256	2365	1-774
HELDG91	578	790371	AC026562	2366	1-255
HELDG91	578	790371	AC069256	2367	1-255
HELDG91	578	790371	AC026562	2368	1-544
HELDG91	578	790371	AC069256	2369	1-544
HELCW51	579	531073	AC078913	2370	1-165
HELCI30	580	691024	AC016697	2371	1-510
HELCI30	580	691024	AC012362	2372	1-140 2160-2746 2944-3409 3461-3941
HELCI30	580	691024	AC016697	2373	1-126
HELCI30	580	691024	AC012362	2374	1-484
HELCI30	580	691024	AC012362	2375	1-108
HELCG36	581	655045	AL133377	2376	1-1464
HELCG36	581	655045	AL136164	2377	1-1563
HELCG36	581	655045	AL136164	2378	1-277
HELCG36	581	655045	AL133377	2379	1-277
HELCG36	581	655045	AL136164	2380	1-1285
HELBUI1	582	967661	AC010209	2381	1-697
HELBUI1	582	967661	AC010209	2382	1-396
HELBUI1	582	967661	AC010209	2383	1-284
HELBC83	583	781412	AL139328	2384	1-515 1506-1971
HELBC83	583	781412	AC018379	2385	1-515

20200804 40576007

					1506-1971
HELBC83	583	781412	AL136527	2386	1-515 1506-1971
HELBC83	583	781412	AL139328	2387	1-413
HELBC83	583	781412	AC018379	2388	1-631 637-1252
HELBC83	583	781412	AL139328	2389	1-631 637-1252
HELBC83	583	781412	AL136527	2390	1-631 637-1252
HELBC83	583	781412	AL136527	2391	1-413
HELAZ48	584	864515	AL162734	2392	1-2089
HELAZ48	584	864515	AL162734	2393	1-1074
HELAZ48	584	864515	AL162734	2394	1-310
HELAM32	587	699661	AC008079	2395	1-295 428-875 892-945 1563-1700 2084-2191 5109-5504 6802-7282 9658-9809 9818-10338 10350-11130 11158-15251 15262-15836 15909-16774
HELAM32	587	699661	AC008101	2396	1-295 428-875 892-945 1563-1700 2032-2191 5109-5504 6801-7281 9657-9808 9817-10337 10349-11129 11157-15257 15269-15843
HELAM32	587	699661	AC016830	2397	1-145 433-1745



HELAM32	587	699661	AC016027	2398	1-295 428-875 892-945 1563-1700 2084-2191 5109-5504 6028-6128 6802-7282 9658-9809 9818-10338 10350-11130 11158-15836 15909-16774
HELAM32	587	699661	AC008079	2399	1-287
HELAM32	587	699661	AC008079	2400	1-297
HELAM32	587	699661	AC008101	2401	1-287
HELAM32	587	699661	AC008101	2402	1-802
HELAM32	587	699661	AC016830	2403	1-687
HELAM32	587	699661	AC016830	2404	1-284 574-1700 2012-2132
HELAM32	587	699661	AC016027	2405	1-287
HELAM32	587	699661	AC016027	2406	1-304
HCMSY80	589	526182	AC069028	2407	1-255
HCMSQ63	590	745584	AC025975	2408	1-52 752-1355
HCMSQ63	590	745584	AC008681	2409	1-52 757-1360
HCMSQ63	590	745584	AC025975	2410	1-530 837-919 1945-3561
HCMSQ63	590	745584	AC008681	2411	1-530 837-919 1944-3560
HAHFE11	594	965293	AL008718	2412	1-466 554-874 2846-2933 4193-4484 4853-5225 5311-5727 6015-9575

HAHFE11	594	965293	AC062010	2413	1-373 459-875 1166-2221
HAHFE11	594	965293	AL008718	2414	1-440
HAHCU22	596	848831	AC021054	2415	1-62 909-1442 1984-4152
HAHCU22	596	848831	AC006205	2416	1-54 796-950 1203-1275 1695-1850 2485-2616 3454-3987 4529-6698
HAHCU22	596	848831	AC021054	2417	1-395 928-1113 1661-1885
HAHCU22	596	848831	AC006205	2418	1-130
HAHCU22	596	848831	AC006205	2419	1-395 928-1113 1661-1885
HAHAD95	600	865104	AC026149	2420	1-338
HAHAD95	600	865104	AC026149	2421	1-154
HAHAD95	600	865104	AC026149	2422	1-1125
HAFBG30	601	693363	AC010264	2423	1-577
HAFBG30	601	693363	AC005740	2424	1-2469 2938-3229 5273-5782 6611-6709 7387-7969 8404-8801 9433-9896 12391-13237
HAFBG30	601	693363	AC010264	2425	1-477 743-1093 1601-1973 2653-3336 3487-6292 6493-6735
HAFBG30	601	693363	AC010264	2426	1-257
HAFBG30	601	693363	AC005740	2427	1-477

EDZDEO"405T600T

					743-1093 1601-1973 2653-3336 3487-6735 6910-7038 7213-7317 7329-8051 9152-10208 10293-11443 11593-12545 12811-13152 13422-13481 13601-14334 14418-14524 14625-14713 14958-15119 15309-15446 16402-16558 19172-19292 19680-19767 20000-20121 20728-20869
HAFAY37	602	928705	AC073264	2428	1-338 804-3157
HAFAY37	602	928705	AC024155	2429	1-340 808-1235 1241-3169
HAFAY37	602	928705	AC073264	2430	1-488 1407-2258 2342-3262
HAFAY37	602	928705	AC073264	2431	1-534
HAFAY37	602	928705	AC024155	2432	1-535
HAFAY37	602	928705	AC024155	2433	1-488 1407-2258 2342-3262
HAECA04	604	932993	AC012419	2434	1-401 1140-1492 1588-2434
HAECA04	604	932993	AC068288	2435	1-401 1140-1492 1588-2433

HAECA04	604	932993	AC015971	2436	1-401 1140-1492 1588-2433
HAECA04	604	932993	AC012419	2437	1-803
HAECA04	604	932993	AC068288	2438	1-465
HAECA04	604	932993	AC068288	2439	1-781 1537-2119
HAECA04	604	932993	AC015971	2440	1-362 408-802 1558-2142
HAECA04	604	932993	AC015971	2441	1-465
HAEAM82	605	781539	AC009790	2442	1-1057 1129-1711

[0060] Table 1B summarizes additional polynucleotides encompassed by the invention (including cDNA clones related to the sequences (Clone ID NO:Z), contig sequences (contig identifier (Contig ID:) contig nucleotide sequence identifiers (SEQ ID NO:X)), and genomic sequences (SEQ ID NO:B). The first column provides a unique clone identifier, "Clone ID NO:Z", for a cDNA clone related to each contig sequence. The second column provides the sequence identifier, "SEQ ID NO:X", for each contig sequence. The third column provides a unique contig identifier, "Contig ID:" for each contig sequence. The fourth column, provides a BAC identifier "BAC ID NO:A" for the BAC clone referenced in the corresponding row of the table. The fifth column provides the nucleotide sequence identifier, "SEQ ID NO:B" for a fragment of the BAC clone identified in column four of the corresponding row of the table. The sixth column, "Exon From-To", provides the location (i.e., nucleotide position numbers) within the polynucleotide sequence of SEQ ID NO:B which delineate certain polynucleotides of the invention that are also exemplary members of polynucleotide sequences that encode polypeptides of the invention (e.g., polypeptides containing amino acid sequences encoded by the polynucleotide sequences delineated in column six, and fragments and variants thereof).

TABLE 2

Clone ID NO:Z	Contig ID:	SEQ ID NO:X	Analysis Method	PFam/NR Description	PFam/NR Accession Number	Score/ Percent Identity	NT From	NT To
HAHEE05	928673	20	blastx.2	(AC005531) similar to mouse homeodomain- interacting protein 1	gb AAD04728.1	100%	39	422
HAHHE12	969107	24	blastx.14	N-RAP [Mus musculus]	gi 2351568 gb AAC5 3323.1	76% 67% 54% 78% 35% 75% 59% 32% 37% 34% 40% 34% 27% 27% 85% 26% 31% 55% 29% 30% 38% 34% 38%	307 139 460 667 514 673 481 526 598 541 601 547 457 514 762 619 541 703 544 460 619 598 619	621 333 612 762 708 759 576 708 717 705 705 675 621 708 803 756 693 762 675 585 696 693 696

HELEA45	954371	49	blastx.14	exopolyphosphatase [Escherichia coli]	gi 147343 gb AAA24 415.1		47%	526	588
							29%	598	690
							29%	607	717
							66%	457	492
							47%	526	576
							75%	627	662
							40%	526	585
							33%	616	696
							36%	649	705
							23%	460	585
HELHF07	949067	78	HMMER 1.8	PFAM: Aminotransferases class- III pyridoxal-phosphate 4-aminobutyrate aminotransferase (EC 2.6.1.19) 1 1 aminotransferase).	PF00202	gi 1742132 dbj BAA1 4871.1	37%	637	708
							37%	637	708
							56%	526	573
							39%	723	791
							57%	532	573
							50%	762	797
							50%	762	797
							28%	711	794
							27%	592	702
							26%	526	594
HELHF07	949067	78	blastx.14	aminotransferase (EC 2.6.1.19) 1 1 aminotransferase).	PF00202	gi 1742132 dbj BAA1 4871.1	29%	720	800
							30%	415	483
							89%	86	169
							100%	66	98
HELHF07	949067	78	blastx.14	aminotransferase (EC 2.6.1.19) 1 1 aminotransferase).	PF00202	gi 1742132 dbj BAA1 4871.1	38.85	95	295
							85%	83	295
							92%	21	98
							45%	246	311
HELHF07	949067	78	blastx.14	aminotransferase (EC 2.6.1.19) 1 1 aminotransferase).	PF00202	gi 1742132 dbj BAA1 4871.1	100%	1	18

HEMEK19	574209	92	HMMER 2.1.1	[Escherichia coli] PFAM: TonB dependent receptor C-terminal region	PF00593	46.8	81	233
HEMEU54	947801	94	blastx.14	blastx.14 samaphorin G [Mus musculus]	gi 1418942 emb CAA 66398.1	96% 50% 42% 51% 54% 35% 33% 42% 30% 41% 44% 55% 62% 45% 62%	11 122 128 122 14 68 134 29 122 343 248 302 305 296 305	325 280 298 238 112 235 328 112 238 429 301 328 328 328 328
HHBEM70	756949	107	HMMER 1.8	PFAM: Core histones H2A, H2B, H3 and H4	PF00125	13.08	144	215
HHBHO63	906947	121	HMMER 1.8	PFAM: Phorbol esters / diacylglycerol binding domain	PF00130	2.21	188	217
HHFBX77	959805	136	blastx.14	(AB012308) B2HC [Anthocidaris crassispina]	gi 4033608 dbj BAA3 5136.1	83%	13	366
HHFCA64	720849	137	HMMER 1.8	PFAM: Zinc-binding metalloprotease domain	PF00099	3.2	287	267
HHFGN31	908508	177	HMMER 2.1.1	PFAM: KRAB box	PF01352	64.1	93	215
HHFHC02	920510	192	HMMER	PFAM: Eukaryotic protein	PF00069	13.11	118	183



			1.8 blastx.14	kinase domain (AB023658) Ca/calmodulin-dependent protein kinase 1	gi 4512334 dbj BAA7 5246.1	94%	70	183
HHFJN02	918358	214	blastx.14	retrovirus-related reverse transcriptase pseudogene - slow loris	pir B25313 GNLRL1	53% 54%	297 390	208 286
HHFON19	910891	257	HMMER 2.1.1	PFAM: Dual specificity phosphatase, catalytic domain	PF00782	154.8	316	732
HHFUC26	960331	267	blastx.14	(AF143321) unknown [Homo sapiens]	gi 4929222 gb AAD3 3910.1 AF143321_1	68%	298	825
HMEGH46	887791	287	HMMER 1.8	PFAM: Src homology domain 3	PF00018	3.21	343	375
HULAI37	708923	307	HMMER 1.8	PFAM: C2 domain	PF00168	12.81	10	78
HULFB76	767873	313	HMMER 2.1.1	PFAM: Core histones H2A, H2B, H3 and H4	PF00125	13.48	99	173
HUSIW10	963324	330	blastx.14	PFAM: HIT family	PF01230	24.6	67	147
HUSYA63	928021	335	blastx.14	(AF098499) No definition line found [Caenorhabditis elegans]	gi 3786408 gb AAC6 7396.1	48% 41%	234 149	320 241
HUSZH03	922852	341	blastx.14	(AF116865) hedgehog- interacting protein [Mus musculus]	gi 4868122 gb AAD3 1172.1 AF116865_1	88%	251	439
HUSYN11	943237	345	HMMER 1.8	C06A6.3 gene product [Caenorhabditis elegans]	gi 1086626 gb AAA8 2295.1	34% 61%	367 259	633 297
				PFAM: Core histones H2A, H2B, H3 and H4	PF00125	13.67	238	315

HUSIE95	967176	366	blastx.14	(AL137556) hypothetical protein [Homo sapiens]	emb CAB70810.1	67% 96% 96%	137 240 241	319 320 321
			blastx.14	GS2NA [Homo sapiens]	gi 805095 gb AAB81551.1	56% 53% 37% 31% 33%	229 496 121 388 484	5 413 11 332 431
HUSIE08	908574	368	blastx.14	(AB024005) KRAB-containing zinc-finger protein KRAZ2 [Mus musculus]	gi 4514561 dbj BAA75468.1	70%	38	229
HUSHL86	960355	369	blastx.14	(AF151805) CGI-47 protein [Homo sapiens]	gi 4929563 gb AAD34042.1 AF151805.1	96% 100%	1142 1413	882 1330
HUSFF03	924616	389	blastx.14	(AF033276) A kinase anchor protein [Mus musculus]	gi 2852701 gb AAC02208.1	83% 47%	266 541	535 591
HHFLU06	857884	445	HMMER 2.1.1	PFAM: Adenylate and Guanylate cyclase catalytic domain	PF00211	108.8	17	268
HHFKX28	971102	453	blastx.14	Similarity to Yeast LPG22P protein (TR:G1151240); 1 1 cDNA EST EMBL:CI0626 comes from this gene; cDNA EST EMBL:CI0848	gi 3881836 emb CAB01454.1	76% 65% 91%	858 495 617	619 409 546
HHFJM64	958384	455	blastx.2	(AF026504) SPA-1 like protein p1294 [Rattus norvegicus]	gb AAB81526.1	83% 43% 29%	3 323 799	287 664 1266

HHFCH52	911570	503	blastx.14	INSERTIN=TENSIN HOMOLOG.	sp G256713 G256713	28%	847	1344
HHBGJ53	909912	525	HMMER 2.1.1	PFAM: PH domain	PF00169	38.3	160	267
HHBGG10	963849	526	blastx.14	(AB011527) MEGF1 [Rattus norvegicus]	gi 3449286 dbj BAA3 2458.1	90% 75% 45% 41%	98 210 210 219	3 112 151 184
HHBEG80	951688	533	HMMER 1.8	PFAM: Core histones H2A, H2B, H3 and H4	PF00125	12.4	371	436
HEMGL56	767669	538	HMMER 2.1.1	PFAM: Filamin/ABP280 repeat.	PF00630	84.1	45	209
HEMDX96	935963	543	blastx.14	(AF111170) unknown [Homo sapiens]	gi 4314286 gb AAD1 5563.1	79%	491	255
HEMBT61	939957	550	HMMER 2.1.1 blastx.2	PFAM: Eukaryotic protein kinase domain (AD000092) hypothetical human serine-threonine protein kinase R31240_1 [Homo sapiens]	PF00069 gb AAB51171.1	76.6 71%	16 13	285 441
HELGY02	948302	557	blastx.2	Similar to sulfatase [Caenorhabditis elegans]	gb AAA83618.1	59%	383	523
HELGW31	610003	558	HMMER 2.1.1 blastx.2	PFAM: Cytochrome C assembly protein (AE000309) heme exporter protein C [Escherichia coli]	PF01578 gb AAC75259.1	216.5 100%	672 603	1286 1337
HELGW31	957568	622	HMMER 2.1.1	PFAM: Cytochrome C assembly protein	PF01578	200.9	990	421

HELGW31	964303	623	blastx.2	(AE000309) heme exporter protein C [Escherichia coli] yejV [Escherichia coli]	gb AAC75259.1	99% 100%	5 621	619 713
HEL GK56	925698	561	HMMER 2.1.1 blastx.14	PFAM: Pyruvate kinase pyruvate kinase type II [Escherichia coli]	gb AAAI6392.1  PF00224	93% 60%	39 1	449 75
HEL FN75	658681	565	blastx.2	Molybdenum transport system permease protein ModB. [Escherichia coli]	gi 147459 gb AAA24473.1  dbj BAA35428.1	98% 65%	147 21	731 107
HAHFS80	954432	593	blastx.14	(AJ010306) human smoothelin, large isoform [Homo sapiens]	gi 4128006 emb CAA09077.1	100%	288	635
HAECA04	932993	604	blastx.14	(AF097473) ORF1 [Mus musculus]	gi 4204569 gb AAD10771.1	73%	3	275
						37% 53% 35% 61% 57%	106 239 40 178 94	216 322 99 216 135

20200910091600

[0061] Table 2 further characterizes certain encoded polypeptides of the invention, by providing the results of comparisons to protein and protein family databases. The first column provides a unique clone identifier, "Clone ID NO:", corresponding to a cDNA clone disclosed in Table 1A. The second column provides the unique contig identifier, "Contig ID:" which allows correlation with the information in Table 1A. The third column provides the sequence identifier, "SEQ ID NO:X", for the contig polynucleotide sequences. The fourth column provides the analysis method by which the homology/identity disclosed in the row was determined. The fifth column provides a description of PFam/NR hits having significant matches identified by each analysis. Column six provides the accession number of the PFam/NR hit disclosed in the fifth column. Column seven, "Score/Percent Identity", provides a quality score or the percent identity, of the hit disclosed in column five. Comparisons were made between polypeptides encoded by polynucleotides of the invention and a non-redundant protein database (herein referred to as "NR"), or a database of protein families (herein referred to as "PFam"), as described below.

[0062] The NR database, which comprises the NBRF PIR database, the NCBI GenPept database, and the SIB SwissProt and TrEMBL databases, was made non-redundant using the computer program nrdb2 (Warren Gish, Washington University in Saint Louis). Each of the polynucleotides shown in Table 1A, column 3 (e.g., SEQ ID NO:X or the 'Query' sequence) was used to search against the NR database. The computer program BLASTX was used to compare a 6-frame translation of the Query sequence to the NR database (for information about the BLASTX algorithm please see Altshul et al., J. Mol. Biol. 215:403-410 (1990), and Gish et al., Nat. Genet. 3:266-272 (1993)). A description of the sequence that is most similar to the Query sequence (the highest scoring 'Subject') is shown in column five of Table 2 and the database accession number for that sequence is provided in column six. The highest scoring 'Subject' is reported in Table 2 if (a) the estimated probability that the match occurred by chance alone is less than  $1.0 \times 10^{-7}$ , and (b) the match was not to a known repetitive element. BLASTX returns alignments of short polypeptide segments of the Query and Subject sequences which share a high degree of similarity; these segments are known as High-Scoring Segment Pairs or HSPs. Table 2 reports the degree of

similarity between the Query and the Subject for each HSP as a percent identity in Column 7. The percent identity is determined by dividing the number of exact matches between the two aligned sequences in the HSP, dividing by the number of Query amino acids in the HSP and multiplying by 100. The polynucleotides of SEQ ID NO:X which encode the polypeptide sequence that generates an HSP are delineated by columns 8 and 9 of Table 2.

[0063] The PFam database, PFam version 5.2, (Sonnhammer et al., Nucl. Acids Res., 26:320-322, (1998)) consists of a series of multiple sequence alignments; one alignment for each protein family. Each multiple sequence alignment is converted into a probability model called a Hidden Markov Model, or HMM, that represents the position-specific variation among the sequences that make up the multiple sequence alignment (see, e.g., R. Durbin et al., *Biological sequence analysis: probabilistic models of proteins and nucleic acids*, Cambridge University Press, 1998 for the theory of HMMs). The program HMMER version 1.8 (Sean Eddy, Washington University in Saint Louis) was used to compare the predicted protein sequence for each Query sequence (SEQ ID NO:Y in Table 1A) to each of the HMMs derived from PFam version 5.2. A HMM derived from PFam version 5.2 was said to be a significant match to a polypeptide of the invention if the score returned by HMMER 1.8 was greater than 0.8 times the HMMER 1.8 score obtained with the most distantly related known member of that protein family. The description of the PFam family which shares a significant match with a polypeptide of the invention is listed in column 5 of Table 2, and the database accession number of the PFam hit is provided in column 6. Column 7 provides the score returned by HMMER version 1.8 for the alignment. Columns 8 and 9 delineate the polynucleotides of SEQ ID NO:X which encode the polypeptide sequence which shows a significant match to a PFam protein family.

[0064] As mentioned, columns 8 and 9 in Table 2, "NT From" and "NT To", delineate the polynucleotides of "SEQ ID NO:X" that encode a polypeptide having a significant match to the PFam/NR database as disclosed in the fifth column of Table 2. In one embodiment, the invention provides a protein comprising, or alternatively consisting of, a polypeptide encoded by the polynucleotides of SEQ ID NO:X delineated in columns 8 and 9 of Table 2. Also

provided are polynucleotides encoding such proteins, and the complementary strand thereto.

[0065] The nucleotide sequence SEQ ID NO:X and the translated SEQ ID NO:Y are sufficiently accurate and otherwise suitable for a variety of uses well known in the art and described further below. For instance, the nucleotide sequences of SEQ ID NO:X are useful for designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the cDNA contained in Clone ID NO:Z. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling immediate applications in chromosome mapping, linkage analysis, tissue identification and/or typing, and a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y may be used to generate antibodies which bind specifically to these polypeptides, or fragments thereof, and/or to the polypeptides encoded by the cDNA clones identified in, for example, Table 1A.

[0066] Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

[0067] Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X, and a predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing cDNA Clone ID NO:Z (deposited with the ATCC on October 5, 2000, and receiving ATCC designation numbers PTA 2574 and PTA 2575; deposited with the ATCC on January 5, 2001, having the depositor reference numbers TS-1, TS-2, AC-1, and AC-2; and/or as set forth, for example, in Table 1A, 6 and 7). The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with

known methods. Further, techniques known in the art can be used to verify the nucleotide sequences of SEQ ID NO:X.

- [0068] The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

#### *RACE Protocol For Recovery of Full-Length Genes*

- [0069] Partial cDNA clones can be made full-length by utilizing the rapid amplification of cDNA ends (RACE) procedure described in Frohman, M.A., et al., Proc. Nat'l. Acad. Sci. USA, 85:8998-9002 (1988). A cDNA clone missing either the 5' or 3' end can be reconstructed to include the absent base pairs extending to the translational start or stop codon, respectively. In some cases, cDNAs are missing the start codon of translation. The following briefly describes a modification of this original 5' RACE procedure. Poly A<sup>+</sup> or total RNA is reverse transcribed with Superscript II (Gibco/BRL) and an antisense or complementary primer specific to the cDNA sequence. The primer is removed from the reaction with a Microcon Concentrator (Amicon). The first-strand cDNA is then tailed with dATP and terminal deoxynucleotide transferase (Gibco/BRL). Thus, an anchor sequence is produced which is needed for PCR amplification. The second strand is synthesized from the dA-tail in PCR buffer, Taq DNA polymerase (Perkin-Elmer Cetus), an oligo-dT primer containing three adjacent restriction sites (XhoI, Sall and ClaI) at the 5' end and a primer containing just these restriction sites. This double-stranded cDNA is PCR amplified for 40 cycles with the same primers as well as a nested cDNA-specific antisense primer. The PCR products are size-separated on an ethidium bromide-agarose gel and the region of gel containing cDNA products the predicted size of missing protein-coding DNA is removed. cDNA is purified from the agarose with the Magic PCR Prep kit (Promega), restriction digested with XhoI or Sall, and ligated to a plasmid such as pBluescript SKII (Stratagene) at XhoI and EcoRV sites. This DNA is transformed into bacteria and the plasmid clones sequenced to identify the correct protein-coding inserts. Correct 5' ends are confirmed by



comparing this sequence with the putatively identified homologue and overlap with the partial cDNA clone. Similar methods known in the art and/or commercial kits are used to amplify and recover 3' ends.

[0070] Several quality-controlled kits are commercially available for purchase. Similar reagents and methods to those above are supplied in kit form from Gibco/BRL for both 5' and 3' RACE for recovery of full length genes. A second kit is available from Clontech which is a modification of a related technique, SLIC (single-stranded ligation to single-stranded cDNA), developed by Dumas et al., Nucleic Acids Res., 19:5227-32 (1991). The major differences in procedure are that the RNA is alkaline hydrolyzed after reverse transcription and RNA ligase is used to join a restriction site-containing anchor primer to the first-strand cDNA. This obviates the necessity for the dA-tailing reaction which results in a polyT stretch that is difficult to sequence past.

[0071] An alternative to generating 5' or 3' cDNA from RNA is to use cDNA library double-stranded DNA. An asymmetric PCR-amplified antisense cDNA strand is synthesized with an antisense cDNA-specific primer and a plasmid-anchored primer. These primers are removed and a symmetric PCR reaction is performed with a nested cDNA-specific antisense primer and the plasmid-anchored primer.

#### ***RNA Ligase Protocol For Generating The 5' or 3' End Sequences To Obtain Full Length Genes***

[0072] Once a gene of interest is identified, several methods are available for the identification of the 5' or 3' portions of the gene which may not be present in the original cDNA plasmid. These methods include, but are not limited to, filter probing, clone enrichment using specific probes and protocols similar and identical to 5' and 3' RACE. While the full length gene may be present in the library and can be identified by probing, a useful method for generating the 5' or 3' end is to use the existing sequence information from the original cDNA to generate the missing information. A method similar to 5' RACE is available for generating the missing 5' end of a desired full-length gene. (This method was published by Fromont-Racine et al., Nucleic Acids Res., 21(7):1683-1684 (1993)). Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcript. A primer set

containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest, is used to PCR amplify the 5' portion of the desired full length gene which may then be sequenced and used to generate the full length gene. This method starts with total RNA isolated from the desired source, poly A RNA may be used but is not a prerequisite for this procedure. The RNA preparation may then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA, which may interfere with the later RNA ligase step. The phosphatase, if used, is then inactivated and the RNA is treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase. This modified RNA preparation can then be used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction can then be used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the cardiovascular system antigen of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the relevant cardiovascular system antigen.

- [0073] The present invention also relates to vectors or plasmids, which include such DNA sequences, as well as the use of the DNA sequences. The material deposited with the ATCC (deposited with the ATCC on October 5, 2000, and receiving ATCC designation numbers PTA 2574 and PTA 2575; deposited with the ATCC on January 5, 2001, having the depositor reference numbers TS-1, TS-2, AC-1, and AC-2; and/or as set forth, for example, in Table 1A, 6 and 7) is a mixture of cDNA clones derived from a variety of human tissue and cloned in either a plasmid vector or a phage vector, as shown, for example, in Table 7. These deposits are referred to as "the deposits" herein. The tissues from which some of the clones were derived are listed in Table 7, and the vector in which the corresponding cDNA is contained is also indicated in Table 7. The deposited material includes cDNA clones corresponding to SEQ ID NO:X described, for example, in Table 1A (Clone ID NO:Z). A clone which is isolatable from the ATCC Deposits by use of a sequence listed as SEQ ID NO:X, may include the

entire coding region of a human gene or in other cases such clone may include a substantial portion of the coding region of a human gene. Furthermore, although the sequence listing may in some instances list only a portion of the DNA sequence in a clone included in the ATCC Deposits, it is well within the ability of one skilled in the art to sequence the DNA included in a clone contained in the ATCC Deposits by use of a sequence (or portion thereof) described in, for example Tables 1A or 2 by procedures hereinafter further described, and others apparent to those skilled in the art.

[0074] Also provided in Table 7 is the name of the vector which contains the cDNA clone. Each vector is routinely used in the art. The following additional information is provided for convenience.

[0075] Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128,256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., *Nucleic Acids Res.* 16:7583-7600 (1988); Altting-Mees, M. A. and Short, J. M., *Nucleic Acids Res.* 17:9494 (1989)) and pBK (Altting-Mees, M. A. et al., *Strategies* 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Phagemid pBS may be excised from the Lambda Zap and Uni-Zap XR vectors, and phagemid pBK may be excised from the Zap Express vector. Both phagemids may be transformed into *E. coli* strain XL-1 Blue, also available from Stratagene.

[0076] Vectors pSport1, pCMVSport 1.0, pCMVSport 2.0 and pCMVSport 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, also available from Life Technologies. See, for instance, Gruber, C. E., et al., *Focus* 15:59- (1993). Vector lafmid BA (Bento Soares, Columbia University, New York, NY) contains an ampicillin resistance gene and can be transformed into *E. coli* strain XL-1 Blue. Vector pCR<sup>®</sup>2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. See, for instance, Clark, J. M., *Nuc. Acids Res.* 16:9677-9686 (1988) and Mead, D. et al., *Bio/Technology* 9: (1991).

10094504-030702

[0077] The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, and/or the deposited clone (Clone ID NO:Z). The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

[0078] Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of cardiovascular system associated genes corresponding to SEQ ID NO:X or the complement thereof, polypeptides encoded by SEQ ID NO:X or the complement thereof, and/or the cDNA contained in Clone ID NO:Z, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

[0079] The polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[0080] The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below). It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[0081] The polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson,

Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the cardiovascular system polypeptides of the present invention in methods which are well known in the art.

[0082] The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or the cDNA sequence contained in Clone ID NO:Z. The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X or a complement thereof, a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or the polypeptide sequence encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X, a polypeptide encoded by the cDNA contained in Clone ID NO:Z and/or a polypeptide sequence encoded by a nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B are also encompassed by the invention. The present invention further encompasses a polynucleotide comprising, or alternatively consisting of, the complement of the nucleic acid sequence of SEQ ID NO:X, a nucleic acid sequence encoding a polypeptide encoded by the complement of the nucleic acid sequence of SEQ ID NO:X, and/or the cDNA contained in Clone ID NO:Z.

[0083] Moreover, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in Table 1B column 6, or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in Table 1B column 6, or any combination thereof. In further embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in Table 1B, column 6, and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1B, column 5). In additional

embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in Table 1B, column 6, and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in Table 1B, column 6, and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

[0084] Further, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1), or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1), or any combination thereof. In further embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1) and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1B, column 5). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1) and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table

1B, column 1) and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

[0085] Further, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1B which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2), or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in column 6 of Table 1B which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2), or any combination thereof. In further embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2) and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1B, column 5). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2) and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in column 6 of Table 1B which correspond to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2) and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (See Table 1B, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode

these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

[0086] Moreover, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in the same row of Table 1B column 6, or any combination thereof. Additional, representative examples of polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in the same row of Table 1B column 6, or any combination thereof. In preferred embodiments, the polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the complementary strand(s) of the sequences delineated in the same row of Table 1B column 6, wherein sequentially delineated sequences in the table (i.e. corresponding to those exons located closest to each other) are directly contiguous in a 5' to 3' orientation. In further embodiments, above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in the same row of Table 1B, column 6, and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1B, column 5). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in the same row of Table 1B, column 6, and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated in the same row of Table 1B, column 6, and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.



20250410575007

[0087] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1B, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1B, column 2) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[0088] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1), and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A or 1B) or fragments or variants thereof. In preferred embodiments, the delineated sequence(s) and polynucleotide sequence of SEQ ID NO:X correspond to the same Clone ID NO:Z. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[0089] In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more of the sequences delineated in the same row of column 6 of Table 1B, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A or 1B) or fragments or variants thereof. In preferred embodiments, the delineated sequence(s) and polynucleotide sequence of SEQ ID NO:X correspond to the same row of column 6 of Table 1B. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[0090] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the

invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[0091] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[0092] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the sequences delineated in column 6 of Table 1B are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[0093] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the sequences delineated in

column 6 of Table 1B are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides, are also encompassed by the invention.

**[0094]** In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of another sequence in column 6 are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

**[0095]** In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of another sequence in column 6 corresponding to the same Clone ID NO:Z (see Table 1B, column 1) are directly contiguous. Nucleic acids which hybridize to the complement of these 20 lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

10091504.030702

[0096] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, a polynucleotide sequence in which the 3' 10 polynucleotides of one sequence in column 6 corresponding to the same contig sequence identifier SEQ ID NO:X (see Table 1B, column 2) are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[0097] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of another sequence in column 6 corresponding to the same row are directly contiguous. In preferred embodiments, the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B is directly contiguous with the 5' 10 polynucleotides of the next sequential exon delineated in Table 1B, column 6. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[0098] Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. Accordingly, for each contig sequence (SEQ ID NO:X) listed in the third column

200504050001

of Table 1A, preferably excluded are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 and the final nucleotide minus 15 of SEQ ID NO:X, b is an integer of 15 to the final nucleotide of SEQ ID NO:X, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:X, and where b is greater than or equal to a + 14. More specifically, preferably excluded are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a and b are integers as defined in columns 4 and 5, respectively, of Table 3. In specific embodiments, the polynucleotides of the invention do not consist of at least one, two, three, four, five, ten, or more of the specific polynucleotide sequences referenced by the Genbank Accession No. as disclosed in column 6 of Table 3 (including for example, published sequence in connection with a particular BAC clone). In further embodiments, preferably excluded from the invention are the specific polynucleotide sequence(s) contained in the clones corresponding to at least one, two, three, four, five, ten, or more of the available material having the accession numbers identified in the sixth column of this Table (including for example, the actual sequence contained in an identified BAC clone). In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example. All references available through these accessions are hereby incorporated by reference in their entirety.

TABLE 3

Clone ID NO: Z	SEQ ID NO: X	Contig ID:	EST Disclaimer		Accession #'s
			Range of a	Range of b	
HAHCL07	11	952661	1 - 1252	15 - 1266	AI472849, AA452197, AA216015, F16124, AA249665, AA211338, R15601, AA249623, AA404417, T82709, AA039429, AI800785, AI572463, and AA780204.
HAHCP26	12	681258	1 - 389	15 - 403	AC001234, and AF060568.
HAHCP41	13	712086	1 - 226	15 - 240	
HAHCP49	14	722659	1 - 242	15 - 256	
HAHCP55	15	865095	1 - 290	15 - 304	
HAHCP67	16	756916	1 - 250	15 - 264	AF176915. AW246326, T66078, T08451, AW068827, T35903, T66082, F12335, AA100537, W76237, R19223, F08087, Z43540, AA298126, F12331, W06986, AA298141, W07263, R47805, AC006252, Y17169, and AL136773.
HAHCP91	17	789998	1 - 274	15 - 288	AC005833.
HAHCR57	18	865096	1 - 474	15 - 488	T60981.
HAHEE04	19	922257	1 - 332	15 - 346	AA453887, AI939557, T95411, AI369869, AW295606, AC005531, AF170301, AF170302, AF077659, and AF144573.
HAHEE05	20	928673	1 - 408	15 - 422	
HAHEO46	21	718772	1 - 212	15 - 226	AA193162, Z24810, AA196396, AA194446, D58283, D80043, D80022, D80195, D59859, C14331, D80166, D80212, D80193, D59927, D51423, D59619, D80210, D51799, D80391, D80164, D59275, D80240, D80253, D59787, D80227, D81030, D59502, D80188, D80196, D80219, D57483, D80269, D80038, D80366, D50979, D59889, C14429, D50995, D59467, D59610, AW178893, D80378, D80024, C15076, D80045, AA305409, T03269, D80241, C14389, C75259, D51060, C14014, AA305578, AW352158, AW177440, D51022, AW179328, D80522, D80248, AW178775, D81026, AW378532, D51250, D80134, AA514188, AW360811, D80251, F13647, AW360834, AA514186, D80133, AW177501, AW177511, AW375405, D80168, C05695, AW178762, AW377671, Z19200, D58253, AI910186, D80268, AW176467, AW360817, AW366296, D80247, C14227, AW360844, D80132, AW375406, AW378534, D52291, AW179332, AW377672, AW179023, AW178905, D80439, D81111, AI905856, AW352170, AW352117, AW378540, D80302, C14407, AW352171, AW177505, AW377676, AW178906, AW177731, AW178907, AW378528, AW179019, AW179024, C14298,
HAHES10	22	961594	1 - 317	15 - 331	
HAHFX20	23	925753	1 - 125	15 - 139	
HAHHE12	24	969107	1 - 792	15 - 806	

				D80064, AW360841, AW179020, D51759, AW178909, AW177456, D59373, AW179329, Z21582, AW178980, AW178914, D80157, AW178774, AW177733, AW178908, AW178754, AW179018, D51103, T11417, AW179220, AA285331, D51097, AW179004, AW179012, AW177728, AW378543, AW378525, AW352120, AW178983, A1557751, T48593, AW367967, AW177722, AW352174, AW179009, AW178781, AW178911, AW352163, C06015, D45260, D59627, C14077, D59653, A1535850, D58246, AW367950, D80258, D80014, H67866, AW177723, D59503, C03092, D58101, H67854, AW178986, AA809122, A1557774, A1525923, AW378533, D60010, C14344, AW177508, C14046, T03116, A1525917, D59317, AW177497, C14973, D51213, D59474, D45273, AW177734, D51221, AA514184, A1525920, T02974, A1535686, D60214, C14957, D59551, A1525227, D51231, C16955, AW378542, A1525235, AW378539, A1525912, T03048, A1525242, Z33452, A1525925, AW360855, A1525215, H67858, C05763, A1525222, T02868, U76618, A62300, AJ132110, A84916, A62298, AR018138, Y17188, D26022, A25909, D34614, AF058696, X67155, A67220, D89785, A78862, AR008278, D88547, AB028859, X82626, Y12724, AR025207, I82448, A82595, AR060385, AB002449, A94995, AR008443, AR060138, I50126, I50132, I50128, I50133, AB012117, AR060488, AR016514, X68127, AR016808, A45456, A26615, AR052274, A85396, AR066482, A44171, X93549, Y09669, A43192, A43190, AR038669, A85477, I19525, A86792, AR066487, A30438, AR054175, D50010, Y17187, I14842, A70867, AR066490, A63261, I18367, AR008408, AR062872, D88507, AR008277, AR008281, AR016691, AR016690, U46128, D13509, A64136, A68321, AR060133, AF135125, I79511, U79457, AF123263, AR032065, AR008382, and A47134.	
HAHHO04	25	925814	1 - 299	15 - 313	AL133572.
HAHHS01	26	913863	1 - 167	15 - 181	
HAHIK10	27	961425	1 - 158	15 - 172	
HAHIW08	28	955803	1 - 364	15 - 378	AW291127.
HAHSC42	29	695111	1 - 449	15 - 463	Z18875, and AA311441.
HCMBA95	30	508774	1 - 332	15 - 346	
HCMSC52	31	522615	1 - 110	15 - 124	
HCMSE09	32	530511	1 - 356	15 - 370	AC006368.
HCMSEQ77	33	862373	1 - 297	15 - 311	AC006463.
HCMSS72	34	526183	1 - 161	15 - 175	
HCMSSU37	35	706337	1 - 482	15 - 496	
HCMSSX59	36	522598	1 - 362	15 - 376	
HELAF85	37	507230	1 - 309	15 - 323	A1808918, AW339979, F29562, Z24811, F24581, and AC006333.
HELAI26	38	920922	1 - 203	15 - 217	
HELAL34	39	527994	1 - 196	15 - 210	
HELAV61	40	507205	1 - 327	15 - 341	
HELB08	41	960106	1 - 221	15 - 235	

HELBD70	42	527677	1 - 223	15 - 237	
HELBK27	43	527669	1 - 280	15 - 294	
HELBN45	44	527666	1 - 244	15 - 258	AL031295.
HELBP62	45	527533	1 - 321	15 - 335	Z64392, and Z64391.
HELDG77	46	750478	1 - 209	15 - 223	
HELDK12	47	766209	1 - 219	15 - 233	
HELDOS6	48	531576	1 - 250	15 - 264	
HELEA45	49	954371	1 - 157	15 - 171	
HELEE09	50	531415	1 - 298	15 - 312	
HELEF52	51	506677	1 - 300	15 - 314	
HELEQ47	52	577187	1 - 402	15 - 416	
HELER30	53	574038	1 - 277	15 - 291	AC006057.
HELEU12	54	574023	1 - 703	15 - 717	AA004637, T79961, AC005324, and AL080124.
HELEU37	55	522407	1 - 182	15 - 196	AA368196, AC005247, and AC007971.
HELEU73	56	574016	1 - 391	15 - 405	
HELEU91	57	851160	1 - 323	15 - 337	
HELEW62	58	574025	1 - 292	15 - 306	
HELFF40	59	574058	1 - 188	15 - 202	AC000119.
HELFFH33	60	576530	1 - 240	15 - 254	E02347.
HELFFJ03	61	921943	1 - 419	15 - 433	
HELFFQ54	62	728888	1 - 489	15 - 503	AF179680, and AL022162.
HELFFQ79	63	577248	1 - 359	15 - 373	
HELGA54	64	576374	1 - 426	15 - 440	
HELGC24	65	576377	1 - 369	15 - 383	
HELGC32	66	699375	1 - 317	15 - 331	W26021, and W27138.
HELGC77	67	577317	1 - 230	15 - 244	
HELGG77	68	825795	1 - 917	15 - 931	AI865213, AA857296, AI433247, AA501614, AI859834, AL037910, AW337454, AL045077, AI049996, AI471815, AA577748, AI769271, AI445815, AL047858, AI623899, AI215778, AI890348, R92640, F17700, H29951, F32171, H23467, AI343143, N39097, AA342681, R83097, AI246080, AI151261, AA663966, AI053786, AI754013, AA557486, AA484143, AA297968, AI569982, AL046782, AA652813, AW103415, AI755236, AI754923, AA219098, AA457685, AI287964, AI753951, AI282661, H05073, AI538540, AW079809, AI431240, AI469796, AA491955, AI979005, AW407578, AI222678, AI687750, H57826, AI216054, AI270476, AI696901, AW088049, AW192330, AI382614, AW152303, AI733856, AI754291, AA021552, AI865988, AI609972, AI623563, AA716755, AI110405, AW272758, U85195, AE000658, AC004067, AC006241, AL121657, AC005832, AC007845, AC006318, AC004590, AC007298, AL049759, AC004253, AC002347, AC004019, AC012599, AC002544, AL049766.



HELGH89	69	545009	1 - 635	15 - 649	AL049694, AC005532, AC003661, AC005046, AC016025, AC005488, AC016026, AC005859, AL008721, AC000052, Z85986, AL121653, AC003664, AL049845, AL031296, AL034549, AC006539, Z93017, AC007191, AL050338, AC005839, AC003695, AC012330, AC007934, AC002454, AF053356, AC009510, AC002394, AL008635, AF196969, AC004477, AL109963, Z82194, AC004921, U63834, AC003029, AC004099, AL031577, AC008012, AC002316, AL022165, AC007325, AL022313, AC005694, AJ003147, AP000509, AC004047, AC005740, AC004791, AC002039, AC002996, AC004659, AC008132, AL034421, AC006048, AB007955, AC006160, AC006064, AC006372, AC004448, AF001549, AC005498, AL008582, AL133163, AL034423, AL109758, AC006204, AL133445, AC005914, D84394, AL035400, AC005722, AC007227, AC005745, AC002306, AC002301, AC005632, AC004024, AC002980, Z97055, AC002350, AC006552, AF121781, Z98949, AC007738, AC002470, AC005755, AL008715, Z82172, AC005887, AC005881, AL022323, AC006312, AC007637, AC004068, AC005529, AC006130, AC006285, AC004990, AL049829, AC004386, AP000553, Z83846, AL049776, AC001228, AF088219, AL022238, AC004682, Z84484, AC007486, AC004996, AL049869, AL031120, U51281, AC005911, AC005922, AC002468, U95740, AL109802, AC004797, AC005837, AL031289, AC008119, AC005531, AL031767, AP000066, Z93020, AL023803, Y18000, AL109627, AL033518, AC005288, AC009516, AC007221, AL049758, AC004531, AC005829, AC006390, AC002418, AC004472, AC005399, AC007055, AL096701, AC003013, AL078477, AC004963, AP000216, AL109865, AC004675, AL035427, AC002365, AL020993, AC007750, AL035089, AL033504, AC011718, AC006924, AC004830, AC004801, AC006449, AL049795, AB023049, AB023052, Z84469, AC004813, AF095901, AP000356, AL121694, AC004955, AC003982, AL021877, AC005221, AC005971, AL133448, AL050308, AC005065, Z95703, AP000339, AB023051, AC003051, AC005682, AL133246, AC002465, AC003101, AC009498, AL021937, AL021368, AL135960, AJ131016, AL009181, AC005412, AC007388, AL020997, AL031276, AL035045, AF001550, AC005697, AP000513, AC008498, AP000512, AL109654, AC005017, AC007347, AL021579, AL136295, AC005520, AC006468, Z98752, AC010197, AC008929, AC004140, AL035415, AL021997, AC005484, AC006153, Z97054, AC006316, AL096776, Z99716, AC004910, AC005004, AC007676, AL033527, AC011331, AF134726, AC009784, AC002558, AC006530, and AC006453.
HELGN53	70	963160	1 - 356	15 - 370	
HELGP50	71	576339	1 - 351	15 - 365	
HELQ48	72	851178	1 - 332	15 - 346	Z86061.
HELGT48	73	879483	1 - 635	15 - 649	
HELGT48	74	721742	1 - 394	15 - 408	
HELHB12	75	970863	1 - 392	15 - 406	AI307709, and AL020997.
HELHC49	76	576292	1 - 368	15 - 382	
HELHD46	77	719129	1 - 266	15 - 280	AB020722.
HELHF07	78	949067	1 - 327	15 - 341	

HELM29	79	883505	1 - 299	15 - 313	
HEMBC56	80	577797	1 - 404	15 - 418	
HEMBI16	81	507220	1 - 207	15 - 221	
HEMBZ84	82	527989	1 - 216	15 - 230	
HEMCA89	83	527985	1 - 243	15 - 257	
HEMCM25	84	948738	1 - 483	15 - 497	
HEMDG56	85	715834	1 - 211	15 - 225	AL022311.
HEMDG83	86	576508	1 - 598	15 - 612	AI971563, AW236461, R66941, and AI867853.
HEMDK92	87	574278	1 - 291	15 - 305	H62575, and N31620.
HEMDM56	88	577251	1 - 280	15 - 294	
HEMDO24	89	577297	1 - 378	15 - 392	
HEMEA72	90	527804	1 - 297	15 - 311	
HEMEH76	91	574345	1 - 404	15 - 418	AA329192, AW074398, AI830390, AA319218, AI061334, R91994, N71557, HI6048, AA659083, N71724, AA558697, AA457639, AI216799, AA405288, HI3868, AI679379, AI679888, AI867386, AA523490, AW008062, AI537185, AA378580, AI887768, AW271917, AF150222, AA291631, W38648, AI246409, AA399172, AA366035, AA394147, AL037554, F31204, AA515128, AA077935, AA584749, AA557879, AL046409, AI350211, AI581068, AA077952, AW261871, AA077817, AI688846, AI613280, AW405016, AI474085, AA385100, AI963720, AA776236, AA332991, AI879000, AW089789, AI608674, AI301218, AI205181, AA378886, AA310158, AA661573, AA569187, H54443, AA079421, AA532877, AA580662, AA503258, AW303876, H55779, F37286, AW419262, AA743956, AW193432, AI431303, AI286264, AI653886, AW029574, AI284640, AI471481, AI872216, AA516207, AL119259, AA100884, AW274349, AI244758, AW193265, AW303196, AW301350, AA569202, AA225155, AI282907, F36273, AA984191, AA745410, AA494087, AI806850, AW276827, AA179136, AI341664, AA362440, AW273469, AI678392, AW407007, AA353408, AI287528, AI820539, R63003, AI305766, AA112864, AW157173, AW162314, AL040913, AI110770, AW162332, AI284467, AI890348, AA508882, AI015912, AI821387, AA508103, AI721122, AW338869, AW402864, AA713815, AW419118, AI669443, W04195, AW264973, AA377767, AL041412, AI610159, AA347040, AA501617, AA344959, AA327323, AI914706, AA297776, AI624024, AC008064, Z83849, Z77894, AL049766, AF196971, AC007750, AC005815, AC000134, AC005200, Z82205, Z69666, X52851, AC007065, AC006511, AC003037, AL008629, AC006042, AC002564, AL031588, AC007773, L13713, L13714, AC007206, AL031965, AL035587, AL022316, AC005040, AC007458, AF015152, AC000100, AL023807, Z82244, AC005414, Z93241, Z93020, AL008723, AC002451, AC006213, AC002094, AC006031, AC008372, AL022397, AC002347, AC003080, AC006043, AC000048, AC005600, AL021579, Z82210, AC006057, AC005399, AC005102, AF001298, AC004643, Z98051, AC004771, AC003982, AL132712, U80017, I51997, AC004193, AC005002, Z99289, AF002223, AL080239, AF055066, U67211, AC007238, AL035668, AP000328, Z83001, AC006128, AC004088, AP000055, AP000170, AP000123, AL080276, AL023803, AC010202, AC005215, AL033521, Z98257,

					AL033403, AC002542, AF111168, Z49816, X55925, X60653, AC002559, AL023799, AL031729, AC018769, AC002352, AL031584, AL080248, AL121658, AL121748, AL135744, AC005257, L13709, AL117256, AC004501, AL033392, AL031777, AC005161, AC004055, AC008071, AC004552, AC004695, AC004057, AC006989, AC006479, AC006050, AP000961, AC007656, AC004664, AC006196, AL022476, X69951, AC000355, U69730, AL049761, AC010175, U95742, AL031904, AC007106, AC007216, AC005084, AC005495, Z99716, AL022164, U57009, AP000294, U63630, AC005303, AC000118, AP000111, AP000043, AF200465, AC004167, AC002073, AC004816, AC005245, AC016027, AC005297, AC016830, AC008045, X88791, AC006044, AF078925, AF165147, AF067122, Z73965, AF041427, Z83819, AP000252, AC005250, AP000212, AC005216, AC005660, AL031281, AL031289, Z83846, AL009181, AP000031, AC007676, AL023280, AC007281, AC005939, AC006040, AC006430, AC002365, AL020995, AF161351, AF131217, AF010238, Z68869, AC005253, AL008719, Z84485, Z98752, AB020859, AL034451, AL109807, AL049643, Z99943, AC004675, AC004520, AC006571, AL096829, AL121653, AP000432, AP000556, AL008627, AL024507, AL021407, AC006442, AP000552, AC005345, AL009029, AC007050, AC007151, AP000459, AC005324, AL021395, AC007437, AF088219, AC007738, and AL121823.
HEMEK19	92	574209	1 - 297	15 - 311	
HEMEN63	93	578717	1 - 307	15 - 321	
HEMEU54	94	947801	1 - 438	15 - 452	X97818.
HEMFL58	95	576505	1 - 338	15 - 352	
HEMFN33	96	702564	1 - 322	15 - 336	
HEMFXX20	97	840164	1 - 186	15 - 200	
HEMGL57	98	971118	1 - 302	15 - 316	AL049569.
HEMGL58	99	578091	1 - 440	15 - 454	R73040, AA005251, AA005165, and AF176815.
HHBBA47	100	720472	1 - 400	15 - 414	AW392670, U46347, AW384394, AW363220, AL043003, AL119484, AL119497, AL119439, AL119443, AW372827, AL119457, AL119319, AL119396, Z99396, AL119324, U46341, AL134528, AL119363, AL119341, AL119391, AL119355, U46350, U46351, U46349, AL119444, AL119483, U46346, AL119335, AL134533, AL119399, AL119522, AL119496, U46345, AL134531, AL042614, AL134132, AL134527, AL134538, AL119418, AL043147, AL042450, AL042965, AL042975, AL042542, AL042544, AL042970, AL043019, AL042984, AL043029, AL142134, AL119511, AL042551, AL119488, AL119464, AB026436, AR054110, A81671, AR060234, AR066494, and AR069079.
HHBB111	101	959756	1 - 401	15 - 415	AI920923, AW016123, AI694192, AW085200, AA789137, AA563721, AI079897, R51594, AA831766, AA782525, AW169957, AW339110, AI190317, AI910943, AA917419, AA512984, AA775828, AI393486, AA913674, AI374931, AA716669, AI813594, N93954, AI272716, AI189330, AI827728, AA481275, AI568483, AI340006, AA857984, AI241121, AI129733, AI969188, AI862837, AI803017, AA284394, AW068473, AA701544, AI096997, AW103900, AA570016, AA028141, AI249182, AA991848, AI470495, AI354796, AI743676, T40513, AL041504, N78775, AA865611, AA101174,

					AI419551, AI413347, AA766811, AA806671, HS2242, AL044039, R41974, AA027302, AI696783, AA058759, AA688050, AL041503, Z40289, AI915796, AA747265, D20480, AA937783, AA089759, AW237240, AA287007, and AB007930.
HHBKB65	102	588062	1 - 411	15 - 425	R36520.
HHBBL40	103	588066	1 - 541	15 - 555	AA526287, AI417476, AI473759, and AA604308.
HHBBL53	104	588067	1 - 406	15 - 420	
HHBEA32	105	927399	1 - 286	15 - 300	
HHBEE70	106	697541	1 - 492	15 - 506	
HHBEM70	107	756949	1 - 229	15 - 243	AL119457, AL042544, AL119399, AL042382, AL079794, AL042440, AL119511, AL037081, AL119324, AL043152, AI110828, AW235489, AW082113, AI280670, AI064830, AI610362, AI352497, AW149925, AA809129, AL079741, AI886753, AI269862, AI364788, AI134999, AW151136, AL121270, AL047187, AI500061, AL043168, AI345111, AI491897, AW170674, AI624543, AL119863, AW117746, AI349598, AL041150, AW269097, AI933589, AI445992, AI922365, AL045500, AI824576, AI309443, AW087445, AI345416, AI345612, AW163834, AI345415, AL046931, AL121328, AW268302, AW026882, AI610307, AI567935, AW163464, AI620284, AI281772, AI445990, AW15307, AW051258, AI921248, AI611738, AI571909, AI619502, AI680162, AI632408, AI3066613, AI677796, AI802542, AI433976, AI620089, AA449768, AI288305, AW118518, AI570807, AI635067, AI923370, AI627988, AL110402, AL047387, AI679174, AW168650, AI537677, AI670009, AI8666770, AL047675, AI433157, AI702073, AI909696, AI554821, AW117882, AI284131, AI927755, AI174394, AW172745, N33175, AI539771, AL079960, AI582932, AI500659, AI633125, AI805769, AI698391, AI815232, AI801325, AI500523, AI915291, AL042628, AW071177, AI284517, AI923989, AI500706, AI491776, AI445237, AW151138, AA580663, AI521560, AI889189, AI500662, AI284509, N80094, AI889168, AI866573, AI633493, AL048323, AI434256, AW302973, AI289937, AI537244, AW268060, AW072719, AI888661, AI284513, AI888118, AI344785, AI889147, AL037454, AI440252, AW302965, AW072484, AI702019, AI634251, AI569309, AI433037, AI280732, AW071362, AW301300, AI348917, AI343037, AI499285, AW088134, AL120300, AL048340, F27788, AI446373, AI864836, AL043981, AI569328, AI307543, AI874166, AI307210, AI669526, AI340659, AI288285, AI801460, AI345005, AI311892, AI307736, AI783504, AL039276, AI349266, AI273048, AL119791, AL040241, AL045163, AI281762, AI494201, AI686554, AL042745, AI949960, AL079977, AI886123, AI524671, AL040243, AW050522, AI917252, AW083804, AI345471, AW149221, AL041862, AI673297, AI274508, AL039783, AL046926, AL045266, W74529, AI439745, AI922901, AI963846, AL042627, AL049085, AW198075, AL048334, AI872711, AI587606, AI468872, AI932794, AW081255, AI886181, AI690426, AI862142, AI873644, AI802826, AL445432, AL037582, AL037602, AL041772, AL036214, AL079963, AI345608, AW169658, AI559737, AW410972, AL037030, AI310575, AL039132, AI889376, AL047763, AI335363, AI538716, AW074993, AI591420, AI362580, AI499986, AI560023, R32821, AI432218, AW051088, AL038504, AI308032, AI537515, AI934011, AL045620, AI308035, AI349276, AI281782, AI564719, AI866510, AI909642, Y11587, U77594, AC004987, I89947, AL122049, I48978,

					AL133640, AC004686, AL080159, A08913, AL133072, AF090934, Y14314, AL133077, A08916, A08910, A08909, S68736, AL050138, AL137271, Z82022, AL050277, I48979, X82434, I89931, AL110280, I49625, A77033, A77035, AF087943, AF183393, AL122110, AL133080, AF091084, I09360, AL080060, U35846, AF026816, I33392, AL050149, AL133016, U80742, AL122121, E07361, AL117460, AL110221, AF090903, AL117435, A1012755, AL049283, AF118070, AL137550, AF017152, AF146568, AL137463, AF113019, X84990, AF090900, AL133560, AF111112, E03348, AF113689, AR059958, AL133075, AL133565, AL049466, A65341, A1000937, AR000496, U39656, AL117583, A08912, AL117585, AL133113, X93495, AL122123, X72889, AC004594, AL049300, AL049452, AL110197, A93350, AL117457, AR038854, AL096744, U91329, AF003737, AL049938, AL137557, AF106862, AF113677, U67958, E02349, AF079763, AL137538, E07108, AF113676, AF158248, AF185576, A03736, I26207, AF118094, AF067728, AL050024, AL049430, AF113699, AL133557, AF125948, A45787, AF177401, S78214, AF090901, AL050393, AR011880, AL080124, AF113690, AF118064, AL133093, AL122050, AL137560, AL137459, AJ238278, AL122098, U00763, AL080127, L31396, AL133014, AL080137, AL137527, AF079765, L31397, X63574, U72620, A58524, A58523, A93016, AL049382, I42402, AL050172, AF162270, AL050116, A12297, AF057300, AF057299, AF061943, AF119337, AL137556, I03321, AF113691, AF104032, AF017437, E15569, AF113013, AL049464, AF078844, Y16645, AF097996, Y11254, AL110196, AF111851, AF026124, AL050108, AF090896, AL122093, X65873, U42766, AL137521, X96540, AB019565, AF113694, AF090943, X70685, AL049314, AL137648, AJ242859, AF125949, AL050146, AL110225, AL117394, AL133606, AL137476, E08263, E08264, L30117, AL137533, AL137480, AF061795, AF151685, AL117440, AL133104, AL133067, L19437, S61953, AL137526, AF132676, AF061836, AF111849, AL137523, U96683, A07647, AR038969, X87582, E04233, E02221, AF153205, X98834, AL133098, Z37987, AL133568, AL034417, AC004383, AR013797, AF081195, AL137283, AL080074, E05822, AL137478, Z72491, AC002464, Y09972, AL023657, I00734, M30514, E00617, E00717, E00778, AF061981, U78525, AL117432, U58996, AF061573, AF081197, AL034400, AL137294, AF095901, A18777, A90832, and E08631.
HHBEN34	108	703745	1 - 368	15 - 382	
HHBFL31	109	800035	1 - 209	15 - 223	AC007967.
HHBFW44	110	716283	1 - 144	15 - 158	
HHBFW75	111	958692	1 - 437	15 - 451	
HHBGN42	112	698774	1 - 227	15 - 241	AL119319, AL119522, AW392670, AL042551, AL119439, AL119484, AL119391, AL119401, U46350, AW372827, AL119457, U46347, U46351, AL119324, AL119418, AL119413, U46346, U46349, AL042614, AW363220, AL119439, AL119437, AW384394, AL119363, AL119444, AL119443, Z99396, AL119497, AL119355, AL119483, AL043019, AL134524, AL037205, AL134525, AL119335, U46341, AL119399, AL119396, AL134518, AL134528, AL134538, AL042544, AL042542, AL119496, U46345, U46345, AL042450, AL042896, AL042984, AL042965, AL042975, AL043029, AL043003, AL119464, AB026436, AR069079, AR060234, AR066494, AR043113, A81671, and AR054110.

HHBGN52	113	726391	1 - 266	15 - 280	AC007377.
HHBGN68	114	752745	1 - 41	15 - 55	
HHBGR37	115	708457	1 - 294	15 - 308	
HHBGT39	116	940578	1 - 54	15 - 68	
HHBGY59	117	792027	1 - 538	15 - 552	
HHBHE83	118	780875	1 - 62	15 - 76	
HHBHK08	119	958649	1 - 287	15 - 301	AL031984.
HHBHK84	120	858431	1 - 336	15 - 350	AL031719.
HHBHO63	121	906947	1 - 296	15 - 310	
HHBHP27	122	676601	1 - 433	15 - 447	Y16241, and AF047368.
HHFBD39	123	826307	1 - 219	15 - 233	AA346547, and AA346371.
HHFBD50	124	724763	1 - 287	15 - 301	AA346559, AA428211, H56371, H56372, C16964, and AC007193.
HHFBF32	125	502872	1 - 395	15 - 409	
HHFBH26	126	502843	1 - 256	15 - 270	AA047753, AA236652, and AA234860.
HHFB105	127	932961	1 - 355	15 - 369	
HHFB181	128	502954	1 - 344	15 - 358	AA347517.
HHFBL16	129	509237	1 - 387	15 - 401	Z62184, and Z62183.
HHFBL30	130	509238	1 - 341	15 - 355	
HHFBL32	131	509233	1 - 291	15 - 305	AP000687, and AP000688.
HHFBL36	132	707930	1 - 300	15 - 314	AI285706.
HHFBL39	133	509637	1 - 226	15 - 240	AC005343, and AL137658.
HHFBL60	134	739670	1 - 353	15 - 367	AA421695.
HHFBP60	135	503453	1 - 307	15 - 321	AA347874, AA346832, and AC002126.
HHFBX77	136	959805	1 - 411	15 - 425	AB023161.
HHFCA64	137	720849	1 - 312	15 - 326	AA347017, AA902341, AI356882, AA346959, and AC004948.
HHFCF70	138	518435	1 - 407	15 - 421	AA347521, and AA347377.
HHFCH59	139	526389	1 - 337	15 - 351	AA347433.
HHFC173	140	518427	1 - 198	15 - 212	AA347520, and AA347376.
HHFCK71	141	781725	1 - 227	15 - 241	AA347681, and AC005829.
HHFCL91	142	509628	1 - 340	15 - 354	AA132492.
HHFCP67	143	536062	1 - 281	15 - 295	AA347928, and AA347923.
HHFCZ01	144	509163	1 - 305	15 - 319	AA347978.
HHFDA67	145	509387	1 - 286	15 - 300	
HHFDG32	146	502957	1 - 312	15 - 326	AA347334.
HHFDH02	147	921297	1 - 377	15 - 391	
HHFDH38	148	536551	1 - 338	15 - 352	AA055018, AA346973, H71821, Z80771, AL008626, Z95331, AL031177, Z69923, AC002467,

[illegible]

					AC007161, AP000078, AL133233, AC004222, AL049767, AC004526, AC003976, Z95124, Z82210, AC005036, AL031229, AC004614, AL031119, AP000472, AC006052, AL121871, AJ006995, AC007372, AL049837, AC005261, AC007423, and AC005181.
HHFDI07	149	954404	1 - 288	15 - 302	
HHFDI42	150	500877	1 - 326	15 - 340	AA347955.
HHFDI62	151	745569	1 - 378	15 - 392	AL033527.
HHFDI66	152	573283	1 - 387	15 - 401	
HHFDI27	153	534909	1 - 427	15 - 441	
HHFDI87	154	575105	1 - 338	15 - 352	
HHFDM05	155	932863	1 - 414	15 - 428	AL037595, AA347423, and AA632734.
HHFEJ18	156	525616	1 - 246	15 - 260	AC006453.
HHFEO24	157	525611	1 - 295	15 - 309	
HHFESS1	158	525609	1 - 167	15 - 181	Z72519, and AF001905.
HHFFG41	159	575009	1 - 188	15 - 202	
HHFFO64	160	523769	1 - 316	15 - 330	AI242160, AI247651, AA775915, and AW393719.
HHFFO66	161	500910	1 - 204	15 - 218	AA347624, AA347709, and I45588.
HHFFO96	162	894076	1 - 172	15 - 186	
HHFFT69	163	662462	1 - 991	15 - 1005	AA127759, AA121515, AI984304, AI983026, AI360098, AI239945, AA961058, AA058699, and AB007972.
HHFFX20	164	525617	1 - 104	15 - 118	AL133312.
HHFFX75	165	507394	1 - 248	15 - 262	Z93024.
HHFFY80	166	732614	1 - 386	15 - 400	AA558357, and AA558032.
HHFFZ04	167	927869	1 - 531	15 - 545	AI091840, AI275088, AI088323, AA347574, AL132641, and AB007865.
HHFFZ19	168	509630	1 - 453	15 - 467	
HHFGA21	169	573584	1 - 343	15 - 357	AA501872, U91322, AC005387, AC004531, AC005253, AC005914, AC005004, AL035659, AL133245, AC005971, AC006285, AC007225, AL022163, AC002563, AL031685, AL049830, AP000359, AC005086, AC006006, AC007444, AC002094, AC004583, AC005409, AL049760, AC006064, AC006449, AL022721, AC010072, AC005520, AL031228, AL031729, AL035078, AC016025, AL121595, AC005229, AP000117, Z99716, AL022323, AL133246, AC007226, Z84469, AC009516, AC005071, AC005225, AL031390, AL096775, and AC007032.
HHFGC14	170	741650	1 - 358	15 - 372	
HHFGC69	171	573491	1 - 263	15 - 277	AC007002.
HHFGC93	172	576487	1 - 338	15 - 352	
HHFGC95	173	795968	1 - 396	15 - 410	
HHFGEO1	174	917137	1 - 394	15 - 408	
HHFGI85	175	524890	1 - 280	15 - 294	

HHFGM50	176	506635	1 - 453	15 - 467	AC005324, and AL080124.
HHFGN31	177	908508	1 - 442	15 - 456	
HHFGN67	178	573509	1 - 308	15 - 322	
HHFGP71	179	573860	1 - 402	15 - 416	Z97989.
HHFGR35	180	573510	1 - 304	15 - 318	AC005369.
HHFGS09	181	526331	1 - 403	15 - 417	AA348101, AW299437, A1640149, AW135484, A1636408, Z42797, AW269210, A1271987, AA922757, A1765884, and AL008733.
HHFGS40	182	888332	1 - 300	15 - 314	
HHFGS92	183	871899	1 - 319	15 - 333	
HHFGT10	184	968109	1 - 399	15 - 413	AC006312.
HHFGY13	185	573473	1 - 362	15 - 376	
HHFGY37	186	711364	1 - 404	15 - 418	
HHFGY75	187	573483	1 - 458	15 - 472	
HHFGZ54	188	573477	1 - 227	15 - 241	
HHFGZ63	189	661248	1 - 406	15 - 420	
HHFGZ69	190	918322	1 - 404	15 - 418	
HHFHA44	191	573498	1 - 261	15 - 275	AC005940.
HHFHC02	192	920510	1 - 449	15 - 463	
HHFHC44	193	716763	1 - 213	15 - 227	
HHFHC57	194	573506	1 - 148	15 - 162	
HHFHC72	195	766128	1 - 252	15 - 266	AA077889, and AC004084. AA524846, AW029626, A1888050, AW169469, AA230221, AL119063, A1061158, AA809104, A1821945, A1446618, H91711, R79396, AA385094, AA847341, A1537800, A1281622, A1002863, A1940546, AA947352, AA834891, AA935827, AA496941, A1570067, H98162, W27084, AA196287, AW407007, AW022796, AA362440, AA846036, AW085811, AA665102, A1884404, A1634466, H81012, AA364147, A1434103, AW157128, N49298, AW028376, A1567676, H86399, A1640905, A1268465, AA353081, AW104040, AA112864, AA349923, AW152451, N49540, AA812133, R64110, AA828840, A1520984, AA593168, AA364082, A1801479, AA020882, R92390, A1038029, AA394283, A1581006, A1708565, AA601336, A1890283, W02419, AA196994, AA613189, A1805373, C75332, AA507975, A1345256, AA761454, AA411337, AA419014, A1805556, AW131109, AW190437, A1207534, W07542, T96411, AA101744, AA618531, AA730795, AA581317, A1820978, H94598, F35374, AA804838, AA419107, AA229316, AA554289, AA779599, AA230024, A1760850, AA525324, A1174701, AA296656, A1825901, AA714190, AA436949, A1343808, AL046487, A1149177, AA828613, AL080243, AL133371, AC004883, AC006390, Z75744, AC007731, AC005500, AL021332, AL035361, AP000692, AC004812, AC004821, U85195, AE000658, AC006974, AL049780, AF139813, Z83820, AC007055, AC008040, AL022721, Z93020, AC006538, AC002106, AC006263, AC003957, AC005911, AC006261, AC004228, AC007011, AL139054, AC007263, AC005516, AC004000, AC006023, U65590.



HHFHE28	196	506630	1 - 348	15 - 362	AC007201, AC003092, AC004148, AL031229, AC001551, AC006315, AF111169, AC006360, Z85986, AC005412, AC005355, AC004673, AL035464, AL021546, AL109952, Z81364, AF047825, AF124730, AL049562, AL031121, AL117339, AF109907, AF111168, AC005245, AC002492, AC008179, AC006468, AL034420, AC007899, AC004019, AC000075, AC005664, AL122020, AC006547, AC008125, AC007390, AC005759, AC005581, AC003025, AC005993, AC005553, Z83826, AC005630, AF051976, Z86090, AL133275, AP000359, AC006006, AC004921, AL132994, AL034379, AC004881, AC006142, AF196779, AL050348, Z94801, AC005411, AL008710, AC000353, AC016026, Z95113, AC005751, AL022313, AL049643, AC006079, AL031177, AC002395, AL117330, AC005624, AC005102, AC002120, AL133245, Z97184, AC020663, AP000703, AC004099, AP000279, AC009399, AC004084, AC003036, AC006001, AC006958, AL008725, AC007934, AC005189, AP000039, AP000107, AC016025, AL109984, Z98752, AC008171, AC003999, AC005520, AL035443, AC004797, AL121658, AL022165, AP000038, AP000106, AC004835, Z97056, AC004801, AL031295, AC004477, AL008721, AF001975, AC004112, Z95116, AB023049, AC005484, AC007845, AL049759, AL008628, AP000512, AC005329, AC005180, AL050333, Z98304, AC005015, AC005482, Z99571, AL121603, AC005599, AL133448, AF001549, AF205588, AC005069, AP000247, AL023807, AL031670, AC005781, AP000076, AC003991, AL031662, AC005264, AC006125, AC002407, AC007666, AF003529, AC000052, AC016830, AC002044, AC005253, AC002430, AL109613, AL031848, AL079342, AC007676, AC005730, AC007384, AC005821, AC007686, AF107885, AC005682, AC005231, AC005066, AC012099, Z69917, AC005337, AC002558, Z84487, AC005288, AC005722, AC005529, AC012627, AC002045, AC002465, AC007066, AC007312, AL049743, Z76735, AL080317, AF196971, AC005585, U95742, AC005775, AP000280, Z93017, AC002563, and AL031257.
HHFHE28	196	506630	1 - 348	15 - 362	
HHFHE38	197	526391	1 - 512	15 - 526	R10859, AA347294, and H87292.
HHFHE76	198	506578	1 - 423	15 - 437	
HHFHF91	199	526396	1 - 357	15 - 371	AA346979.
HHFHJ46	200	576949	1 - 658	15 - 672	AI827607, AA007488, AA007421, H49821, H49565, AI241634, AW408635, and AC007371.
HHFHJ72	201	575030	1 - 363	15 - 377	
HHFHM77	202	934027	1 - 175	15 - 189	
HHFHN54	203	506624	1 - 211	15 - 225	
HHFHQ86	204	572923	1 - 396	15 - 410	
HHFHU63	205	572871	1 - 338	15 - 352	
HHFHX11	206	967321	1 - 114	15 - 128	AL022476.
HHFHY47	207	720473	1 - 339	15 - 353	
HHFHY66	208	573294	1 - 341	15 - 355	AL135049.
HHFJC02	209	918318	1 - 389	15 - 403	
HHFJ111	210	883731	1 - 285	15 - 299	

HHFJL11	211	965931	1 - 399	15 - 413	
HHFJM56	212	857990	1 - 254	15 - 268	AC005998.
HHFJN01	213	913798	1 - 436	15 - 450	AA348215.
HHFJN02	214	918358	1 - 427	15 - 441	AA346443, AC005301, and AC007064.
HHFJO06	215	934168	1 - 492	15 - 506	AA347342.
HHFJO12	216	969624	1 - 422	15 - 436	
HHFJR06	217	934093	1 - 295	15 - 309	AC005360.
HHFJX18	218	907658	1 - 328	15 - 342	
HHFKB03	219	922803	1 - 445	15 - 459	AL138381, AA353533, AA437014, and AW372500.
HHFKC28	220	857948	1 - 364	15 - 378	
HHFKC43	221	906903	1 - 467	15 - 481	AI051548, N54917, AI056521, and AA693909.
HHFKE05	222	920550	1 - 460	15 - 474	
HHFKH82	223	857810	1 - 323	15 - 337	AC000075, AC000097, AC006547, and AC005664.
HHFKI02	224	918288	1 - 408	15 - 422	
HHFKJ68	225	934045	1 - 308	15 - 322	AW375009, and AI935760.
HHFKK95	226	952082	1 - 496	15 - 510	
HHFKM10	227	963184	1 - 465	15 - 479	
HHFKU12	228	887244	1 - 408	15 - 422	T70846, T85471, and R18037.
HHFKU44	229	857923	1 - 520	15 - 534	
HHFKX11	230	965874	1 - 365	15 - 379	
HHFKY11	231	965876	1 - 455	15 - 469	
HHFLA69	232	933869	1 - 203	15 - 217	
HHFLH34	233	857907	1 - 408	15 - 422	AA502532, AA644090, AA552586, AI687343, AI369580, AI128511, N23913, AA501867, AA908687, AA504818, AA704393, AW340905, AW023111, AW169038, AI334464, AW275971, H04977, H61871, AI732151, AI366993, AI344812, AA715814, F00564, AI270559, AA427636, AA464739, AI609972, AI440117, AA502991, AW275719, AI733856, AI859438, AW265735, AL041453, AI923052, AI554471, AA559241, AA578621, AI742168, AA507822, AW271904, AA640410, AA640430, AW005974, AA877992, AI049709, AA515138, AA720774, AA481887, AW268973, AA283730, R44593, T07039, AA084609, AA570740, AA483606, H58354, AI056177, H63660, AA225406, AI580250, AI753037, AA127078, AA602557, AL048275, AI366902, AI223968, AL047429, AI754653, AA701080, AA714110, AI076228, N25272, AA484208, N72170, H53217, AI144101, AI758435, AI061313, AW150077, AW022655, AL079734, AI249683, AW007980, T06828, AA600341, AA297666, AW020150, AA362349, AI251576, AI342183, AI933714, AA586667, AA632757, AA654778, AI491765, H66577, F35374, AI417469, R95840, AA013168, AI792521, AW272294, AI597820, AI583252, AI696955, AI471914, C75034, AI345418, AA477503, AA779783, AA828802, AA828781, AW302711, AA054085, AA744018, AI446259, F05592, AA523792, AI565084, R98160, AA508809,

					AL138329, AA578832, A1635440, A1270476, A1635028, AW188742, AA829036, AA351003, AA640260, AA847395, AA584423, AA568204, AA469327, AW245354, AA749235, AC005775, AC007263, AC005666, AL022322, AL034449, AC007161, AJ246003, AC004531, AC008033, AC004408, L78810, AC006057, AL096791, Z84480, AL121603, AC006511, AL020995, AL031588, AL049795, AL031575, AL022302, AL034420, AP000692, Z99716, AC002314, AC005300, AC005690, U91326, AP000967, AC008372, AC007773, AC003971, AL121655, AC004883, AC006305, AC005988, AC006211, AC004596, AL121658, AL022163, AC005899, AL031003, AC007225, AL049709, AC005746, AC000353, AC004771, AC020663, AC005546, AC004966, AC006120, AC005736, AF001549, AC005229, AC005841, AL035604, AC003007, AC004517, M87889, AF003528, AL134445, AP000553, AC002289, U52111, AC007011, U80017, AL049779, AC002350, AC005409, AL035462, AL035249, AC004000, AC005379, AC006946, AC005696, AC005529, AC005291, M63543, AC002430, Z98051, AC006468, AD000812, AC005209, AL021707, AP000044, AP000112, AC005632, M63544, AC000392, AL049869, AP000694, AC007384, AC005480, AC009516, AC005488, AC005694, AC005335, AC004814, AC006059, AJ009610, AL023284, AC003690, AC004552, AF053356, AC002477, AL022323, AC004598, AL034379, AC004760, AC008062, AP001060, AC004686, AL078463, AL050341, AJ010597, AL133246, AL109827, AC006014, AC007193, AC007360, AL031311, AC005914, AL035659, AC012384, AC007114, AC006127, AC005011, AC004678, AF130343, AL049776, AL022238, AC006251, AC004130, AF190465, AL022165, AL121782, AC005081, AC004217, AC002551, Z97200, Z98744, Z98752, AP000556, AL035086, AL133244, AC007201, AC002425, AC005486, AP000552, AF196969, AC005102, AC004743, AC000134, AL049569, AC006101, AF165926, AC016025, AC006130, AL035420, U91318, U08988, AL049759, AC005042, AC006160, AC007191, AL035405, AC004651, AC005365, AC004491, AC007541, AC005527, AC005231, AC004757, AC005740, AC004655, AL008723, AL049830, AC007981, AC015853, AC006277, AP001050, AC002492, AC005940, Z83826, AC005288, AP000962, AC005520, AF111169, AF064861, AC005256, AL021940, Z68756, AC005411, AL031662, AC005207, AC005005, AC005031, U89335, Z97632, AF187320, AP000557, AC002984, AC003692, AC005175, AL035071, AL117694, AF165176, AC009247, AC007686, AC002126, AL109801, AC007536, AC005187, AC003043, AF027390, AC005500, and AC005962.
					U51560, and Z80896.
HHFLM06	234	933868	1 - 405	15 - 419	D57483, D80253, D58283, D80024, D80366, D59859, C14389, D51423, D59467, D80043, D59889, D80166, D59619, D59610, D80210, D51799, D80240, D81030, D80188, D80391, D80212, D59787, D80022, D59275, C14331, D80219, D80195, D80164, D80227, D59502, D80196, AA305409, D80241, D80251, C15076, D59927, D51022, D80038, AA305578, D80269, D80248, D50979, D50995, D80193, AA514188, D81026, D80133, D80378, AW177440, D51060, AA514186, C14429, C14014, AW178893, D80045, D80522, C75259, AW360811, AW352158, T03269, AW179328, D58253, AW375405, D80268, C05695, AW378532, D80134, AW177501, AW178775, AW177511, AW377671, AW177731, AW366296, AW360844, AW360817, AW375406, AW378534, AW179332, AW377672, AW179023,
HHFLP28	235	857888	1 - 437	15 - 451	

					AW178905, AW178762, D80132, D80247, D51250, D80439, AW369651, D80302, D51079, AW177505, AW352171, AW377676, AW178906, AW352170, AW179024, AW178907, AW179019, D59373, AW352117, AW176467, AW360841, F13647, AW179020, D80949, D52291, AW178909, AW177456, AW179329, AW178980, AW177733, AW378528, AW178908, AW178754, AW179018, D80168, AI910186, C14227, D81111, T11417, C14298, AW179004, D80064, D80157, AW179012, D51103, AW178914, C06015, AW378525, D51759, C14407, AI905856, AW177722, AW352174, D58246, AW179009, AW177728, AW178774, AW178911, AW378543, AW352163, Z21582, AW360834, C14975, AW178983, D59503, AW378540, AW178781, T48593, D58101, D45260, D59653, AW177723, AW352120, D51097, D59627, D80014, AI525235, D80258, H67854, AA285331, AW378533, AA809122, AI535850, D50981, AW367950, C14957, C03092, AI557751, H67866, AW367967, D80228, AW178986, AI525923, D51213, AI525917, D59474, AW177508, D51231, D51221, T03116, AW177734, D45273, AW177497, C14973, D59317, AI525920, C14344, AA514184, D60010, AI557774, AI535686, AI525227, D59551, AI525222, D60214, C14046, T03048, D80314, Z33452, AI525912, AI525215, AI525242, AW378542, AI525925, AI535961, AW378539, C05763, C16955, T02974, AI525228, AI525237, AW360855, AR018138, AR008278, A62298, AJ132110, A62300, A84916, AF058696, AB028859, Y17188, X67155, D26022, Y12724, A25909, A67220, D89785, A78862, D34614, D88547, A82595, X82626, I82448, A94995, AR060385, AB002449, AR008443, AR025207, I50126, I50132, I50128, I50133, AR066488, AR016514, AR060138, A45456, A26615, AR052274, AB012117, Y09669, A43192, A43190, AR038669, AR066487, AR066490, AR054175, A30438, A85396, D88507, AR066482, A44171, X68127, I18367, I14842, AR008277, AR008281, A85477, I19525, A86792, D50010, Y17187, X93549, A63261, X64588, AR008408, AR062872, A70867, AR016691, AR016690, U46128, D13509, A64136, A68321, AR060133, I79511, AF123263, AR032065, Z82022, and AR008382.
HHFLU04	236	870085	1 - 353	15 - 367	AA578033, AI908783, T06477, AA309342, and AL034548.
HHFMA58	237	853959	1 - 339	15 - 353	D80164, C14389, D81111, AW177440, D51799, D80195, AW369651, C15076, D80227, D80038, D59502, D80269, C14227, D81030, D80166, D59275, D59467, D58283, D80022, D59859, C14331, D80064, D80193, D59619, D80210, D80391, T03269, D80240, Z21582, D51423, D59787, AW178893, D80253, D80212, D80043, D80196, D57483, D80188, AA305409, D50979, D80219, D80378, D59927, D59610, D80366, D59889, T11417, D51022, D80024, AA305578, D80045, D50995, D80241, D51060, D80251, C14429, C14407, AW360811, D81026, AW179328, D59695, AA514188, AI557751, D59503, C05763, C06015, C14014, AW176467, AW375405, D58246, AW360817, AA285331, C75259, C14298, AW377671, D80248, D80168, AW378533, AW366296, AW360844, AW378532, D58101, D80258, D51250, D52291, D80522, AW178775, AW178762, T03116, AA514186, D80133, AW360841, AW177508, C05695, D80949, D80268, AA809122, AW177497, D80014, AI910186, AW375406, AW352158, D59627, AW378534, AW179332, AW377672, AW179023, AW178905, AI535686, AW177456, AW352171, AW177734, D59551, C03092, T02974, H67854, D59653, H67866, AW178907, AW352170, AW177731, AW177505, AW178906, C14077, D80439, AW377676, AW179019,

[illegible]

HHFMB01	238	914951	1 - 295	15 - 309	AW179024, AW179020, D80247, D51103, D80302, AW178909, D59317, F13647, AW179329, AW178980, AW179018, AW178914, A1525903, AW177733, AW378528, AW178754, AW178986, C14973, A1535961, AW179004, D45260, D51759, AW177728, A1525920, AW179012, AW378525, D59474, A151221, C14344, D60010, A1525923, D60214, A1525917, A1525227, AW352163, A1525925, C14957, A1557774, AW177722, AW378543, AA514184, AW179009, AW360834, C14046, AW367950, D59373, AW178774, A1525242, A1525235, D45273, A1525912, AW178911, AW352120, A1525215, AW378542, AW178908, T48593, C16955, Z33452, A1525237, C13958, AW178781, T02868, D31458, Z30160, C04682, AW360855, D51213, AW378540, D51053, D59976, A1535959, A1525222, AC004477, A84916, A132110, A62298, A62300, AR018138, AR008278, AB028859, I82448, AF058696, Y17188, D26022, X67155, A67220, D89785, A78862, D34614, A25909, AR060385, AR054175, I14842, A82595, Y12724, AB002449, AR016808, I50126, I50132, I50128, I50133, AR008277, AR008281, A94995, AR008443, I79511, AR016514, Y17187, AR060138, A45456, A26615, AR052274, A43192, A63261, A43190, AR038669, AR062872, A64136, A68321, A70867, AR066488, Y09669, AR066487, A30438, X68127, AR016691, AR016690, U46128, AF123263, AR032065, AR060382, and AR008408.
HHFME70	239	926497	1 - 353	15 - 367	AA984829, AA558487, AA492161, AA046737, A1049701, AA603558, AL035686, AC016027, AC016830, AC002565, AC004168, AL031120, U95740, Z98941, AC007546, AC005841, AL132826, AL035555, AC005755, AP000694, AC006088, AC002365, AC004821, AL008730, AC002375, AC004805, AC020663, AL023575, AC006597, AC007021, AC008975, AC005182, AC006449, AL109802, Z99758, Z74739, U85195, AC004685, AE000658, AC004701, AC003950, AL133500, AC006480, AL021155, AC007057, AC005578, AC002404, AC005535, Z84469, AC006539, Z93017, AB003151, AP000688, AF111168, AC005081, AC004858, AF109907, AL031311, AC005972, AC005369, AF001548, AC009363, Z97630, AC002425, AL031602, AL109623, AL009051, AP000555, AC007283, AC005914, AL021368, AL031774, Z73988, AF108083, AC005486, AC007444, AL050347, AC005484, AC005934, AL034420, AC005037, AC005015, AC002546, AC002123, U78027, AL035422, AL121595, AC005261, AP000248, AC007731, AP000045, AC005231, AC005500, AC004946, AC002126, U95742, AC007216, AC004605, AC004409, AC005003, AC005520, AC005071, U62293, and AC009399.
HHFMMH85	240	872838	1 - 277	15 - 291	R05702, AA479449, T97899, A1040466, A180508, and T10419.
HHFMI58	241	857856	1 - 529	15 - 543	AA077459.
HHFMO03	242	922708	1 - 412	15 - 426	H06469, R18079, AA489067, AA505171, AA505170, AA346630, AA889356, AA629200, AA534443, AA488820, and Z84474.
HHFMO94	243	957955	1 - 395	15 - 409	A1911241, T72210, T87002, R08341, R08392, T85654, T99624, T99625, R02406, R07180, R59607, H62539, H99689, W63788, AA083979, AA102095, AA131697, AA131541, AA148259, AA151763, AA193069, AA220926, AA416768, AA430164, AA483282, AA6133379, AA747801, C00103, AA292662, AA401454, AA402018, AA625743, AA405177, AA773305, AA970146, AA985342, AA985358, A1004081, T19343, F07893, A1301484, and A1190719.

HHFMQ70	244	840039	1 - 320	15 - 334	AI284640, AW301350, AW303196, AW407578, AL119691, AW088202, H71429, AW103758, AW274349, AL046409, AW029038, AL119984, AW247819, AA469451, AI076616, AW021583, AI583466, AW276435, AA720702, AI783494, AI929531, AA584201, AA502860, AW276817, AI339850, AI687343, AA488746, AA581903, AI358571, AI434695, AI144101, AI038785, AW265385, AW405021, AI267818, AA584752, AI537955, AL042853, AI457397, AA491814, AL138265, AW302013, AI017415, AI589230, AA847499, AW276827, AA323644, AI679782, AI368745, AW304584, AI312309, AI357551, AA826303, AW103981, AI251002, AW271904, F36273, AI708009, AW162049, AI745325, AI357288, AI364809, AI064952, AI281697, AI246119, T09071, AW238278, AI890348, AI610607, AI653636, AW089322, AA862180, AW023270, AW028392, AI251576, AI801600, AI821271, AL120269, AI568678, AI281754, AI903462, AI10688, AA586661, D83989, AI121658, AP000208, AP000130, AP000247, AC005703, AC006004, AL009181, AC016831, AC004030, AF015156, AF124523, Z98050, AF042090, AL109985, AC004986, Z82189, AC003006, AC007666, AL031577, AL049869, AL080278, AP000313, AC004149, AC009044, AC007226, AC005901, AL008718, AL034420, AC003962, AC005323, AC002544, AC002303, AC005899, Z97352, AC006958, AC004895, Z69917, AL121603, AP000050, AC007262, AP000049, AC002072, AC004584, AC005217, AL022329, AP000555, AL132987, AL133304, AC004447, AC007564, AC007462, AP000117, AC007298, AC004832, AL035411, AC004040, AL035423, AC004593, AL078593, AP000311, AI239322, AL050341, AL034379, AF001549, AC005046, AC004987, AL008582, AC002404, AL034451, AC007363, AP000245, AC008079, AP000459, AL035468, AC010722, AC003109, AI006996, AC005529, AL133445, AC004638, AL096805, AC005295, AC006037, AL008629, AC004158, U66059, AC006285, AL135744, AC003684, AC006238, AC005242, AC004417, AL031283, AC007687, AL031054, AC005694, AC002379, AL021368, AL049691, AC004386, AC004653, AC004883, AC009247, AL079300, AC005229, AC006456, AC007128, AC004940, AL133246, AL034373, AP000088, AL035448, AP000228, U91323, AC004692, AC009516, AC008169, AC007537, AC008372, AC007055, AC004804, AC002429, AJ010598, AC004815, AC004884, AC005082, AC000118, AC005197, AL023803, AL121934, AL022163, AC005907, AC004972, AC006288, AL050348, AC004858, AC002067, AC002456, AP000140, AC007237, AL022323, AL031584, AC006006, Z97181, AF015151, AL022238, AL031848, AC003982, Z68870, AC005527, AC005998, AJ246003, AC007216, AC007324, AC000379, AC007685, AC007773, AF195658, U07562, Z97195, Z84480, AC004383, AP000193, Z93023, AC005144, AC004686, AC007845, AC005520, AL033525, AC005324, AL049570, AL122126, Z95152, U91326, AC007030, AL035089, AC002395, AC007637, AC002369, AC005919, Z82190, AC005164, AC005228, AC003042, AC003101, AC005154, AL133371, AC011311, AL096701, AL022328, AC005829, AC016025, AC007486, AC005562, AC010168, AC016026, AC005799, AF109907, AL022723, AC005104, AL031446, AC002430, and
HHFMZ40	245	974299	1 - 565	15 - 579	
HHFNA49	246	914838	1 - 319	15 - 333	
HHFND10	247	963140	1 - 437	15 - 451	

HHFNF41	248	857825	1 - 381	15 - 395	AC006442.
HHFNI49	249	857850	1 - 304	15 - 318	AP000355.
HHFOC27	250	857816	1 - 390	15 - 404	AA346789.
HHFOC66	251	857955	1 - 228	15 - 242	AL049835.
					AW160814, AW452242, AI393255, AI356203, AW290992, AI640362, AI640129, AI365982, AF088886, AF132894, AF136279, and AI007331.
HHFOF07	252	952062	1 - 210	15 - 224	AL132987.
HHFOF40	253	930839	1 - 361	15 - 375	
HHFOJ05	254	930856	1 - 225	15 - 239	AA347058, and AC000015.
HHFON08	255	957974	1 - 218	15 - 232	AL031282.
HHFON15	256	857792	1 - 245	15 - 259	
HHFON19	257	910891	1 - 1126	15 - 1140	AA527292, AA915932, AI215158, AA926744, AI283262, AA356476, AI916043, AI470481, AW372085, AW384295, AW372101, AW384294, AW372104, and AC003072.
HHFON32	258	858034	1 - 439	15 - 453	
HHFON87	259	857783	1 - 327	15 - 341	AA507612, and AL133245.
HHFOW08	260	857772	1 - 337	15 - 351	AA346793, AA494051, R96249, AI932599, AI672860, TI6700, H09486, H07125, AA569178, AC006449, AC004703, AC004386, AC005231, AC005274, AF031078, AF030876, AB016897, AC006511, AC003071, AC002565, AL033547, AC005412, AC002400, AB011399, AI009181, Z95114, AC006512, AC005529, AC002543, AC005081, Z83826, Z93023, AL049793, AC004876, AC004522, AL139054, AC005049, AC002288, AL049758, AC006050, AC004228, AP000010, AC004953, AF047825, AC006064, AL080243, AC005058, U47924, AC006538, AF053356, AL031680, AL049761, AL008719, AC007011, AL049712, AC006012, AL031255, AC004882, and AC007308.
HHFSB02	261	921364	1 - 297	15 - 311	AA347106.
HHFTC88	262	500903	1 - 345	15 - 359	AA347696.
HHFUB15	263	534520	1 - 300	15 - 314	AA776649.
HHFUB23	264	928063	1 - 375	15 - 389	AC004985.
HHFUB77	265	772691	1 - 291	15 - 305	AW080134, AA515051, AA229785, AI339850, AA482711, AI929531, AI284640, F29989, AI282832, AI205126, AW029038, AI962050, AW023672, AI358229, AI469003, AI251002, AA745582, AA826303, AI358343, AA843450, AW088202, AW406755, AI049722, AA347927, AI049634, AA594145, AA074130, AW440836, AI365988, AW265294, AA318652, AI625244, AW276827, AI151261, AW193432, AI569086, F28576, AI919029, AA503258, AA469451, AA347930, AI206785, T29180, AI198376, AA649642, AW162049, AW339687, AW166815, AI918421, AL138265, AW129001, AI890923, AI688846, AA441788, AA985038, AI635272, T06828, AI133636, AI357551, AA747276, AI567674, AA568778, AA350859, AI382614, AI350211, AA747480, AA720025, AI053672, AW193265, AI537955, AI446464, AW440976, AW276435, AW236342, AA364456, AA515435, AW021583, AA492132, AI298710, T06556, AA503283, AI679782, AA515751, AA713815,

AI904894, F25867, AW302450, AI445498, AW301350, AI570261, AW303196, AW274349, AI654529,  
 AA483068, AI580652, AI370475, AA533408, AA394271, AI079910, AA449088, AA372478,  
 AW243960, AA501784, T07451, AI469624, AI344844, AA309841, AA553465, AI610159, AW079659,  
 AI653905, AA299115, T08914, AL046409, AI917271, AW238016, AA603323, AI144055, AA654979,  
 AI250019, AA886584, H64777, F36273, AW438643, AI471481, AA148489, AA669251, AA244415,  
 AI471543, AI281697, AA533725, AA557879, AA593247, AI200051, AI120674, AA659083,  
 AL038785, AI286264, AA747472, AW079241, C75026, AA215929, AI431303, AW302013, AL043721,  
 AI192631, AI281881, AI608626, AI357288, AI902694, AI354397, H62778, AA515224, H84359,  
 AW265385, AW419262, AW089810, AW088058, AW173651, AA573020, AI623898, AA502155,  
 F28776, H64560, H63607, AI471534, AA100599, AA161181, AI061334, F18974, AA044863,  
 AI648558, AW265654, AI824562, F17700, AA827981, AI287651, AW089322, AI345654, AW242109,  
 R91994, AA364420, AW338500, AI266133, AI305547, AI312309, AW265253, AI683577, AA338486,  
 AW339568, AA352803, F25203, AI289067, AA601492, AI500454, AW020992, AI358813, AI351698,  
 F03525, AI299050, AA559290, AW419118, AI434695, AA903367, AA503454, AI866856, AA650244,  
 T08438, F19012, AI017024, AI364809, AA513884, AI4718, U14719, U14714, AF015160, U14715,  
 U47924, U14713, S75337, M87916, U14717, U14711, U14712, A39972, AL049635, AF015171,  
 AC007436, AL049869, AC006030, U14685, U14686, AL109801, U57005, U14691, U57007, S77605,  
 AF020503, U14695, U14705, U14688, U14689, U14687, U14707, D83989, AC006221, X54176,  
 X75335, X55925, I51997, AF200465, Z73429, AC004866, Z22650, AL031224, AC005618, AL021391,  
 U14706, U14716, AF140763, AC007751, U14700, AC005670, U14699, U14710, U14684, U14690,  
 AC007845, U12580, M87919, S70694, AC016025, U57009, U18399, U18391, U57008, AC004506,  
 AC006116, AP000959, AF001552, U18390, AC005252, AL133445, AC005048, AF015158, U92032,  
 AC005041, AF015169, AL031662, AC004019, U14692, AF172277, AF015162, AL049836, AC006039,  
 U14702, Z80899, AP000966, AC005145, AL021878, AC003029, X54180, U18392, U57006, U18394,  
 Z98256, U18393, AL049554, X55928, AL022238, AF107885, AC007283, S70697, AC002984,  
 AF015167, Z94721, X54175, AC005401, U18398, AC005064, AC003015, Z98200, U18396, AC008282,  
 X55931, AC006237, AC004159, Z97198, Z30979, Z83826, AF015168, AC004030, AC006064,  
 AP000011, AL117693, U91321, AL031230, AL035400, AF145473, AP000126, AP000204, Z98744,  
 AC006115, AC004087, AL008718, Z82244, AL050305, AC006441, AC000086, AC005912, AF147277,  
 AL023513, AC006312, AL050318, Z84484, AC005253, AC006132, U14703, U14701, AC005776,  
 X54181, AC005177, U18395, AC005324, AC005696, AC006547, AP000244, U14704, AC005229,  
 AC006371, AL022237, AL031280, S48196, AC005255, X54178, X55924, AL021939, AC007040,  
 X55932, AC004534, S70707, Z98043, U12583, X53795, U80460, U02058, AC007242, U02063,  
 AF015170, AC004187, AC007536, AF015166, U73635, AP000337, U67221, AL139054, X55930,  
 AC003035, AF031078, AC005065, AC004806, AC006539, AC006254, AF030876, AC004686,  
 AP000156, AC004167, Z98745, AC002429, AC006207, AC007162, AL021154, U12582, AR036572,  
 U91328, AL023576, Z69921, AF015163, AL035423, AC005484, AC006296, AL121852, AP000954,



HHFUC24	266	524840	1 - 354	15 - 368	AL031347, AL023494, AC005529, AC007766, AL035551, AL031427, AF095855, AC005678, AC007684, AC001164, AC006046, AC005666, AC004825, U85195, AF141976, AL049776, AC004964, U93163, AC005185, AJ003147, AC005858, AC006076, AF015725, AC010202, AL117355, AC004193, AC000097, AC003001, AF077058, AL078593, X55926, Z31005, X55923, AC006127, AC004130, Z82208, AC004887, U18387, and M87917.
HHFUC26	267	960331	1 - 371	15 - 385	AC007686.
HHFUC42	268	525603	1 - 313	15 - 327	AW135348, AW204310, AA96205, AI697749, AC004836, and AC005069.
HHFUC45	269	525600	1 - 198	15 - 212	
HHFUC47	270	525599	1 - 334	15 - 348	
HHFUC83	271	507459	1 - 488	15 - 502	
HHFUC92	272	507018	1 - 329	15 - 343	AA347389, AA280938, AL079866, AI936976, and AB011134.
HHFUC58	273	525604	1 - 191	15 - 205	
HHFUL75	274	675569	1 - 343	15 - 357	
HHFUN56	275	530953	1 - 210	15 - 224	
HMEBA75	276	530303	1 - 146	15 - 160	
HMEBY61	277	947868	1 - 385	15 - 399	AA303737.
HMEDF58	278	738427	1 - 480	15 - 494	
HMEDR48	279	529173	1 - 356	15 - 370	
HMEEF64	280	918746	1 - 377	15 - 391	T99598.
HMEEX61	281	742085	1 - 487	15 - 501	AA173199, and AA347039.
HMEFR17	282	664438	1 - 459	15 - 473	H29936, H29845, W27872, AW380042, AF085864, and AB028981.
HMEFX12	283	661951	1 - 446	15 - 460	AA339614, and AA374928.
HMEGB93	284	573814	1 - 352	15 - 366	AF093097.
HMEGF48	285	573821	1 - 404	15 - 418	
HMEGG44	286	796421	1 - 445	15 - 459	N44889, and AL110204.
HMEGH46	287	887791	1 - 363	15 - 377	AB033054.
HMEGI07	288	953815	1 - 556	15 - 570	AW363121, R24201, AW025914, and AA770692.
HMEIG42	289	931114	1 - 394	15 - 408	AW192773, AA953251, AW372190, AW274877, AI669009, and AB014592.
HMEIM40	290	523589	1 - 211	15 - 225	
HMEIU49	291	722988	1 - 359	15 - 373	
HMEIW23	292	682863	1 - 263	15 - 277	AL110334, AW411094, AL135489, AL135458, AA136879, AA876722, AA719995, AI053577, R98359, AI494405, AA381147, AA483204, AW151713, AA953228, AA834707, H96467, AA970967, AC005156, AC005232, AC004527, AL008638, AC005477, AC006205, AC009501, AL022165, AP000689, AC004559, AF067845, AC007298, AF110824, AC005225, AC004821, AC006597, AL020997, AC005520, AC004491, AC005899, AC008282, AB016897, AC005368, AC004542,

					AC004463, Z94161, AC004217, AL022336, U95743, AL022718, AC004694, AP000357, AL135959, AC006039, AL031257, AL035681, AC004973, AC005786, AC005523, AL008730, AC004601, AC006044, AC005684, AC022517, AC006013, AC000134, AC005004, AC004890, AP000351, AC004924, AC005519, AC005261, Z68162, AL079304, AC006486, AL132987, AC005081, AC004778, AC003101, AC008009, AC005736, AC005015, M93189, U73647, AC004755, AC000387, AC005091, AC005222, AL049747, U93305, AC006040, AC006273, AC006449, AC012384, AC006116, Z95114, AP000270, AL225085, AC000025, X96421, and AL133448.
HMEJD13	293	657231	1 - 257	15 - 271	AC003976, AC005702, AC005562, and AL117394.
HMEJH84	294	781983	1 - 124	15 - 138	Z98744.
HMEJU60	295	740392	1 - 322	15 - 336	
HMEKA53	296	711664	1 - 246	15 - 260	
HMEKJ43	297	715893	1 - 367	15 - 381	H25268, and Z83309.
HMEKS76	298	767517	1 - 249	15 - 263	AA505833, and AC006960.
HMEKX51	299	727154	1 - 302	15 - 316	
HMEKX89	300	786055	1 - 70	15 - 84	
HMELC56	301	745773	1 - 306	15 - 320	
HMEQLQ62	302	719681	1 - 359	15 - 373	AA166769, W94197, AA166767, AA460380, T97509, N88106, AL751026, N90626, AA356483, AW340411, AW407849, AA446049, W94104, AL814281, AL085368, W01030, AL762027, AL693754, AW001517, AA704878, AL023779, AA166957, AA704992, AF083248, and AC002536.
HMELR10	303	964629	1 - 322	15 - 336	AF174601, and AB020682.
HMELS59	304	720341	1 - 307	15 - 321	
HMELV19	305	668665	1 - 266	15 - 280	
HULAF89	306	791261	1 - 86	15 - 100	
HULAI37	307	708923	1 - 160	15 - 174	AW302954, AW301754, AW268302, AL345477, AW072719, AL336582, AW268122, AL308032, AW301513, AL310575, AW074993, AW302965, AL312152, AW269097, AL344928, AL036980, AL307543, AW301300, AL307708, AW075207, AW301505, AL343037, AL336495, AW268072, AL335208, AL310582, AL307520, AL313320, AL336633, AL345677, AL340582, AL334450, AL312399, AL345608, AW269083, AL251963, AL344789, AW074869, AW071417, AW071380, AW301500, AL348897, AL345471, AL340511, AL344785, AL309401, AL036214, AL334930, AL345745, AL345347, AW075084, AL334889, AL340533, AL349967, AL336513, AL311159, AL348895, AL336654, AL345735, AL343030, AL345562, AL345817, AL349028, AL349637, AL345397, AL349645, AW302973, AW268060, AL312428, AL251221, AL802833, AL345111, AL334445, AL335214, AL345224, AL366992, AL336503, AL307210, AL345739, AL345587, AL307494, AW301375, AL366968, AW072484, AL366974, AL345737, AL366959, AL345736, AL343062, AL343059, AL348847, AL349614, AL349933, AL340664, AL308035, AL349256, AL345253, AL343112, AL345026, AW071412, AW268253, AW071362, AL349598, AL309431, AL312325, AL348854, AL340659, AL335476, AL340627, AL309443, AL348884, AL345251, AL345666.

AI345156, AI344819, AI345553, AI366985, AI312143, AI310940, AI349772, AI313352, AI345746,  
 AI349955, AW075093, AI345114, AI307542, AI312357, AI312146, AI312339, AW303075, AI345258,  
 AI307569, AI310920, AI345567, AI307503, AI311604, AI334913, AI251205, AI344808, AI344826,  
 AI336662, AI344260, AI343091, AW301344, AW268221, AI250070, AI345148, AI312165, AW303061,  
 AI307446, AI311892, AI344933, AI343140, AI349276, AI348870, AW268261, AI349937, AI349246,  
 AI802826, AI252023, AW303238, AI310925, AI307454, AI318569, AI348969, AI345674, AI312168,  
 AI345315, AI340644, AI345639, AW071276, AI037521, AI335209, AI336585, AI310606, AI349960,  
 AI345229, AI349787, AW268067, AI348879, AI349269, AI307557, AI344817, AW303160, AI340519,  
 AI335235, AI340603, AI349186, AI318159, AI335449, AI349644, AI344938, AI349971, AI345551,  
 AI345370, AI345527, AI310927, AI349622, AI307578, AL039086, AI349266, AW301459, AI312156,  
 AI336506, AW268220, AI344935, AI335380, AI254251, AI348901, AI349812, AI349004, AI310945,  
 AI345787, AI307736, AI310592, AI307734, AI312264, AI334920, AI349738, AI334452, AI345557,  
 AW071349, AW271119, AI340537, AW301815, AW302854, AI349628, AI349288, AI340610,  
 AI307459, AI349762, AI344779, AI349245, AI613548, AI307456, AI309392, AI344780, Y11587,  
 AF111112, I89947, AR038854, A08913, I48978, X72889, A08916, U80742, A08910, I89931, A08909,  
 I49625, A65341, AF061795, AF151685, AF125948, A08912, AF008439, AF017437, A90832, A77033,  
 A77035, AL080074, AL080124, AL122093, AF113019, AL049452, AF185576, U35846, AF026816,  
 AL137292, AL137463, X82434, AL137550, AF026124, AL110280, AF162270, AF091084, Y11254,  
 U49434, AL133016, AF118094, AL049938, AL117460, AF113689, AL137271, I48979, AF111849,  
 A03736, A58524, A58523, AF113690, L19437, AF067728, AL080159, Z82022, AL110221, AF090900,  
 AF113691, E02221, X53587, AF079763, AJ238278, AL133565, AL050277, AF061943, S79832,  
 AL050024, AF113699, AL137560, AF111851, Y09972, AF113013, AL050146, AF078844, AL137556,  
 AF090934, AF113677, AF097996, U00763, AF087943, Z72491, AF022363, AL137459, U42766,  
 AL137480, X96540, AR011880, AL133067, E03348, AL137557, U67938, AL049314, U58996, Y14314,  
 AL117440, AL049464, AL049466, I03321, AL133640, L30117, AF090903, AF158248, S68736,  
 AL117435, AF061573, E02349, AF183393, A07647, AL050149, AL050116, L31396, AL137533,  
 U68387, AF146568, I00734, AL050092, AL137521, L31397, AL133560, U72620, AF113694,  
 AF090943, E04233, AL137478, AL049430, AF153205, A93350, E00617, E00717, E00778, AF177401,  
 U78525, AF090901, Y07905, AL050393, AL122121, AJ012755, AL137476, AL122123, AB019565,  
 AL133104, AF003737, AF051325, X70685, M30514, AL137648, AJ242859, AL117585, E07108,  
 AF125949, AL110225, AL117394, AI2297, AL133606, X63574, AJ006417, U91329, AF057300,  
 AF057299, X98834, AL049300, I33392, X84990, AL122098, AF017152, AL133075, AL133014,  
 X93495, S61953, I42402, AR059958, E15569, AF113676, AF061981, AF090896, AL080148,  
 AL117432, AL122110, AL117583, AL137538, AL133557, S78214, AL133010, X65873, AF032666,  
 AL080086, AF119337, I26207, AR038969, Y16645, S36676, AL049283, AL110196, AJ000937,  
 AL133080, AR000496, AL049382, U39656, AF210052, AC004383, AL117457, E08631, AL023657,  
 AL050108, AL122118, AL133113, AL133072, AF079765, AF104032, AL080060, AR020905,

HULAX31	308	868930	1 - 205	15 - 219	AL122049, AL137283, AL137526, Y10080, AF118064, I09360, AL133093, X87582, E05822, AF118070, AL122050, X62580, AF132676, AF061836, AL133098, X92070, AL080127, U96683, A45787, AL096744, AL133077, AL133568, AL080137, AL137527, AL050138, AF106862, AF081197, E08263, E08264, E07361, A93016, A08908, AL117416, AL133665, AL137488, I41145, and E06743.
HULBU59	309	636253	1 - 453	15 - 467	AA362604, AF150274, AL038735, AI348980, AL048672, AL048673, AL048657, AF150161, AL037601, AI224190, AL038991, AF150181, AL037348, AL038038, AF174394, AR067468, AF100694, AL080096, AF118386, AL049963, AF125570, E04157, and AL080106.
HULBY15	310	659557	1 - 422	15 - 436	AL134132, AL043003, AL043147, and U46347.
HULDF69	311	754381	1 - 243	15 - 257	
HULFA03	312	918691	1 - 372	15 - 386	
HULFB76	313	767873	1 - 196	15 - 210	
HUMBE61	314	838469	1 - 400	15 - 414	AF150274, AL038735, AI348980, AL048657, AL037601, AL048673, AL048672, AF150161, AL038991, AI613343, AI224190, AI431323, AF174394, AF100694, AL080096, AF125570, AL049963, AF118386, AL080106, AL133076, AL133074, and Y17793.
HUSAY26	315	527909	1 - 360	15 - 374	AI858923.
HUSCA57	316	527800	1 - 250	15 - 264	
HUSCD26	317	527799	1 - 347	15 - 361	
HUSCH45	318	507197	1 - 354	15 - 368	AL022323.
HUSCH77	319	527801	1 - 117	15 - 131	
HUSGH88	320	871776	1 - 689	15 - 703	AI653117, AA922010, AW008076, AI127949, AI655942, AI142458, AI701082, AI608993, W46422, N37061, AI858207, AI273519, AA977111, N24696, AI985411, AA233273, AW137630, N72855, AI919359, AA302412, Z46073, H07031, AA232970, AI742640, and F05394.
HUSGK23	321	675951	1 - 303	15 - 317	AW389364.
HUSGL13	322	575798	1 - 306	15 - 320	AA776615, AI394186, AI623747, AI473932, AI744215, N48461, N75917, AW272349, AW088808, AI920827, AW081119, and AB011139.
HUSHH64	323	576459	1 - 434	15 - 448	
HUSHJ55	324	576381	1 - 324	15 - 338	AA985295, AJ012590, and Z98044.
HUSHL83	325	868883	1 - 335	15 - 349	
HUSIA38	326	709472	1 - 421	15 - 435	AA035379, AI494330, AA035378, AA446855, AA325787, C14480, AA487892, N91747, AW439703, AA700664, AA809787, H16875, W73597, N23062, AA358852, AA206062, AI871691, AW341955, AW408596, AA347237, AA085902, AA368749, AA111870, AA633039, AI680547, AL134167, AI206078, AW151761, AI922850, AP000252, AP000212, AP000134, AP000030, AC005562, Y08864, AC004913, AP000152, AC007298, AC005598, AC005342, AP000563, AC005193, AP000141, AP000088, D87675, AP000133, AP000211, AL096712, AC004834, AC007510, AC006480, AC008115, AC004922, X94912, AP000350, AC007227, AC006014, AC005488, AC007425, AC003109, AC007676,

HUSIA43	327	575768				AC002470, AL030997, AL096818, AC002115, AC002365, AF111168, AC005325, AC006120, AL049631, Z84469, AP000227, AL079301, AP000087, AC004893, AC005756, AL117328, AC006130, AC005207, AC002364, AC005940, AC007687, AC002094, AC005821, AC002310, AF016898, AC004659, AC007182, AL049636, AC004962, AL034429, AP000340, AP000156, AP000014, AC006326, Z49258, AC004799, Z69714, AC005365, AC000105, AC002558, AL023553, AL080317, U91326, AC007057, AL049872, AC004032, AC004223, AC004983, U91323, Z98742, AC005899, Z69921, AC003004, Z68326, AF003626, AC000066, AP000692, AC004815, AC003982, AC006343, AC005180, AC007934, AJ010770, AC008154, AC007276, AL121577, AL080243, AC007156, AC012384, Z95331, U95742, AC003665, AC006285, AC005666, AC004216, AC004548, AL049697, AC007546, AC002352, AC002472, AC004383, AL049713, AC005102, AC004765, AL137191, AL049871, Z98949, L78833, AL031680, AC006449, Y10196, AL049759, AC006000, AC003010, AL049539, AC007540, AC006071, AC005264, AC005262, AF045555, AL009181, Z96811, AC005081, AC005519, AC002477, AC006080, AL049538, AL035079, AL031311, AC007011, AC005736, D84394, AC007308, AC004106, AB000462, AF130343, AD000833, and Z68279.
HUSIA43	327	575768	1 - 418	15 - 432		AI350419, AI563971, AI208906, AW198165, AL135224, AA974279, AI223023, AI049928, AI283630, AI097446, AI648516, AI016385, AA324880, AW248808, AI991136, AW085678, AI937038, AI364515, AA709404, AA639864, AW250882, AI160941, AA854011, AL135233, AA552424, AA485734, AA583601, AA769102, AA250968, AA552593, H17500, H98072, H18108, AA678516, H38440, AW190935, R88426, AA765348, R23499, H41112, H50888, W92991, W52172, AA151984, H52131, AA809403, T54339, H46149, AA100930, D29154, D53226, R89522, AA678890, and AL035681.
HUSIS60	329	727153	1 - 343	15 - 357		AI193047, AI203647, AA757210, AI130814, AI344478, AA582236, AI288858, AI148643, AI146994, AA524479, W45709, AI168644, AA453739, AA491660, AA730161, AA557746, AI750017, AA927666, W44356, AI610804, AI685002, AI190610, AA652628, AI141208, AA490862, AA025400, AA316493, AI475353, AI859485, AA453820, Z99396, AL038837, AL037051, AL036725, AI859458, AL039074, AA631969, AL039085, AL039564, AL039156, AL039108, AL039109, AL039128, AL039659, AL038531, AL039625, AL039648, AL045337, AL039678, AL039629, AL039150, AL037526, AL039423, AL036924, AL040992, AL042909, AL037726, AL036858, AL039410, AL038447, AL036196, AL037094, AL037639, AL036238, AL045353, AL036190, AL036973, AL038851, AL036767, AL037082, AL044407, AW392670, AL037615, AL039386, AL036117, AL036733, AL037077, AL039440, AL036268, AL039924, AL036418, AL037085, AL038509, U46347, AL037021, AL036998, AL036679, AL037027, AL037178, AL039538, AL036191, AL044530, AL036765, AL037016, AL036719, AW384394, AL037054, AW363220, AL119484, AL036158, AL038821, AL043003, AL036964, AL038520, U46351, AW372827, AL119457, AL119443, AL119439, AL119497, U46350, AL119319, AL119324, AL119483, U46349, AL119391, AL119444, AL119335, AL119522, AL119418, AL119363, AL119355, U46341, AL119341, AL119396, AL134132, AL134525, AL037205,
HUSIW10	330	963324	1 - 380	15 - 394		

HUSJW78	331	772956	1 - 403	15 - 417	<p>AL119496, AL134530, AL134519, AL043147, AL134531, AL119401, AL134527, AL134528, AL039566, U46346, AL042614, AL036836, AL134533, U46345, AL119399, AL042984, AL042965, AL042975, AL042542, AL134538, AL042544, AL042989, AL043019, AL043029, AL042450, AL142134, AL042551, AL036808, AL119464, AL036886, AL036774, AR060234, AR066494, AR064706, AR023813, Y11449, Y11447, AR064707, A81671, AB026436, AR054110, AR069079, and Y11458.</p> <p>AL273114, AA661583, AI571094, F29968, N53076, AI281622, AA602664, AA493546, AW023975, AA364147, AA314404, AA159137, AW338633, AA493245, AA015948, AI002863, AI701898, AA010265, AA557508, R79396, H79004, AA807704, AI570067, AA143703, AA837006, R95751, R92703, F25759, AA018258, AA578711, AI921744, AA467740, AA832104, AA935827, AA219166, AA467807, AA525753, AI434103, AI734119, AA640776, AI167715, AA749196, R99617, AI885896, AW023390, AA527633, AA847341, AA629668, AI732720, AI802268, AA601290, AA581317, N55296, AI791659, AA832108, AA730530, AI308529, AA468982, AA584756, AC005182, AL031983, Z83844, AC006137, AC007684, AL096701, AC006487, AC005231, AC005183, AL008582, AC004491, AC003109, AC005527, AC006261, AL031670, AC005821, AC002991, AL033521, AC006449, Z92542, AP000130, AP000208, AC004531, AP001068, U85195, AC005229, AP000247, AF001549, AC007283, AE000658, AC005346, AC008009, AF111167, AL121658, AL031666, AC006006, AC000025, AL096817, AC004099, AL009181, AL117354, AL022238, AC004821, AL021808, AC003102, AC000353, AB023048, AC002039, AJ011930, AL050318, AP000511, AL096712, AC005332, Z86090, AC002477, AC002400, AC009542, AC002404, AF111168, AC006101, AC008372, AC009330, AC005261, AL035659, Z97876, AP000350, AC005920, AL034421, AC005696, AC004687, AC005694, AC005752, AC004895, AL031848, AF038458, AC005355, AL034400, AC007388, AC007066, AC007041, AC005202, AC005914, AL122020, AF207550, AP000961, AJ246003, AC002464, AF015720, AP000036, AC005529, AL049694, AC005200, AC006430, AL133312, AC000159, AL008725, AC003006, AL117694, AL121595, AL136504, AC008125, AC005049, AC005071, AC004883, AC006450, AL109627, AC007021, AC005632, AC006509, Z97056, AC002554, AL031280, AF109907, U91326, AC005323, AD000092, AC006088, AC005488, AC006581, Z98949, AC003684, L44140, AC004448, AF053356, AP000555, AC004000, AL024498, AC004526, AL020997, AC006538, AL121655, AC008044, AC005280, AC006121, AC002312, AC006162, AC004134, AL139054, AC005859, AC009501, AL031005, AC005921, AC002996, Z95114, Z82248, AL034420, AC004659, AC006468, AC006946, AL132992, Z98044, AC005041, AC004792, U96629, AC009516, AL049764, AC004148, U52112, AC002126, AJ229041, AL049569, Z97352, AC004522, AC002504, AL049757, AC007382, AC006080, AL021397, AC005089, AC004859, AL049761, AD000812, AC004896, AL080317, AP000359, AC005011, AC007536, Z69837, AC010849, AL132987, AC006441, AC020663, AC005519, Z85986, AL049538, AC003101, AF196779, AC004049, AC006014, AC007993, AC004815, AC005971, AC004383, Z82195, AC004771, AL022329, AC005225, AC002094, AC005755, and AC008101.</p>
---------	-----	--------	---------	----------	---

HUSK176	332	914084	1 - 425	15 - 439
				AI805657, AW105425, AI684850, AA609319, AI445561, AW087416, AI569649, AI052449, AW005296, AI884876, AI683390, AI821774, AI857542, AI375889, AI498223, AI937172, AI052455, AI126807, AI276854, AW300679, AI148078, AI812099, AI079310, AI301571, AI832389, AA448176, AA634578, AA813907, AI159912, AA583197, AI081042, AI130045, AA960856, AA781740, AA078817, AI500194, AA037280, AA757372, AA577413, F17150, AA604543, AI694019, AA405506, AI025798, AI127345, AA890888, H78344, T50735, AI814129, AA002152, AA625807, AI352262, T23478, AI280471, AI240763, AW004806, H26970, H14637, AA933720, F03913, AA203660, AW327749, AA715028, M78972, M78956, AW074558, F36821, AA428254, AA427399, AI475077, F29918, T98945, R11215, R46235, R85210, F29924, AA878694, AA865297, AA910724, AA132925, R11273, AI147818, AA844613, AA564700, AA653527, AI214070, AW151132, AL042686, AA580663, AI432644, AI623302, AI431307, AI866786, AI431316, AI431238, AI955221, AI358271, AI440260, AI537677, AI494201, AI804505, AI500659, AI815239, AW058275, AI866465, AI815232, AI866691, AI801325, AI500523, AI538850, AI887775, AI582932, AI872423, AI590043, AI284517, AI923989, AI500706, AI491776, AI445237, AI926593, AI289791, AW151138, AI421662, AI889189, AI521560, AI500662, AW151974, AI285417, AW172723, AI539800, AI284509, AI582912, AI538885, AI889168, AI440263, AI927233, AI866573, AI633493, AI434256, AI866469, AI434242, AI805769, AI888661, AI500714, AI284513, AI888118, AI285439, AI859991, AI436429, AI355779, AI889147, AI623736, AI581033, AI371228, AI491710, AI440252, AI440238, AI567971, AI860003, AI610557, AI242736, AI539260, AI828574, AI887499, AI539781, AI889191, AI539707, AI702065, AI654286, AI885949, AI285419, AI559957, AW089557, AI696583, AI521571, AI469775, AI866581, AI567953, AI815150, AW074057, AI446495, AI867068, AI952433, AI225248, AI798359, AI698352, AI282249, AI890907, AI371229, AW084105, AI687944, AW194509, AI431315, AW151979, AI868931, AI924051, AI539771, AI872315, AI866458, AI687588, AL047422, AI540354, AI371251, AI889157, AI866510, AA829775, AI620864, AW151136, AI866461, AL047398, AI923046, AL042655, AW406745, AL039390, AI432666, AI561170, AI567954, AI469764, AI690946, AI648567, R20540, AI433157, AI932620, AA853473, AI249936, AI049859, AW029457, AL048403, AI362495, AL047611, AI799313, AI309306, AI801286, AI554821, AL045166, AI888022, AI433011, AI521596, AW129310, AI371243, AW151970, AL046681, AL039287, D50913, AL031732, AC003042, AR050959, AL133084, AL133070, AL049423, U30290, AL133047, AC002540, AL133053, AL133607, AL133049, AC004213, Z93784, AF124728, AF032666, L30117, E13998, X97332, A77033, A77035, AL133655, AL133076, AF081571, AL033523, AC007012, AR038854, AF090940, AF090432, AL034417, AC007114, Z82206, AC005968, AC007172, A41579, X83544, E12580, AL133015, AL133608, AL133088, AL122101, AR034821, I89947, AL133665, AI8777, X82434, AL133619, I48978, Z82250, AC006221, AL034374, AC002416, AC009113, AF206503, AF095901, AC005886, X98066, AF044323, AF139986, AF162270, A08910, AL080124, A08909, AL122049, A65340, D83989, I42402, X63162, A1005870, AL117435, AC002538, AL031656, AL022723, A08908, S59519, AL122118, S77771, AC002564, AC006203, AF000167, AF003737, AF003738, AR015970, D38178, U69730, U95739, AC005353, AL096776, AC005250,

					AP000020, AP000161, AC005091, AL122021, AL022147, AC005048, AC004686, Z94277, AL022722, AC004066, AC006501, AC008067, AC004057, AC018767, AC006222, AC008014, AC007056, AC006112, AC005291, AC007298, U66059, AF002985, A65341, AP000458, Z98744, Z49216, AL117587, AL050138, AF008439, AC005815, AF053356, AE000664, A86558, AF215669, AC007748, Y14314, AL110228, X66366, AL133051, X83508, AL133623, AC006373, AL034376, S70057, AL110280, AP000208, AP000130, AP000247, Z83840, AL049557, AL031274, AL020994, AC004690, AL137530, AL137574, AF172400, M64936, Z49258, U89335, AL080245, AC004989, AC006336, L05367, AC002301, AC003977, AL034400, AC007877, AL136520, AC002531, AL031054, AC007392, A21103, X00474, AL137478, A76337, Z13966, AF183393, AF026008, X93328, I48979, AF116573, Z98036, AL049742, AC006944, AP000697, AC005411, AC004554, AF013214, D44497, E03168, and X70514.
HUSXK92	333	848959	1 - 474	15 - 488	AA081241, AW014362, AW262136, R82455, AW058401, A1422211, A1394480, AA983672, A1342274, W35127, AA181256, A1246708, AA411749, A1291504, A1417724, A1304761, A1952055, A1653254, W02704, A1348150, A1242604, A1284964, W20374, A1318380, A1699314, A1933232, AA676646, A1249266, AA063542, A1167529, A1648485, AA908300, C00986, W25058, AA384416, A1553597, AA410298, N70938, AA410480, and AA411748.
HUSXO30	334	973266	1 - 540	15 - 554	AA421248, AA706819, A1831655, AW001127, A1659943, A1936705, AA482102, A1601264, AA609988, AW157288, A1249746, A1673452, A100758, A1168496, A1334090, AA291557, A1300503, AA838311, A1032723, A1446090, N92238, AA548418, AA458972, A1273811, A1243467, A1608877, A1720973, A1744302, H99292, A1762325, A1445510, A1248324, A1354517, N33480, AA999919, A1080097, AA837993, N26709, A1299967, AA459641, AW406362, AA523728, A1076417, W86074, A1222030, AA699459, A1052485, A1149225, A1018107, A1091825, AA961603, AA464265, A1983959, A1356935, AW263743, A1762657, D53076, AW070434, AA304075, R54623, AA484009, AA953293, AA806590, AA835477, A1309964, AA481610, AA464376, A1086019, W42436, AA641534, AA533309, AA303091, A1523186, A1969104, AA550930, AA506454, AA479918, AW182773, T57831, AA459188, A1077915, H78028, T40539, R81765, AA065082, A1248883, D30930, H67864, W42839, AW238626, A1278707, AA373915, A1749543, AA502726, AW068093, AA508653, AA480137, AW015934, AW338581, H67188, A1673126, AA514511, H46429, H78027, AA878251, AA952963, AA352423, AA428506, N55859, A1381737, T63660, A1419779, AA496458, H52178, AA658417, AA303090, A1286174, H25782, N43871, AW407913, AA758370, AA494208, AA552063, A1991234, AA961889, AA301776, A1609606, AA506757, AA370233, AW132067, A1654782, AA459415, W42551, H25788, A1434926, AA364016, A1866432, A1078057, A1825290, D81186, AA314590, AA291876, A1720956, AW103581, AA532661, AA353183, AA853831, AA481544, AA595944, A1468311, AA897049, AA151954, R71996, R54886, AA327953, AA371679, T82388, AA860424, A1128633, H52592, AA100822, AW170175, W42749, AA285031, AA331050, H19087, AA938023, H28635, N34590, AA373337, N25824, A1922550, A1887381, AL121286, T30322, AW161202, AA916133, AA825548, AA908294, A1690738, A1870192, AW025279, A1565172, AA464646, AW162194, A1559752.



20200740516007

					AI114703, AW020419, AI500714, AI554343, AL079963, AI537076, AI285439, AW104056, AA975952, AI433611, AW020693, AI494201, AI539800, AI801325, AW163834, AA828415, AW366372, AW082532, AI805688, AI491710, AI289791, AI524652, AI382670, AI357049, AW129117, AW163554, AW071349, AI812003, AW162189, AI933992, A761557, AA808175, AI656270, AI310575, AI633061, AL046466, AI340533, AA279795, AI345396, AI445620, AL039858, F27788, AI333104, N99092, AI537863, AI309306, AL037454, AI813986, AI499570, AI27565, AW022636, AW021717, AW089844, AI539723, AA806719, AI973236, R65859, AI590043, AI690813, AW262983, AI586931, AF031147, Z82022, X80340, AL133665, AF017437, AL117416, U96683, E01314, AI8777, AI242859, I32738, AL133049, AF000301, I89947, E04233, A08913, AR038854, L19437, Y11587, AL110296, AL050116, A08912, AL137556, I48978, A93350, AJ005690, X96540, U92068, AF132676, AF061836, AF106697, A45787, AF057300, AF057299, S76508, AL137530, AL133619, A65341, A77033, A77035, E12806, AF026124, A23630, AF032666, Z97214, A08910, AL133067, A08909, A08907, I89931, A65340, A08908, AF113676, AF106657, S77771, AF091084, I49625, AL122100, I09499, AL080126, AJ006417, AF115410, AL137271, M86826, AF047716, E06743, M27260, AF061795, AF151685, X72889, AI5345, A08911, I89934, AL133558, X63410, AL137276, AL080154, AF183393, AL117460, AL117578, X57961, AF106862, AF000145, AF008439, A58524, A58523, I79595, AF002985, AL080086, AF119337, AF110329, AL110221, AL080148, AF081197, AF081195, AF114170, AF036941, AL122106, AF195092, AL096751, AF079765, S54890, AJ001838, X06146, Y11254, AF038847, AL133062, AL050155, AF030513, AF002672, AR034821, I00734, X66862, Y10823, AL117463, A08916, AF159615, AJ238278, AF090903, AL023657, U00686, I66342, AF040751, E00617, E00717, E00778, AJ003118, AL137660, AL137495, AL122093, X67813, AL137641, U35846, AJ012755, X83508, I68732, D83032, AL049324, A07647, A12297, AF026816, AF012536, AF118094, S36676, AF215669, AF107847, E01812, AF111849, AL117649, Z37987, AL133075, AL110158, X57084, AF125948, AF044323, AF061981, E02221, AL133112, AL137479, AL137537, I89944, AR020905, X66871, AL049460, A76337, I33392, AR029490, AL137640, AF153205, AF017790, AL133016, U37312, AF177401, AJ010277, AF004162, AL050138, AL133010, DI6301, M92439, AF055917, Y08769, AF111112, AF162270, AF113019, E03348, X87582, E05822, E03349, L13297, AF069506, X84990, AL122098, U57352, A90832, X72387, U80742, S79832, AL137548, L04504, AF022363, U89295, A18788, E12747, L04849, AL080159, AL137560, AF115392, AF017152, AL137281, U87620, AL137256, AL049452, AF185576, AL122118, AL117435, AB008792, AF061573, U77594, AB008791, AL110280, AR053103, I29004, AL133560, AB019565, AF067790, AR013797, AF028823, AF126247, AF175903, AF090943, AF097996, AF067728, AL137478, Y10936, AL049430, AF051325, AL137258, E02349, X63162, AF176651, U55017, X67688, AL080162, AL117440, U78525, Y07905, U42766, E01614, E13364, and U51587.
HUSYA63	335	928021	1 - 428	15 - 442	AA909334, AI056548, AA723669, AA156125, AA303333, AW409843, AW263540, AA156120, AA157141, and AA151564.
HUSYB16	336	868843	1 - 423	15 - 437	AC008975.

HUSYD15	337	699195	1 - 401	15 - 415	AI264637, AW440517, AA045417, AA308065, AA127626, T96809, W39584, F22528, AA304050, AA082078, AI333981, AI087224, AI289816, N32838, AI042559, AA513003, AA716353, AA121528, AA302332, AI208270, W44935, AI805206, AA112905, AI024761, AA677140, AI087291, AA448463, AA101212, AA704993, AI097092, AA704961, T97914, T97458, AA203274, R09226, AA045311, AW451722, AI685445, N42284, AA062780, T97730, AI094396, AA932240, AA973273, AA062735, AA331180, T96692, AA431571, AI309788, AI248836, AI911056, and AI086794.
HUSYO46	338	868827	1 - 433	15 - 447	T96489, AI820673, T96482, AI732251, T96405, and T96398.
HUSYP67	339	575787	1 - 105	15 - 119	AW008247, F03615, H14592, H16768, AC007425, AP000519, AB023056, AF055066, AL049869, AC004172, AC004192, AB023058, AP000521, AL022723, and AC003001.
HUSZV72	340	851170	1 - 398	15 - 412	N79736, AW149788, AA187127, and AB011095.
HUSZH03	341	922852	1 - 785	15 - 799	R36618, R82444, R26337, and AC006443.
HUSYX03	342	922840	1 - 398	15 - 412	HI4379, R44811, H08865, F03387, and AL135783.
HUSYO86	343	784691	1 - 443	15 - 457	AI373275, T89685, AI718639, AA041276, R89363, AI192953, W80851, AW262845, AI861933, AI913971, AI810948, AI524062, AI970561, W23739, W35338, AA652221, H39865, AI871006, AI122618, AA197299, C75047, AA004293, AA594157, AP000356, AL031005, AC003684, AC006160, AL022721, AL109963, AL031003, AL035684, AC004678, U91323, AL022476, AC006312, Z78022, AC004983, AF001552, AL034548, AF135025, AL049569, AF047825, AF196969, AC007546, AC005412, AL050307, AC005280, AL050318, AL109758, AB023049, AC003043, AC006965, AC005914, AC005529, AC005746, AC003950, Z93930, AC003104, AL035086, Z92542, AL035413, AC005324, U91318, Z95331, AL133448, AL035455, AC007842, AC002350, AF067844, AC003003, AC004821, Z98044, Z85987, AC005482, AP000563, and AC007021.
HUSYN33	344	651293	1 - 450	15 - 464	R42079, AI215849, AI860946, AI024831, AI095766, AI523478, AW168267, AI554112, AI333039, AW263148, AI199955, AI708228, AA909572, AW264918, AI568327, AA936422, AA251096, AA010338, AA133384, AA731460, AA557232, AA058742, AA085448, AI097188, N52329, AA608709, AA031496, AA018410, AI123990, AI124972, AA416589, AI961604, N64588, R68411, R52688, AA593634, AI289991, AI284433, R51734, AA442321, AA975850, AA861668, R23934, AI735426, AI523604, AW317004, AA227638, AI263691, AA480016, AI699675, AW377524, AI499248, AI273115, AA889622, C01392, AA621372, R68066, AA377337, Z40503, W19549, F01878, AI203798, F01877, H88553, N24559, AA214352, AA085784, AW006661, AA814174, Z19952, AI955892, AL048323, AL048340, AL043152, AW157096, AI041393, AI954721, AI074008, AA830839, AI085582, AL046944, AI682932, AI266652, AW167328, X68497, AJ001202, AF129131, AL137478, AJ003118, and AF131821.
HUSYN11	345	943237	1 - 348	15 - 362	AI570590, AW391281, AI524309, AW103371, AI868831, AI687375, AI567351, AI538716, AI680113, AI281779, AI433976, AI568870, AI673256, AI366549, AI753683, AL119791, AW162071, AI866002, AW074869, AI497733, AI682841, AI952114, AW169653, AW071349, AL036759, AL036146, AI340582, AI521012, AI349614, AI859733, AI343112, AI349645, AI349598, AW274192, AW074993, AI800453, AI800433, AW238730, AI702406, AI498579, AI349004, AI349772, AI312152, AI349937,

AL040243, AW068845, AI349933, AI590128, AI866608, AW301409, AL135661, AL036980, AL119049, AI500553, AI433157, AI863014, AW148320, AI500077, AL045903, AI249257, AL036396, AL038605, AI275175, AI432813, AI492540, AW071417, AI857296, AW268253, AI702433, AI345735, AI564719, AW089572, AL047763, AI309401, AI620284, AW302965, AA613907, AW195957, AI583316, AL045500, AL036274, AI349256, AI687362, AI636456, AA640779, AI873731, AW301300, AI636445, AI568854, AI758437, AI678302, AI969567, AI633419, AI679724, AL036802, AW269097, AI690835, AI282655, AI343059, AL046849, AI597918, AI250293, AI686926, AI281773, AW303152, AI608667, AA938383, AI906328, AI439087, AI813914, AI285735, AI340519, AL121270, AI348897, AI224992, AW235035, AI889203, AL038779, AI499463, AI567632, AI866687, AI679764, AI823670, AI671679, AI064830, AL047042, AW117882, AI440426, AI334902, AL119748, AL121365, AI307466, AI969601, AI687728, AL038778, AL120736, AI349226, AI493248, AI597750, AI570384, AI921379, AI699857, AA572758, AI475134, AI690751, AW268072, AI625079, AI613017, AL120854, AI540832, AI635461, AI631107, AI800411, AI818683, AI499393, AI610645, AI745485, AI445432, AI815383, AW302992, AL048871, AI436456, AA508692, AI207510, AI307558, AI920968, AI909662, AI434281, AI687376, AI934036, AI469532, AI889839, AI610307, AI866780, AI580190, AW149287, AI345111, AL040169, AI697137, AW071412, AI564247, AI874109, AI475371, AI439745, AI609592, AI269696, AW166645, AW167776, AI500659, AI811845, AI283941, AI919058, AI628205, AI631057, AI446606, AI687415, AI539771, AA493647, AI818206, AI580984, AI683684, AW080838, AI799199, AW075351, AI312542, AI799305, AI754897, AI569616, AI909666, AI624668, AW118557, AI811863, AL036260, AI907070, AA528491, AL047041, AA585422, AI149592, AI802542, AI620868, AW026882, AW085799, AL038604, AW005858, AI922665, AW132121, AW087445, AI888953, AL043326, AI282281, AL042753, AI696846, AI696398, AW193134, AI811509, AI432229, AI318280, AW183130, AI273142, AI818977, AI598061, AL121014, AW002342, AL036247, AI499131, AI609580, AI922901, AI569583, AL036240, AI431424, AB007963, I48979, AF118070, AL049452, AL049938, A08916, I89947, AF113676, AF125949, S78214, AF090934, AL110221, AF078844, AL133606, A93016, AF118064, AL080060, Y11587, AF113694, AF104032, AL050146, AL137527, AF113013, AI242859, L31396, L31397, AL133640, AR059958, AL133016, S68736, AF090901, AF090903, AF090943, AF113691, AL117457, AL050393, AF113690, AL110196, AL117460, A08913, AF090900, AL050149, I89931, AF090896, AL122050, AL050108, X84990, AL049466, AF106862, AL133075, AL049314, AB019565, AF113677, AF017152, AL133557, AF113019, I48978, I49625, AL122123, AF158248, AL080124, AL133093, AL050277, AL080137, AL122093, E03348, AL096744, AL137557, AF113699, AJ238278, AL133565, AL122121, AL133080, AL117585, E07361, AL117583, AF125948, AL137283, A08910, AL110225, X63574, AL117394, AL050138, AL049464, AL137459, AF146568, AF091084, Y16645, AF111851, X82434, AJ000937, AF017437, Y11254, AL137550, AL133560, AR011880, A65341, E02349, AF177401, AF079765, A08912, E07108, U91329, A08909, AL049430, X93495, AL049300, U72620, U00763, A58524, A58523, AL137463, AF118094, U80742, X70685, AL049382, AF183393, AL117435, X96540, AL137648, AL122098, AL133113,		
---	--	--

					X65873, AL137338, AL080127, X72889, Z82022, A12297, I03321, I09360, AF097996, A77033, A77035, AL137271, U35846, AL050024, I33392, E15569, AL122110, I26207, A03736, AL080159, AF087943, AF067728, E08263, E08264, X98834, U67958, AL122049, I42402, AL137521, AL133077, AL133014, AF061943, AL137556, AL049283, AL133072, AL121603, AL110197, AR038969, AC004690, S61953, AC006840, AF153205, AL133568, AF111112, AC007298, AR000496, U39656, Y14314, AC006336, AJ012755, AF119337, AC007390, AL096776, AL137560, A90832, AF026816, AC005488, AF026124, AL110280, AF003737, AF162270, AL133098, A93350, U58996, AC004987, AC006371, AF000145, AL133104, AL137526, E08631, AF185576, U96683, A45787, AL117440, AL137476, A08911, AL080074, L30117, M30514, Y09972, X83508, AR038854, I17767, I00734, E02221, AF057300, AF057299, AL133067, AL050172, AF079763, AC004200, A07647, E00617, E00717, E00778, X87582, Z72491, U66059, AL035587, AF051325, AF132676, AF061836, AF095901, AF109906, X92070, U68387, AJ006417, AC006222, E04233, E05822, AL137523, AC002467, AF091512, AF111849, I41145, AF106827, and X53587.
HUSYM49	346	723015	1 - 257	15 - 271	AA525157, H54430, AI370548, N68565, T91791, AC007263, AC007676, AC004616, AC005089, AC005225, AC004757, AC002550, AC006966, AC004531, AC006441, AC005412, AC002485, AC002044, AL020993, AC002073, AC005792, AC005338, AC005015, Z84487, AL079342, L44140, AC006121, AC004890, AC007934, AC003035, AC004587, AC002350, AC007279, AL121653, Z93023, AC005049, X55448, AF111168, AC006120, AC004865, AL049712, U91323, Z94161, AL109963, AB015355, AC003037, AC008115, AP000512, AP000553, AC005666, Z83822, AC006511, AC007226, AL049795, AL080243, and AP000691.
HUSYM37	347	464221	1 - 324	15 - 338	AI251034, AI251203, AI251284, AI250552, AI251944, AI349748, AI284543, AI354397, AW268231, AL046746, AW303098, AI345629, AI307588, AA469327, AI345394, AI250577, AA837715, AI223626, AI254770, AI225179, AI271217, N69399, F25761, AI267823, F33126, T52745, AI246061, F23326, AI754105, AI755214, AI249853, AC004560, AC007358, AC005479, AC004030, AL031280, AC005777, AC006126, U91323, AC007792, AC005332, AC005900, AC005632, AC002389, AL035461, AC006930, AC002044, AC004125, AF207550, AF111169, AC005765, L78810, Z98044, AC005264, AC005620, AC005952, AC003070, Z97832, AC003101, AC002492, AC002350, AL121603, AC002290, AC003007, AC002303, AC002045, AC005778, AC005305, AL049749, AC004841, AL008583, AC004382, AC000039, AC005291, AC004821, AC004832, AC008372, AL022315, AC004887, AL035683, AL031595, AC002352, AC002316, AC000070, AC004517, AL035684, AL049538, AL034423, AC008040, AC005562, AL023879, AC005335, AC004112, AC005775, AC003684, AL035659, AL049712, AC005578, AC004525, AC005197, AC006125, AC007193, AC005880, AL136295, AL022323, Z98752, AC004703, AC003982, AC002375, AL023807, AC000026, AC007934, AC006952, AF196779, AC007191, AC004967, AC005177, AC007878, AC004938, AC005480, AC005786, AC003957, AC005157, AL031283, AL022316, AL109939, AC002477, AC003664, AP000142, AC005015, L47234, AF002223, AC010205, AC006079, AC005913, AC004671, AC005944, AL050317, AC004032, AL008735, AC003026, AC005210, AL031289, AC004685, AF001548.

					AC002544, AC000063, AL022314, AL021528, U91326, AL049576, Z92547, AL021391, AC004851, AC004885, AF123462, AL034429, U91318, Z83851, AC002070, Z93017, Z97987, AL021155, AL049871, AC005790, AC006312, AL034430, AL035588, AC005779, AL049759, AC005932, AF047825, AC004104, AC007386, AC005839, Z86090, U80017, AC016831, AC004216, AC005339, AC004584, AC002072, AC004067, AC005668, AC006071, AL020993, AL117694, AF107885, AC005821, AC002554, AL035685, AL109628, Z81369, AC004024, AL031585, AC005520, AC005488, Z85994, U62293, AL050307, AC004983, AC005764, AC005031, AC005355, AC000353, AL035089, AC002467, AC007073, AL022324, AL022476, AC007226, Z85996, AL049611, Z97056, AF001549, AC006511, AC005859, AC006013, AC004973, AJ003147, AL079295, Z83826, AC002511, AC007425, AC003956, AC006531, AC007052, AL031284, AC005594, AD000092, Z83840, Z86064, AC006023, D87675, AC005822, AC003010, AC005011, AC005722, AC002059, AC008041, AC004590, AC006014, and AC002378.
HUSY113	348	657288	1 - 802	15 - 816	N25018, H98802, AW294590, AA553572, AW236050, AI268116, AA777178, AA417117, AW236049, AA463827, and AA445940.
HUSYG24	349	677258	1 - 662	15 - 676	N63953, N40973, H23529, T78668, F08173, Z44774, F08664, and F08221.
HUSYF74	350	554723	1 - 465	15 - 479	AA688133, W15472, AA603105, AA773036, H05310, AI242746, AI279200, AI573250, T03740, R38903, AI342482, AI818093, AI221800, AA577570, AI285452, AA872045, T03417, W04515, AI758880, AL041772, AI539153, AW102785, AL039086, AW023590, AL038505, AW161579, AW071417, AI634224, AW238730, AL079963, AI358701, AL041150, AI635067, AI344785, AI567582, AI349645, AI284517, AI923989, AL119836, AL042538, AI345608, AI921379, AI446606, AI312428, AI812080, AI436429, AL110306, AI345471, AI612759, AI929108, AI872910, AW161156, AL120853, AI521012, AI627988, AW089179, AL121328, AL036403, AI874166, AA580663, AI670009, AW059828, AI635942, F27788, AI625589, AI567351, AL048656, AI348897, AW082113, AL119791, AI567993, AI815232, AW302965, AL038605, AA640779, AI340603, AA613907, AA572758, AI950892, AW068845, AI923370, AI612920, AI783504, AI499285, AI620284, AI859464, AL119863, AI445992, AI537244, AI538342, AI866770, AI433157, AL036631, AW188539, AI702073, AI538829, AL049085, AI340511, AW269097, AL036396, AI269580, AI499986, AI174819, AI500061, AI689420, AI251830, AI288285, AI633125, AI345347, AI310575, AI698391, AI251221, AI801325, AI500523, AA493647, AI915291, AI582932, AL121365, AI500706, AI340533, AW026882, AI521560, AI889189, AI500662, AI343059, AI345745, AI284509, AI312542, AL041220, AI866573, AI830029, AI343091, AA833760, AW022682, AI866798, AI888661, AW403717, AI340627, AI308032, AW074993, AI349933, AI349614, AI431424, AW022699, AI343112, AW193134, AW302992, AW105601, AI919345, AL036980, AI686906, AL047422, AW268253, AI862144, AW301300, AI567612, AI349598, AA225339, AI440263, AI890806, AL036664, AW075207, AI349256, AI554821, AI589267, AI312152, AI636719, AI955906, AI345735, AW268072, AW075084, AW191003, AI950664, AI468872, AW152469, AI698427, AI349937, AI499920, AI364788, AW089572, AI334884, AI335426, AI348777, AI036274, AI307708, N71180, AI312325, AI500659, AA641818, AL038445, AI890507, AA579232,

					AI933589, AI313320, AI089748, AI612885, AL134999, AA494167, AI432040, AI863321, AI609911, AI445990, AW020693, AI307520, AL036638, AW074869, AI445237, AW151138, AW243886, AI312146, AI312339, AI345258, AI889168, AL040241, AA508692, AI745713, AI434256, AW151136, AW150578, W33163, AW195969, AW020419, AW163834, AI8666608, AL048644, AW050522, AI570807, AW081255, AI471361, AL045500, AI267502, AL036802, AI457113, AI343037, AI862142, AI174394, N71199, AI866002, AL038959, AL036214, AI784230, AI679724, AI446373, AI340582, AW073898, N33175, AL134259, AI539687, AI345737, AI537677, AB002372, A08916, X62580, A08910, I89947, A08909, X92070, A08913, I48978, AL122050, X70685, AL137459, AF113691, AL117460, AL137658, I89931, I49625, AF113690, I48979, AF061943, AR011880, AL080060, AL110196, AR038854, AF177401, AL049430, S78214, AL133568, Y11254, AL117583, AF090901, AL133113, AL133565, AF026816, E03348, AF017437, A65341, AR059958, AL110221, AF125948, AL133077, AF158248, AL050108, AF113013, X93495, AL133606, A58524, A58523, AF078844, AF118094, AF097996, AL133640, I42402, AF111851, E08631, AF113676, AL137527, AL133093, U00763, AL137557, I33392, E02349, AL137648, AL080158, A93350, AF090886, AF090900, AF090903, AF185576, AL117432, AB019565, AF090934, AF113677, AF118070, AL049452, Y10080, AL133557, AL122093, X63574, AL122121, X96540, X98834, AL050277, AF113694, AF113019, AF113689, Y16645, AL049466, Y11587, AJ000937, Z72491, Z82022, AF183393, U91329, AL122123, AF091084, AR013797, AL133104, AR038969, AF118064, AL050024, AJ238278, E07108, AF125949, AJ003118, AL117435, U42766, E01614, E13364, X72889, AL133560, A08912, A93016, AL137283, X87582, AL049283, AL133080, AF113699, AL137550, AL137538, AL117457, AL050116, AL050146, AF090896, AF079765, X82434, AF067728, A77033, A77035, AF087943, AL049382, AL137271, A07647, AL096744, AL117394, AL050393, A03736, AF057300, AJ012755, AF057299, AF104032, AL122110, AF119337, AF003737, I26207, A18777, AF017152, S68736, U80742, S79832, AF022363, AL049464, AL049314, AF079763, AJ242859, AL117585, AL122118, AL133072, X65873, AJ006417, U72620, AL080086, X72387, AF090943, E06743, AF132676, AF061836, AL050149, L31396, AL110225, L31397, AL122049, U67958, I03321, AF118090, AF153205, D16301, AL133075, E15569, AL080137, AL050155, AL137521, AL110280, AL080124, AL137429, E04233, AL137560, X84990, AL080127, AL137705, U35846, AL137300, AF106862, AL133098, AL122098, A08908, AL133016, AF146568, I00734, AL137463, X81464, S61953, AF176651, AF031147, AF026124, E00617, E00717, E00778, AL133014, AL137479, AL137476, I41145, AF162270, Y10655, AL137526, AL049300, I09360, AL049938, AR029490, A90832, AL122111, AF111849, U68233, I92592, AL050092, AL050138, AL137480, AF061573, E07361, AR020905, AL133067, AF065135, AL080159, AF051325, M30514, U58996, U68387, AL096751, AF028823, AF126247, L19437, AL110197, I17767, AF142672, Y09972, I09499, U88966, AF111112, and AL080074.
HUSYA27	351	934423	1 - 624	15 - 638	AI810054, AI004966, AI953417, AA204830, AA700282, AA101580, AA127230, T77885, AI215410, AI049842, AI383026, AW193598, AW195356, and AL137534.
HUSXW61	352	741856	1 - 623	15 - 637	AW439843, AW172765, H96902, H95980, and T15387.

HUSXM28	353	703326	1 - 418	15 - 432	T84537, R92899, and AC007999.
HUSX171	354	760415	1 - 155	15 - 169	N77614, and AL034417.
HUSXH57	355	859907	1 - 656	15 - 670	AA894530, AI336165, AI379833, AI870419, AI339336, AW173013, AI660120, AI697018, AI418065, AA946777, AI372055, AI961630, AI281293, AI018252, AA918864, AI804228, AI280166, AI191797, AA115687, AI476152, AA938579, W58372, AI124079, AI123356, AA460235, AA961355, AI201697, AA806449, AA181956, AA633371, N54918, AI284979, AA975500, AA948109, AI262806, T86966, AA864307, AA126483, AA670154, AA187579, AI285113, AA928294, AI282777, AA856633, AA554905, W04639, H01916, AA952898, AA358260, H04478, AA939258, AI498851, AA283086, AA070685, AA503091, AI473325, T56012, AA553782, AI718566, N21275, N78134, H78232, T31842, AA651925, AI583297, W31735, AA383311, AA383310, R70784, D20722, AA296522, AI381360, N40501, AA282901, W58050, C14037, AA932915, AW188939, AA770225, AI284357, AA969062, AA918735, AI658500, and AB034206.
HUSKA86	356	784887	1 - 411	15 - 425	N72485, H71940, AF072873, and AB012911.
HUSKA65	357	868860	1 - 490	15 - 504	AA551165.
HUSJW03	358	923035	1 - 700	15 - 714	AI057455, AA699325, H75865, H75866, AP000247, AC004216, AC005071, AL035415, AC005790, and AL121603.
HUSJN66	359	886987	1 - 766	15 - 780	AA426047.
HUSIT75	360	679416	1 - 453	15 - 467	AA040679, AI669256, AI763048, AI902903, AW245958, AW237088, AF039235, U25751, AI312767, D80899, AI571378, AA362530, AW296090, AI354976, AI093187, AI680102, AA625161, C01220, AW119180, AI140327, AA934553, AI201269, AA559846, AA040597, AI991982, AA359868, AI203368, AW087508, AA497000, AW268813, AW152391, AI243335, AI185996, D80898, AI887841, AA489585, F31194, and AI700879.
HUSIS59	361	739327	1 - 217	15 - 231	
HUSIS54	362	730734	1 - 188	15 - 202	AI149061, AW130563, D87675, AP000089, AP000142, AC005182, AC005089, Z83822, AC004832, and AC007229.
HUSIS08	363	959536	1 - 421	15 - 435	AA663486, AA635417, and AA196796.
HUSIR04	364	709228	1 - 422	15 - 436	H77990, H90576, and AI267628.
HUSIN12	365	970759	1 - 481	15 - 495	AI689603, AL039203, and AW104130.
HUSIE95	366	967176	1 - 482	15 - 496	AI186511, AI905032, AW248242, W30830, AI908041, AA287850, and AF212940.
HUSIE18	367	666523	1 - 404	15 - 418	H47045.
HUSIE08	368	908574	1 - 964	15 - 978	AA736450, AW205616, AI492501, AI275895, AI806754, AA426438, AA437170, AI825464, AI984990, D81011, D81007, AI954689, AI962153, and H26441.
HUSHL86	369	960355	1 - 1399	15 - 1413	AA703915, W19899, AI926481, C06034, AL120774, AI926508, AI889895, AI125240, AA508101, AI140921, AI620543, AI889904, AA557537, AI198792, AI222307, AA744258, AI400233, AI630644, AI266510, AI832176, AI699153, N91246, AA312954, AA258943, AI718296, AA053015, and AF151805.

HUSHE34	370	703409	1 - 549	15 - 563	AA253268.
HUSHB71	371	766060	1 - 329	15 - 343	H79732, and H79731.
HUSHB60	372	746560	1 - 462	15 - 476	AA132883, AA253947, and AB033047.
HUSGW06	373	935574	1 - 625	15 - 639	AI375142, AI041354, AI739576, AW449593, HI2704, AI684781, HI2705, AW151325, AL137763, and AL031431.
HUSGV84	374	813557	1 - 433	15 - 447	N22167, AI469417, AA186334, AI084312, AL048280, AL048279, AA845906, AA542871, AA164384, AI927103, N92214, AA482803, AA482818, AI206093, AA884457, AA167321, AA548958, AI925703, AI537994, AI973197, AA985451, AW072297, AW054739, N64610, AW151542, AW105365, N93598, AI866140, AI399865, AA525978, AA516263, AA587848, AA663007, AI857882, AA594987, AA173554, AA156124, AA830434, AA125898, AA834145, AA620469, AA843345, AA844552, AI023983, AI041347, AA045631, AA854158, AA582010, AA618385, AI872043, AA101123, AA187231, AI924029, AI126870, AA595697, AI338472, AA524686, AA469343, AA865364, AA652110, AA657769, F24916, AI720143, AI698664, AA862692, AA469127, AW263715, T34217, AA501497, AI687698, AI253667, AI745562, AI669479, AA748114, AW440479, AI033755, AI598103, AI282490, AI567685, AA527738, AW304954, AA046680, AI630921, AI687740, AA927656, AW169855, AA195478, AI301902, AW243866, AI253769, AA541719, T51486, AI984248, AI360533, AA224320, AA176133, AA121062, AA635576, AA635568, AW151767, AI709168, AA112117, AI052129, AA227540, AA565009, AW275898, AI275685, AA662712, AA527766, AI744375, AA514849, AW316620, AA971114, AA227680, AA931889, AI199685, AI306714, F33796, AI954557, AI609269, H73139, AA480503, T93062, AA469196, AA834505, T92777, AA167699, AI523421, AA853136, AI472161, AI560670, AI567024, AI358189, AI244950, AI222622, AA593985, AI922122, AI253759, AW071035, AA223905, AI890437, HI9976, AA224846, AI460153, N64480, AI890697, AI444954, AI972952, Z28473, AI889384, AI708079, H06213, T41240, AI671926, AI041981, DI2312, D51107, AI038910, H30023, AI687614, AI869809, AA716547, AA530894, AI749102, AI833109, AI749897, AA661582, F19628, R71226, AI720476, AW440666, AI719996, AI091016, AI961304, AI954378, AI274494, F28330, AW078568, AI807358, AA508357, AA570600, AI493435, AA507860, AA425416, AW167098, AA687857, AA724960, AW167109, F27045, AW363237, AW363233, T93147, AI422648, AA167142, AI312755, D56789, AI380771, F25929, AI676180, AI312752, AA738139, C16070, AI783456, AA775238, D57103, AI932746, D57956, F01287, AA962120, F28415, F29404, AI269032, AI871878, AI434478, AI925288, F30481, AW198053, N75639, AA541559, AI718158, AI750135, F30485, F20314, AA487950, AI932857, AW374950, F35602, AI632607, D54610, AI864772, N55135, AA480577, AA247574, F00097, AI131951, and AJ225898.
HUSGU08	375	959540	1 - 429	15 - 443	AI745055.
HUSGT01	376	916620	1 - 509	15 - 523	AA610520, AW296489, AA179163, AI824978, AA827817, AA483268, AC004966, AL049761, AC004216, AL031591, AC002059, AC004626, AF109907, AC006014, AF064861, Z84466, U62317, AL031904, AC000026, AL022324, AP000099, AP000260, AC005775, AC003662, AC004973, and AC015853.



HUSGS35	377	707777	1 - 666	15 - 680	AI271425, AI223830, AA564436, AA251799, AI167485, N26127, AA143235, AI051406, AA975269, AI927242, AI565174, AI052812, AA405571, AA251929, AA610292, AA826503, AW371603, AW103340, AI289757, AI472362, and AI431964.
HUSGQ62	378	745727	1 - 492	15 - 506	R93146, AA704227, AI340024, and AC005021.
HUSGM24	379	425180	1 - 773	15 - 787	R81604, AI928098, and AI347338.
HUSG168	380	753059	1 - 352	15 - 366	AA767257, AI799645, AA906290, W57567, AA729117, AA171892, AA526416, AI732502, AI791301, AA205911, AI791458, T71257, AA658087, AA743698, AA551400, R36518, H47143, AA330897, AA446645, AI057588, AA225380, N91747, AI748842, H69661, AI339725, H73306, H52510, AA358852, AI475758, AA424008, AA167159, AA487281, AC003003, Z68277, AC002430, AC007384, AP000699, AC002420, AC007786, AC007536, Z99755, AC011504, AF111167, AC005071, AC006047, AP000509, AC002350, AC007066, AC005180, AC005821, AC002288, AC004644, AC006111, AL117344, AL023883, AL049697, AC006204, AL034377, AC007488, Z98304, AC004885, M89651, AL008725, AC004910, Z95152, AC005839, Z94277, AL022316, AL049742, AC020663, AC007371, AL022721, AC006071, AC005911, AC005323, AL137191, AL222316, AL049776, X90590, AC004638, AC007993, AL049871, AC004587, Z99754, AD000092, AL022726, AC005933, AC004750, AC002077, AC000387, AL022315, AC006449, Z82176, AC005702, and AC005204.
HUSGH09	381	625647	1 - 726	15 - 740	T98941, AA777690, AA704182, and AF057709.
HUSGF79	382	775309	1 - 424	15 - 438	AA262313, R67992, AA262917, AI420959, and AI695670.
HUSGF59	383	576784	1 - 302	15 - 316	F34498, AA491814, AL119691, AA661921, AA689526, AI284640, AA493708, AA527630, AA714453, AA502103, AA063173, AA580808, AA613591, AA613397, AA658362, AW303196, AW274349, AI754955, AA507547, AI281697, AA502860, AA316905, AI338350, AA491862, AW270382, AW028392, AA559290, AA618452, AA847409, AI282511, AW168618, R77905, AA525824, AA362349, AA848134, AW103758, AI687343, AA857486, AA346458, T94842, AI358571, AA806796, AI367975, AA514854, AA491650, AA469451, AA847516, AL046409, N27763, AA491284, AI499487, AW301350, R89294, AI084294, AA362698, AW265385, R98359, AA491831, AA364429, AA219225, AA812281, F17891, M78005, AI821271, AA641052, AA483256, AA531079, AI678316, AA828227, F17555, D52587, AA713891, AI798489, AI281881, D51681, AI537077, AA501600, AA630925, D83989, X75335, X55923, AF077058, U67827, U67829, M87925, AF015151, AF015156, M87919, AF010317, AL133246, S70707, L10641, AL031255, AC006292, AC006596, AC000353, AC004638, AL031650, AP000555, U62293, AC002564, L49046, S43650, AC009510, AL132800, Z49769, AD000684, AL133276, Z69666, AL031054, AL031542, AC005288, AL031281, AF001549, AL031668, Z70042, AC002045, AC000118, U49740, AB019437, AC004016, AL096765, Z97205, AC004931, AC005696, AC000379, AP000431, AL132712, AC005257, AB020859, AC005913, AC006213, U66059, AC007682, AC005603, AL110120, U38950, AC004134, AC008079, AL080242, AC007285, AL023882, AL031657, AC010382, AC007043, X14448, AP000159, AC004987, AC006128, AC006130, AL035668, Z98742, AC006057, AL023755, AL133494, AC005597, AC005837, AL022302, U85195, AC005839, AC006315, AC004675, AC005862, AL023800, AL049544, AC005740, AC006155, AL020993,

Variable	Mean	Standard Deviation	Minimum	Maximum
Age	34.2	10.5	22	55
Gender	0.5	0.5	0	1
Marital Status	0.7	0.5	0	1
Education	12.5	1.5	10	16
Income	3500	1500	1000	8000
Health	0.8	0.4	0	1
Stress	4.5	1.5	1	7
Depression	2.5	1.5	1	5
Life Satisfaction	5.5	1.5	1	9
Work Satisfaction	6.5	1.5	1	9
Family Satisfaction	7.5	1.5	1	9
Community Satisfaction	8.5	1.5	1	9
Overall Satisfaction	7.5	1.5	1	9

						AC007324, AC003103, AL078615, AC005026, AP000557, AC004041, AC007919, AF045448, AC008101, AE000658, AC004946, AC007319, Z74022, AC007666, AL031682, AC007263, AL009179, AC007064, AC004690, AL078581, U91321, AC004972, Z98051, AC006538, AF042090, AP000114, AP000046, AC002350, AC004650, AC006045, AF064861, AC009247, X69907, AF088219, U72787, AC009069, AP000459, and AP000090.
HUSGF10	384	964844	1 - 564	15 - 578		
HUSGE22	385	888829	1 - 446	15 - 460		
HUSGB36	386	572924	1 - 418	15 - 432		R33929, and AC004099.
HUSGB01	387	916804	1 - 423	15 - 437		T70191, AC004812, AC007263, AC005412, AC004686, AF111168, AC004253, Z84469, AC008044, AC006111, AC000353, AC006128, Z98941, AC004216, AC005057, AC005484, AC005911, AL031005, AC016027, AF067844, AC006312, AB023051, AL049843, AC016830, AP000512, AC006449, AF196779, AL035413, AC008040, AC005785, AC005667, AL009183, Z98884, AC004491, AC006948, AC004150, AC005037, AC004526, AC005004, Z95329, AC004019, AL133445, U62293, AC005792, Z99716, AC004972, AC007386, AL109623, Z84487, AC005736, AC005345, U63721, AC005081, AL031681, AC005776, AC007207, AL031283, AF053356, AC005086, AL022315, AC004752, AC005696, AF001549, AC004382, AL121655, AC005067, AL022336, AL031846, AL031602, AF045555, AC008372, Z85987, AC005585, AC020663, AL049569, AL031666, AC004685, AC004605, AC007773, AC006511, AC007114, AC006974, AL034549, AC005874, AF134471, AC002316, AP000689, Z84466, AC002565, AL109798, AC006057, AC003665, and AC006571.
HUSFH89	388	786970	1 - 500	15 - 514		H53955, and T95446.
HUSFF03	389	924616	1 - 577	15 - 591		AA769109, AA283754, AA883471, AB023137, AF033276, AF033275, and AF033274.
HUSFE05	390	932106	1 - 796	15 - 810		AA251680, and AA219333.
HUSDA09	391	461656	1 - 475	15 - 489		AA180076.
HUSAY21	392	920403	1 - 508	15 - 522		R66556, R66555, AA305729, AI903265, AA648467, and T03611.
HUSAO27	393	955287	1 - 320	15 - 334		
HUSAM87	394	529783	1 - 134	15 - 148		
HUSAM35	395	558191	1 - 245	15 - 259		M63005, M63543, M63480, and M63544.
HUSAM22	396	523674	1 - 100	15 - 114		
HUSAL04	397	927719	1 - 316	15 - 330		AC004547.
HUSAJ57	398	678932	1 - 339	15 - 353		
HUSAJ15	399	522056	1 - 243	15 - 257		R45895, T18597, R28735, R29445, AA585325, D54897, D60765, R28892, D60844, AA585098, AA170832, R29657, R29218, AA585101, R28895, D61185, Z32887, D59751, C15406, D53161, C16315, D61254, AI546999, D55233, D57491, Z32822, R28965, AA585439, AA283326, AI557262, Z28355, AI546875, C15069, R28967, AI557864, AI541356, D59436, D53472, C16294, AI557734, C16293, AI525316, C15120, D52835, AA585476, AA585155, D53447, AI526140, AI546971, AI541365, AI525500, AI557740, C15762, AI541383, AI525431, AI547250, AI547202, AI546945, AI541374,

					AI525306, C16292, AI541205, AI547039, AI526184, AI557763, AI526078, AI540903, AI541517, AI541013, C15737, C16300, AI526109, AI557731, AI557718, AI557727, AI526016, AI541307, C16305, AI541535, AI525556, AI541034, AI526194, D60730, Z33559, AI546891, AI557084, AI525320, AI557602, AI557787, C16296, R29179, Z30131, AI547006, AI546921, AA585356, AI557807, AI525339, AI557758, AI541346, AI546996, AI540967, AI526195, AI557809, AI541523, AI535639, D57186, D51433, AI526191, D54850, AI526113, AI557408, AI557533, D59458, C14208, AI557155, AI547196, R29262, AI546831, R29172, AI557852, C14391, AI557808, C16290, AI541527, R29177, AI547137, AI556967, T41289, AI540974, AI541027, AI525856, AI526024, AI526117, AI541321, AI526112, AI526158, AI557279, AA585453, AI546901, AI541422, AI526146, AA514191, AI526073, AI524904, AA585430, AI535660, AI541492, T19407, AI525286, AI541506, AI546829, AI557799, AI540920, AI557264, AI526180, AI525332, AI524890, C14322, AI525076, T41329, AI546828, AI541510, AI557039, AI541515, AI525168, AI541415, AI525656, AI546954, AI540882, C14210, AI557317, AI547189, AI536138, AI526205, AI541423, AA174170, AI540944, AA585434, AI557810, AI541345, C14723, AI557786, AI525114, AI557041, AI541390, AI526196, AI541017, AI5255040, AI557238, AI557082, AI524891, AI557785, AI557285, AI541514, AI546841, AI526169, AA585420, AI526125, AI541508, AI547138, AI046024, AI526187, AI526026, AI540928, Y09813, AR062871, A25909, AR038855, X82786, X76012, AJ244005, Y16359, AF082186, AR038762, Z32836, AR054723, X81969, D50010, D13509, AJ244004, A20702, AR062872, AR062873, A20700, D78345, A43189, A43188, X55486, AR017907, AJ244003, AJ244006, AJ243486, AR031358, AR031365, A98420, A98423, A98432, A98436, A98417, A98427, AR003381, AC005913, X82834, AR017826, A98767, A93963, A93964, I63120, and Z30183.
HULAG30	400	788577	1 - 461	15 - 475	R21151, AA053860, AA256918, AA054011, AA258280, AA026671, R72975, N48017, R69294, D79117, W47268, W78713, W56158, AA417682, H03608, AA379320, AA019166, AA026702, R34494, H84115, AA131247, AA044911, AA164369, AI301065, N36665, N73920, and AA131515.
HMELV25	401	678120	1 - 491	15 - 505	T97544, AA811921, AW237809, AA256699, and AL031591.
HMELV14	402	876087	1 - 385	15 - 399	AI026625, AA897168, AA846392, H30053, and AA989218.
HMELR45	403	717696	1 - 178	15 - 192	AL078634, AF031075, AF030876, U52112, AC004213, AL022723, AC006199, AL122021, Z68869, AC002425, and AC002352.
HMELM86	404	784702	1 - 235	15 - 249	
HMELM85	405	783536	1 - 211	15 - 225	C06367, and AL034374.
HMELM03	406	924168	1 - 302	15 - 316	H70012, H69999, AA188729, AI623442, AI078409, AA584765, AA729384, AW237905, AA302690, H73550, AA595770, AW304580, AA833896, AA833875, AA713705, AI298079, AI742168, AW265688, AA601728, AA582746, AI272052, AA536040, R91816, AA481887, AA654874, AA279649, T60666, N71729, AI369580, AI491755, AL040430, AA995373, AI282629, AW300749, AW194046, AA912287, AA663461, AA564642, AI024339, T17332, AI963679, AA455252, AA680253, N35306, AA525293, AI792439, AA689351, AW002831, AA525379, AI917132, Z30294, AA360245, AL045476, AA582374, AA346436, AA493808, AA715277, AA485343, H52397, AI583448, C14480,

Variable	Mean	Standard Deviation	Minimum	Maximum
Age	34.5	10.5	20	55
Gender	0.5	0.5	0	1
Marital Status	0.5	0.5	0	1
Education	12.5	1.5	10	15
Income	35000	15000	10000	60000
Health	0.5	0.5	0	1
Exercise	0.5	0.5	0	1
Stress	0.5	0.5	0	1
Sleep	0.5	0.5	0	1
Diet	0.5	0.5	0	1
Alcohol	0.5	0.5	0	1
Smoking	0.5	0.5	0	1
Family Size	2.5	1.5	1	5
Work Hours	40	10	30	50
Job Satisfaction	0.5	0.5	0	1
Life Satisfaction	0.5	0.5	0	1
Depression	0.5	0.5	0	1
Loneliness	0.5	0.5	0	1
Self-Esteem	0.5	0.5	0	1
Resilience	0.5	0.5	0	1
Optimism	0.5	0.5	0	1
Gratitude	0.5	0.5	0	1
Forgiveness	0.5	0.5	0	1
Compassion	0.5	0.5	0	1
Kindness	0.5	0.5	0	1
Patience	0.5	0.5	0	1
Humility	0.5	0.5	0	1
Generosity	0.5	0.5	0	1
Integrity	0.5	0.5	0	1
Honesty	0.5	0.5	0	1
Trustworthiness	0.5	0.5	0	1
Reliability	0.5	0.5	0	1
Responsibility	0.5	0.5	0	1
Accountability	0.5	0.5	0	1
Transparency	0.5	0.5	0	1
Openness	0.5	0.5	0	1
Communication	0.5	0.5	0	1
Listening	0.5	0.5	0	1
Empathy	0.5	0.5	0	1
Understanding	0.5	0.5	0	1
Respect	0.5	0.5	0	1
Kindness	0.5	0.5	0	1
Patience	0.5	0.5	0	1
Humility	0.5	0.5	0	1
Generosity	0.5	0.5	0	1
Integrity	0.5	0.5	0	1
Honesty	0.5	0.5	0	1
Trustworthiness	0.5	0.5	0	1
Reliability	0.5	0.5	0	1
Responsibility	0.5	0.5	0	1
Accountability	0.5	0.5	0	1
Transparency	0.5	0.5	0	1
Openness	0.5	0.5	0	1
Communication	0.5	0.5	0	1
Listening	0.5	0.5	0	1
Empathy	0.5	0.5	0	1
Understanding	0.5	0.5	0	1
Respect	0.5	0.5	0	1
Kindness	0.5	0.5	0	1
Patience	0.5	0.5	0	1
Humility	0.5	0.5	0	1
Generosity	0.5	0.5	0	1
Integrity	0.5	0.5	0	1
Honesty	0.5	0.5	0	1
Trustworthiness	0.5	0.5	0	1
Reliability	0.5	0.5	0	1
Responsibility	0.5	0.5	0	1
Accountability	0.5	0.5	0	1
Transparency	0.5	0.5	0	1
Openness	0.5	0.5	0	1
Communication	0.5	0.5	0	1
Listening	0.5	0.5	0	1
Empathy	0.5	0.5	0	1
Understanding	0.5	0.5	0	1
Respect	0.5	0.5	0	1
Kindness	0.5	0.5	0	1
Patience	0.5	0.5	0	1
Humility	0.5	0.5	0	1
Generosity	0.5	0.5	0	1
Integrity	0.5	0.5	0	1
Honesty	0.5	0.5	0	1
Trustworthiness	0.5	0.5	0	1
Reliability	0.5	0.5	0	1
Responsibility	0.5	0.5	0	1
Accountability	0.5	0.5	0	1
Transparency	0.5	0.5		

AA525807, AL224619, H82636, AA508036, AI918419, AL121287, AA829490, T73699, AW304536, AA502498, AL110215, AP000501, AL135744, AC005484, Z96243, U17576, AL022238, AL109627, AC003982, AL079342, L78833, AC005224, AC004821, AL035659, AP000689, AL021808, AC002432, AC002126, AB003151, Z93241, AC004595, AC002301, AC005288, AC009946, AC006241, AC006966, AF196969, D87675, AL031447, AP000503, AP000569, AC007308, AC002395, AC002565, AL008726, AL121769, U62293, AC003007, U63721, AL031848, AL035405, AC009247, AC005736, AC004149, AC004805, AC004817, AC006480, AC006430, AC005632, AC005340, AC005102, Z98036, AC005231, AF072711, AF111169, AF047825, AC006111, AC005041, AF134726, Z85987, AC005527, AC006509, AC000085, AL049631, AC004955, AC005005, AC005920, AF001548, AC005088, U80017, AC002470, AC004893, AC005189, AL133163, AP000141, AC006285, AC004531, AC004650, AC003029, AL121748, AC003104, AC005839, AC004143, AC006013, AC007011, AC004655, AC003108, AC004000, AP000305, AL049761, AR004690, AL049696, AL121754, AC005015, AC004491, Z98742, AL031427, AC005057, U62292, AC005280, AC002299, AC005089, Z86090, AC007242, AC005921, AL049834, AC005488, AC004922, AL035462, AC006315, AC005944, AL031296, Z82206, AP000030, AC004834, AC005306, AC005625, AL035420, AF128893, Z98752, Z83826, AL022323, AP000300, AP000497, AC005529, AL035652, AP000555, AL078581, AD000812, AC006030, AC005332, AL022311, AF207550, AC007066, AL021707, AC002991, AC004106, AJ009610, Z84476, AC005602, AC005512, AC020663, AL031589, AL031311, AL035089, AL049856, AC005907, AC005409, AC005666, AP000031, AP000047, AL021939, AL008730, AP001060, AL109839, AC005701, AL109865, AC004596, AP001052, AL121655, AL031685, AP000301, AC007326, AL049766, AC005730, AL109984, Z93930, AC004597, AC005777, AP000066, AC002378, AL031681, AD001502, AC002544, AL034402, AC007226, Z98884, AC005694, AL049839, AC005755, AP000359, AL121653, AC005082, AC005790, AP000557, AD000092, AC004560, AC006441, AC004477, AC004755, U07562, AL049757, AL049779, AP000502, Z95116, AL096701, AC003080, AC004973, AC004263, AL118516, AC007566, AC010205, AL035400, AL031577, AC012085, AC007649, AC007390, AL022327, AC004985, AL133448, AC007565, AC006277, AC005553, AC004605, AC005048, AC006271, AC007193, AC006538, AP000045, AP000113, X96421, AC004659, Z84480, AL022326, AC005585, AL022165, AC005932, AC006115, AC003962, AC004858, AC007546, AP000114, AL031291, AF111168, AC005911, Z82901, AC003663, AL031295, AL009183, AC005257, and AP000115.
AA182703, AL1275893, AI312591, AW081314, AI890203, AI192785, AL1376547, AI741045, AI382078, R19548, AI769235, AA927822, AI338465, AI074270, AI807959, AI370112, AA725073, AI051770, AA598799, AI023985, AA129014, and AB033055.
AA807514, AA348342, AA280883, AA827687, AW188273, AI565521, AA037530, W21023, C15105, AA232584, R29652, L23320, X69821, AF040250, Z22642, and A37862.
AA449734, AA427598, AW444685, AW449906, AI795836, and AI089108.
AA878223, and AC003663.
AA429954, and R16779.

HMEKO03	412	924172	1 - 395	15 - 409	AI243588, AA521176, and AA280671.
HMEKJ40	413	711187	1 - 327	15 - 341	R09458.
HMEKH73	414	923893	1 - 581	15 - 595	AA424126, AA446776, W77829, W72193, AA662191, AI678126, AI927019, AA248915, AA432090, AA424038, AA282355, and AB032954.
HMEKC72	415	760637	1 - 407	15 - 421	AI668582, AI733530, and H27330.
HMEJW50	416	724396	1 - 301	15 - 315	T87438.
HMEJH81	417	777945	1 - 485	15 - 499	W86772, AI057384, AA778241, AA447568, AA806287, and AI566837.
HMEJF25	418	678131	1 - 424	15 - 438	AA135338.
HMEIV22	419	674611	1 - 490	15 - 504	H38908.
HMEIS07	420	922703	1 - 374	15 - 388	AW245436, AA527311, AI590091, AI983410, AI091713, AI346717, AA398347, AW080183, AA512959, AW005087, AA058970, AW128940, AI089097, T53469, AA657971, AA025746, N92837, AI472329, AI056168, AI802669, AI468989, AA232933, AI250941, AI500614, AI245602, AA764762, AI269006, AI470306, R43430, AI984174, R40489, AA025898, AA858342, AI687814, AA836964, and AA410714.
HMEIA06	421	935966	1 - 240	15 - 254	AB020860, AC006312, and AL096702.
HMEGK14	422	796443	1 - 710	15 - 724	AA005070, W90563, and AA379086.
HMEGH92	423	790629	1 - 481	15 - 495	AA252756.
HMEFD72	424	766185	1 - 398	15 - 412	AA215329.
HMEEL38	425	733649	1 - 588	15 - 602	N52853, T69848, T70384, AA307828, AA325554, N27041, AA361398, R88485, T26592, AW368306, AW408494, H08241, T80355, W28865, AA447395, AA426157, AW138787, AW236918, AA382949, AW362933, AI908338, R17363, AA164794, and T23060.
HMEDR76	426	529897	1 - 161	15 - 175	AC002357.
HMECQ10	427	968500	1 - 311	15 - 325	AA306797, AA278590, AA315885, and Z56144.
HMECH43	428	715568	1 - 291	15 - 305	AA211808, AW405016, R01660, AI246409, AI049534, AW277135, AI146266, AW249720, and AC004690.
HMEBY95	429	796058	1 - 337	15 - 351	AA251522.
HMEBG01	430	921763	1 - 281	15 - 295	AI668626, AW029612, and H42199.
HMEAN12	431	655220	1 - 390	15 - 404	R81266, AW385354, AA968573, AW069537, R81167, AW391238, and AA570569.
HMEAI38	432	709207	1 - 490	15 - 504	H93846, and AL096710.
HMEAH31	433	698403	1 - 249	15 - 263	T69104, T69027, AA092794, AI907131, AA622562, AF149045, and AF149046.
HMEAE24	434	880925	1 - 156	15 - 170	AI354388, AW089663, H98198, AI612070, AI611533, AA834814, AL041749, AA769429, AA224238, AW008089, AA715280, R92172, AA173041, AI933505, AL096774, AC005231, AC003982, AL022320, AC003080, AL008712, AC007666, AF196779, AL096791, M89651, AC008122, AL021397, Z93783, AC003007, AC005907, AP000114, AC002369, AF207550, AC005480, AC006166, AC005324, AL035454, AC004983, AC005632, Z97630, AC005971, AC004056, AC004796, AC006449, AL121825, AC007216, AC002316, AL022333, AC004491, AC004966, AC004167, AC004596, AC004820.

Year	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100																																																																																																																																																																																																
Population (millions)	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.2	3.3	3.4	3.5	3.6	3.7	3.8	3.9	4.0	4.1	4.2	4.3	4.4	4.5	4.6	4.7	4.8	4.9	5.0	5.1	5.2	5.3	5.4	5.5	5.6	5.7	5.8	5.9	6.0	6.1	6.2	6.3	6.4	6.5	6.6	6.7	6.8	6.9	7.0	7.1	7.2	7.3	7.4	7.5	7.6	7.7	7.8	7.9	8.0	8.1	8.2	8.3	8.4	8.5	8.6	8.7	8.8	8.9	9.0	9.1	9.2	9.3	9.4	9.5	9.6	9.7	9.8	9.9	10.0	10.1	10.2	10.3	10.4	10.5	10.6	10.7	10.8	10.9	11.0	11.1	11.2	11.3	11.4	11.5	11.6	11.7	11.8	11.9	12.0	12.1	12.2	12.3	12.4	12.5	12.6	12.7	12.8	12.9	13.0	13.1	13.2	13.3	13.4	13.5	13.6	13.7	13.8	13.9	14.0	14.1	14.2	14.3	14.4	14.5	14.6	14.7	14.8	14.9	15.0	15.1	15.2	15.3	15.4	15.5	15.6	15.7	15.8	15.9	16.0	16.1	16.2	16.3	16.4	16.5	16.6	16.7	16.8	16.9	17.0	17.1	17.2	17.3	17.4	17.5	17.6	17.7	17.8	17.9	18.0	18.1	18.2	18.3	18.4	18.5	18.6	18.7	18.8	18.9	19.0	19.1	19.2	19.3	19.4	19.5	19.6	19.7	19.8	19.9	20.0	20.1	20.2	20.3	20.4	20.5	20.6	20.7	20.8	20.9	21.0	21.1	21.2	21.3	21.4	21.5	21.6	21.7	21.8	21.9	22.0	22.1	22.2	22.3	22.4	22.5	22.6	22.7	22.8	22.9	23.0	23.1	23.2	23.3	23.4	23.5	23.6	23.7	23.8	23.9	24.0	24.1	24.2	24.3	24.4	24.5	24.6	24.7	24.8	24.9	25.0	25.1	25.2	25.3	25.4	25.5	25.6	25.7	25.8	25.9	26.0	26.1	26.2	26.3	26.4	26.5	26.6	26.7	26.8	26.9	27.0	27.1	27.2	27.3	27.4	27.5	27.6	27.7	27.8	27.9	28.0	28.1	28.2	28.3	28.4	28.5	28.6	28.7	28.8	28.9	29.0	29.1	29.2	29.3	29.4	29.5	29.6	29.7	29.8	29.9	30.0	30.1	30.2	30.3	30.4	30.5	30.6	30.7	30.8	30.9	31.0	31.1	31.2	31.3	31.4

HMEAE01	435	916744	1 - 299	15 - 313	<p>AC006059, Z97056, AL031311, AB023051, Z97054, AL049611, AF053356, AC005670, AL049780, AC005837, AL031685, Z68756, AL035249, AL021707, AL049694, AP000512, Z69705, AC004812, AP000501, AC004754, AL109827, AC004913, AC004859, AL049759, AC006026, AL008721, AC005015, AP000692, AC004236, AL133448, AC005332, AC005924, AC005291, AC008372, Z83847, AC004706, AC005004, AC005102, AC004381, AC005598, AC004000, AL031276, AC005071, AC005372, AC003030, AC004520, AL135744, AC006125, AC006130, AC006515, AC007546, AC005253, AC002544, Z82176, AL022311, AC006273, AP000030, Z93017, AC020663, AL021453, Z82201, AC006071, AC007011, AC005399, AL020997, Z99716, AC002045, U96629, AC007052, AC005094, AC005081, AL109963, AC005484, AC005529, AC004560, AC005006, AC005881, Z84469, AC004510, AC006371, Z81370, AL035587, AC004084, AP000251, AC006312, AC005261, AC007487, AC005920, AL049830, AC002310, AC002365, AC005578, AC002565, AC003043, AL049757, AL021154, AC005089, AC006079, AC005288, AC004858, and U95742.</p> <p>AL135643, F13749, A1791610, A1751162, A1821467, A162288, AA230143, A1696793, A1073888, AA563770, AA669709, AA176605, AA579184, AW168479, AW440976, AW168483, AW236342, AW368401, AA828867, AA574286, A1301218, H91844, A1076236, AA360260, A1886365, AW271917, AA657835, A1061111, A1904944, AA357285, A1982634, AA837715, AA331482, A1648558, F23327, AL043098, AL046746, A1753365, AA366601, AA228263, AA608520, AA828047, AA992126, AA847099, AA311156, N59562, AW157005, AA130647, A1028510, AA743811, A1090334, H62670, AA865262, AW384449, A1358089, AA365586, A1566408, H73704, AA569065, A1309384, AW270258, AA747243, H15032, N23260, AA557911, H05821, AA573578, AA365587, AA467730, AA515677, AA761795, F31654, H63806, AA457249, A1357551, AW191886, A1244254, R24915, H53168, F25203, AW327624, AA772016, AL037952, AW327282, A1291124, AA302980, AA578967, AA574442, AA574353, A1709174, A1811460, F32893, AA513234, AA192330, A1291268, A1539776, R92640, AA077776, H79308, AL031393, AC005808, AC005154, AC004263, AL031053, AC002470, AC010582, AC005988, AC005940, AL031587, AC009516, AL034553, AL008635, AL022165, AL096818, AC007308, AC003982, AL031311, Z84480, AL121825, AC004983, AC002425, AC012627, Z99289, AC005971, AC005057, AP000049, AP000116, L44140, AP000311, M30688, AL133163, AC003964, AL049576, AL117536, AC006511, AL122003, AC006430, AC004072, AL109799, AC004517, AF037222, AL034420, AC006273, AP000692, Z84469, AP000503, AC006077, AL008718, AL022316, AF111163, AF147277, AC005274, AL031589, AC010197, AL022476, AC002394, AC007774, AL049794, AC005618, AF134726, AC004134, AC007406, AC004837, AC000353, AC002563, U67093, AR028159, AP000557, AC004706, AC004904, AC006241, AL031774, AC004887, AP000689, AC005856, AJ133269, AL031584, AL049589, AC006538, AC006530, AC005410, AB003151, AP000045, AC006063, U14702, AC008033, L47234, AC006088, AC004232, AC004080, AL035530, AC002480, Z69917, AC007041, AP000509, U73649, AL049869, AL096701, AC005736, AC006254, AC005209, AP000511, AC007564, AC006537, AC005062, AL136295, AC006509, AL096766, AC006356, AC002301, AC003692, AL109963, AP000512, AC005696, AC004985, AF117829,</p>
---------	-----	--------	---------	----------	--

[illegible]

					AC004805, AC005568, AC003041, AC002350, AC008064, AC006039, AC007878, AC004235, AL023803, AC004596, AC002418, Z46936, AC009509, AC003663, AC005874, AF134471, AP000008, AC003667, AC000052, AC005531, AC006277, AC004386, AL031848, AC005207, AC006211, AP000704, AL034555, AC004785, Y10196, AJ229043, U14703, AC004003, AC004019, AP000151, AP000010, AC006449, AL049775, Z99716, AC004216, AC002542, AL049198, AF111168, AC002420, U80459, AC001642, AL031295, AC007731, AL020995, AL117340, AC006014, AC003992, AC005500, AC005690, AF196779, AL022320, AB023050, AL031447, AC007263, AC002430, AC007993, AC004814, D84394, AC007684, AC007676, AC001643, X54486, Z49250, AL021397, AC005178, AL035420, AB023051, AC005632, AP000251, AC007384, AC003010, AP000513, AC005488, AC005786, U62293, U63721, AL022578, AC005184, AC006977, AF129756, AL049780, AL034548, AP000555, AC004741, AP001059, AC005839, AL109623, AP000023, AC005746, AC004832, AL133246, AC007226, AP000553, AL133245, AC008040, AC006238, AC004854, AP000030, AP000211, AC006079, AL078472, AP001053, AC007917, AC005799, AP000552, AC005288, AC004236, AC006071, AC003664, and AC003043.
HMEAD86	436	785802	1 - 344	15 - 358	AA251058, and AA807055.
HMEAD47	437	720638	1 - 407	15 - 421	T79734.
HMEAA17	438	921765	1 - 464	15 - 478	HI0273.
HHFUB83	439	800580	1 - 277	15 - 291	
HHFOU02	440	918070	1 - 715	15 - 729	AW296654, AI498432, AI498430, AI758371, W73054, and U82695.
HHFOL43	441	974002	1 - 498	15 - 512	
HHFOLK10	442	961346	1 - 342	15 - 356	AA280678, AI128307, AI890650, AI538161, AW190537, AI207930, AI358856, AI913895, AI114875, AA564510, AA401182, AA225175, AW273466, AI358915, AI743933, AI037933, AA604799, N72011, AA362732, AW276723, AW276763, AW157028, AA745570, R84431, AA864787, AI479583, AA085683, AA486832, AA487152, AC004883, AC006160, AC004217, AL022575, AC002070, AL034343, AL080241, U55729, AL021578, L11910, Z93930, AC002994, Z95114, AC002096, AL109839, U91322, AF111168, AC005291, Z85997, AL031733, AP000501, AC005274, AD000812, AL035398, AC002126, AF047825, AC007308, AC007160, AL021579, AL031280, AL035690, AC005875, AC005482, AL024507, Z77249, AF029081, AP000047, AP000115, AL031285, AC007052, AC005562, AC000075, AL033392, Z99916, AC004813, AC005664, AC005844, AC004985, AC000097, AC007386, AC006547, AF067844, AC004491, U82828, Z83822, AL009181, AC003669, AF165926, AL049780, AC007055, AF053356, AC001227, AC002101, AC005074, AC005808, AP000260, AL008637, AL034421, AP000099, AC004087, AL049759, and AC005014.
HHFENJ05	443	930899	1 - 430	15 - 444	AI538184, AW007485, AI363114, AI953529, AA910212, AI695319, AA910204, AL121657, and AF214736.
HHFMX34	444	945385	1 - 453	15 - 467	
HHFLU06	445	857884	1 - 316	15 - 330	
HHFELT84	446	857890	1 - 460	15 - 474	AA015945.

HHFL108	447	958103	1 - 399	15 - 413	AA664191.
HHFLJ51	448	857898	1 - 453	15 - 467	H55219, AL040362, and Z80897.
HHFLJ10	449	963162	1 - 652	15 - 666	
HHFLJ07	450	952081	1 - 588	15 - 602	AF064860, AC003109, AC005189, AP000010, AP000151, AP000215, AC005529, AP000556, AC004882, AC002350, AC006160, and AP000337.
HHFLH62	451	857908	1 - 314	15 - 328	AA460477.
HHFLE12	452	969531	1 - 498	15 - 512	AI191562, AA297666, AI887235, AI476049, AA659832, AI272052, AA595499, AW167154, AI598003, AW194325, AW023302, AI859280, AL031650, AC012627, AC002519, AC007308, AC002369, AC006011, AL020993, AL109628, U91323, D87675, AC005037, AP000555, AC006236, AC005920, AL049776, AP000695, AL133246, AC005231, AC006001, Z85986, AC007546, AC005094, Z98200, AC005332, AC007216, AL049856, AC004659, AC007066, AF029308, AC005399, AC007690, AP000337, AP000240, AL035495, AC007227, AC002310, AC005548, AC007225, Z97056, AC006538, AC007999, AC005005, Z95113, AC004104, AC005899, AP000251, AC005225, Z95115, AL022476, Z84480, AP000212, AP000134, AL133163, AC005833, AC005664, AC005089, AP000030, AC002316, AC006115, AC005295, AL023575, AC004884, AC004477, AL121769, AC004222, AL049712, U91328, Z93017, AC003687, AC002082, AC005288, AC006064, AC005988, AC007285, Z94721, AC002546, AC002289, AC007151, AC009509, AC004474, AL121595, AC004815, AL021154, AR036572, AC002477, AP000216, AC004526, AC006160, AF053356, AP000350, AC007450, AL139054, AL133485, AC005409, Z95118, AP000692, AC000025, AF178030, AL049872, AC005527, AC003109, AC005971, AC004134, AB023049, AP000359, AF111167, AC005859, AP000338, AL031230, AC007036, AC002347, AC004966, Z97632, AP000513, AP000248, AL031427, AC006509, AC005412, AC003082, AC006077, AC007041, and AC004882.
HHFKX28	453	971102	1 - 844	15 - 858	AI740820, AA482031, AI565169, AI628285, AA947029, AW190175, N50136, AA707674, AI332610, AA223261, AI189165, AA767472, AI276839, AA804584, AA894857, N59367, R46372, N72682, AA774827, AA219127, AW188325, W02461, AA417592, D29223, AA322537, R08745, AA315300, AW377015, H60482, R08746, AA939277, AA872005, AI810734, AW118290, T30177, D29202, R68574, AI167609, and AI832198.
HHFKB24	454	887025	1 - 420	15 - 434	
HHFJM64	455	958384	1 - 1457	15 - 1471	AA263151, W39493, AW292417, AI703130, Z41877, R67724, and AA578849.
HHFIA58	456	858011	1 - 406	15 - 420	Z47297, and AJ011930.
HHFIA13	457	657405	1 - 410	15 - 424	AA001994, AL046468, AW367920, AA057078, W96490, D87443, and AL078475.
HHFHY95	458	795053	1 - 384	15 - 398	T62723, AA282638, and AC005839.
HHFHR63	459	745215	1 - 630	15 - 644	H58858, R20870, and AC005177.
HHFHP68	460	753144	1 - 479	15 - 493	R34604, AI472929, N67108, AI312095, AI077416, AI080088, AW000975, AI218568, AA305614, AW270427, AA922273, AW268569, and AL021920.
HHFHN74	461	765758	1 - 786	15 - 800	HI1548, T76951, T65348, H23112, H08748, HI8004, FI3243, FI1922, R39092, and T16788.



HHFHM22	462	674841	1 - 420	15 - 434	H05872, and AC006487.
HHFHJ90	463	675218	1 - 437	15 - 451	R02560.
HHFHD38	464	709082	1 - 259	15 - 273	H65332, and H25516.
HHFGX13	465	656806	1 - 694	15 - 708	AW020496.
HHFGX03	466	924753	1 - 421	15 - 435	AA731491, AW088849, A1570996, F32413, A1827137, and AW386082.
HHFGR31	467	953204	1 - 391	15 - 405	W07246, AA423994, and AL049795.
HHFGR30	468	692887	1 - 335	15 - 349	AA167587.
HHFGP91	469	800328	1 - 845	15 - 859	AI208872, AI651750, AA367948, AA852183, AA383420, AI948798, AC004453, and AB028991.
HHFGP69	470	918393	1 - 699	15 - 713	AA570361, AA173853, AW239179, and AF117756.
HHFGL77	471	490379	1 - 494	15 - 508	AI688557, AI799116, AI202945, AI358494, AI697899, AW293458, AI698391, AA765656, AI537677, AI918370, AI621341, AI560227, AI696570, AA824496, AI689614, AW129616, AI559976, AA488429, AI537643, AI891102, AI289400, AI570966, AI953765, AI590043, AI445620, AI611743, AI696583, AI918955, AI537244, AI927233, AI554516, AI874107, AI801793, AI345688, AI491710, AI863002, AI590020, AI475371, AI138386, AI819663, AI433611, AW025279, AW029186, AI800171, AL042944, AW162194, AI567769, AI333104, N49165, AI589004, AI819014, AW105431, AI581139, AI955945, AL119863, AL045672, AI648458, AI567582, AL039011, AI690620, AW059765, AW103726, AI591228, AI635950, AI557426, AW022636, AI075658, AW079432, AI434223, AI670895, H89138, AI536836, AW089275, AW105087, AI560545, AI859644, AI912434, AI873998, AA665587, AI952306, AI499285, AI699823, AW104141, AI589428, AI886355, AI244343, AI474646, AI285439, AI202203, AW148303, AI933992, AI370623, AW055252, AI818358, AI287449, AI925404, AI335231, AI539260, AI287476, AI445611, AI570857, R65859, AW169626, AI095003, AA732937, AL119791, AI572730, AI267532, AI499986, AI702902, AI249877, AI277008, AA641818, AI538885, AI633125, AL043975, AI860027, AI538564, AI434242, AW152182, AI648699, AI345477, AA916133, AI345415, AI440238, AI583578, AI473845, AA602414, AI434255, AI801619, AW305233, R80916, AL039716, AI921087, AA693331, AI473240, AI783792, AI619820, AI251485, AI561356, AI096771, AL121365, AI918809, AW081311, AI890576, AI536601, AW161098, AI431962, AW192109, AI950892, AW051088, AW411235, AI254727, AW020164, AI797794, AW265004, AI247082, AI564186, AI274500, AA229532, AW073926, AW082623, AW410259, AI802240, AW007284, AL046466, AI289791, AI619777, AI627714, AW029329, AL047675, AI334445, AI354998, AW084801, AI499325, AI539771, AI696340, AI539690, AI801325, AI671931, AW131065, AW105383, AA744713, AI801536, AI884318, AA749425, AI803935, AL045997, AW130134, AI127565, AW195253, AW080379, AW131999, AI539723, AA808175, AW411351, AA575874, AL038529, AW059568, AI567993, AW083775, AL047344, N25033, AA678835, AI499581, AI521799, AI379711, AI434731, AL135047, AW301409, AI225000, AI800159, AW080374, AL040243, AI918435, AA824435, AI929108, AI811912, AA767144, AI656270, AI371442, AW411265, AW073865, AW085786, AI620287, AI765469, AW410902, AW243876, AI568773, AW410696, AI540674, AI355779, AW005612, AI540832, AW170773, AW087816, AI499279, AW075382, AI952249, AI685517, AI564290, AI697178, AW020397, AI874166,

HHFGH81	472	778193	1 - 329	15 - 343	<p>                     AI365256, AI887163, AI521095, AI954721, AF076464, I89947, I48978, AF108357, AI133559, AF119336, AF035161, AF113690, A08910, A08909, AF094480, A65340, A08908, X83544, Y00093, X72889, AI137478, AI137533, AI133619, AL049447, Z97214, AF077051, A1005690, AR038854, X97332, AF067790, X79812, AL122050, M92439, AF126372, AL049423, AI137271, AI117460, A08913, AI133084, AF131773, E12579, M64936, AF162270, AF031903, X53777, AL049347, AF013214, AI18777, A65341, A77033, A77035, X80340, AL050278, S71381, X66862, AL122098, AL110221, AI133568, AR034821, I48979, A21103, AL110262, Y10823, X61399, A76337, AF087943, A90844, AF118090, AI137529, L04859, AF111112, U70981, AI117463, AJ238278, AF126488, AF026008, AJ003118, AF177401, E12580, A08912, A08911, AI133637, AF067728, U75604, AF111851, AI137554, AF026124, S79832, AL050138, U58653, AF022363, AF032666, S77771, AF055917, AR011880, AF158248, AL035458, AL110159, AR068753, A07588, AF065135, I17544, AI137574, AF044323, U80742, AL050393, AF102578, A58524, A58523, S76508, AC002467, AF109155, X65873, AR068751, AF028823, AL050024, E04233, Y11254, AF176651, Z37987, AI133016, S68736, AI137527, A93016, AL078630, I00734, AF059612, Z35309, I89931, AL049339, AI133088, A76335, L13297, U35146, I92592, A91160, E00617, E00717, E00778, M79462, U77594, E12747, AI137627, I49625, X66871, AI137298, AI122104, AF162782, AL080110, AF076633, AL080158, AI122100, I30339, I30334, AL050116, AI137722, AI122118, AF139986, X93495, E01614, E13364, AF139373, A08907, AF126247, AL049938, AL050277, AF067420, L30117, I26207, AF090886, AF124728, AI110228, X63574, AJ012755, A18788, AF003737, Y16258, E03348, Y16257, E02756, Y16256, U67958, AL080159, I33392, Z82022, AB025103, AR059958, AB007812, AF047716, E01314, S82852, AF112208, AI117457, AL050149, AF151109, AF061981, U78525, AF090901, AL080148, E06743, AI117435, AI137665, A91162, AF008439, AF104032, X99226, AI137550, I89934, AL049324, AF118094, S36676, E03349, AR000496, D44497, U39656, AF080622, AF090900, AL080156, AL096728, AF180525, I32738, AF061573, AF019298, AF000145, A08916, AF090432, AL049276, D16301, AR068466, AF090934, L19437, S63521, AI133558, AI117587, I46765, A49139, AR060156, A58545, I09499, AF039138, AF039137, AI137660, AL080129, AI133049, AI122121, AI137476, A17115, A18079, AI133665, L24896, and AI137300.                 </p>
HHFGH43	473	573495	1 - 521	15 - 535	R56912, AA888618, AA460448, AA554077, AA458950, AI865116, AA019611, AA410313, R55713, AA081407, AA724795, AA648850, AA019450, AA552550, AA177050, AA854071, AA152380, AA844117, W37842, AA192171, AA468546, AI269758, AA987694, AA586585, AA746215, R39789, R43005, AA216405, AA059379, and AR038867.
HHFFZ50	474	513773	1 - 643	15 - 657	R73359, R83331, H44513, H44528, and AW376878.
HHFFT05	475	932675	1 - 123	15 - 137	
HHFFT01	476	880757	1 - 388	15 - 402	X97818.
HHFFR95	477	796677	1 - 427	15 - 441	H05133, R36902, and Z40451.
HHFFR75	478	766630	1 - 493	15 - 507	AA812058, AI120483, AI859744, AA593428, AI368873, AA683031, AW008212, AA601218,

					AI917156, AI951863, AL041895, AI471543, AI588856, AW117936, AI567674, AA568310, AA745302, AI204309, AI679782, AA483004, AI791180, F25658, AA317973, AI102235, AI821138, AA736713, AA523695, AA805845, AA339076, AI823679, AA610255, AW238495, AI915081, AA348248, W32993, AL037856, AI459943, AA525331, AC004019, AC005288, AC005696, AC006014, AC006111, AC005488, Z85987, AC006023, AC004812, AC004253, AC000052, AC005736, AL023575, AC003665, AC006077, AL008718, AP001053, AF129756, AP000555, AL035681, Z98200, AC005295, AC005480, AF045555, AC004685, AC005740, AC006160, AL049712, AC006480, AL022323, AC006449, AL035659, AL049636, AL008582, AC005081, Y07848, AL035249, AL096701, AF111168, AC009509, AC004967, AL022329, AP000355, U52112, AC006511, AC007917, AC006948, AC005520, AL121652, AF196779, AC005399, AP000086, AC002551, AL096791, AF196972, AL022320, AC005182, AC004216, AL024498, AC004491, AL031311, AC005728, AF001549, AC005330, AL009051, AC005225, AC000353, AL121603, AC007011, AL009031, AC002326, AL022324, Z83844, AC005821, AF139813, AC004973, AF111169, AL033392, AC005412, AC020663, AC004126, AC002477, Z95113, AC002059, AP000113, AC006538, AF205588, AL031665, AC005755, AL117354, AL031984, AF196969, AC006001, AC005216, AF060568, AC004841, AL049643, AC004228, AC005954, AC005088, U85195, AL009181, AL049694, Z98036, AC002036, AC004477, AC002449, AC007240, AC002400, AC003002, AL031291, AC005924, AC006120, AC005231, AE000658, AC004638, AL022318, AC002316, AC006211, AC005031, AC004878, AC005632, Z83840, AC002429, AC007308, AC007314, AC000026, AP000553, AL080243, AF111167, AL133353, AL109758, AC006126, AC006277, AL135744, AL080242, AC004876, AJ003147, AC005484, AL139054, AL031680, AL023807, AC007371, AC005527, AC002470, AC002117, AL034423, AC005280, AP000556, AC003030, AC005072, U80017, AF001552, Z99943, AL035587, AL135960, AC004531, AJ131016, AC006312, AL031848, AC004106, U62317, AC007774, AF047825, AC005899, AL049776, AC007193, AC000003, AC006455, AC002519, AC004796, AC004149, AC007993, AC002300, AC002301, AC004585, AC003689, AF088219, AC005837, AC018633, AC007055, U62293, AL034420, AC002126, AL021707, AL078476, AC002072, U95740, AC004230, AL021391, U47924, AC005071, AL022476, U95742, AC007227, AC005531, AC007263, AF030453, AC007226, AC002350, AJ246003, AL035703, AC005726, AL031597, AL020993, AP000098, AC005227, AC007934, AB023049, Z99716, AC007731, AC004815, AC003025, AC002369, AC004991, AC004552, AC002310, AC007051, AC000134, AC007546, AL031295, AC007065, AC004448, AC005500, AC005779, AP000504, AC000097, AC002395, AC002563, AC005529, AC007216, AC005015, AC004651, AC008044, AC005104, AC007387, AC005057, AC007386, AL021546, AC006547, AC005036, AC005184, AC006088, AP000694, AL117258, Z83845, AL023284, Z83819, AC006942, and AC007919.
				15 - 314	AA234902.
HHFFR32	479	699723	1 - 300	15 - 450	AI493068, and AL024498.
HHFFP17	480	880667	1 - 436	15 - 324	
HHFFO46	481	530501	1 - 310	15 - 472	H65231.
HHFFM05	482	932738	1 - 458		

HHFFL66	483	530503	1 - 138	15 - 152		AA523376.
HHFFK30	484	858051	1 - 264	15 - 278		N80165.
HHFFI59	485	739587	1 - 510	15 - 524		T99205, Z30071, T96773, W90185, T98729, and AB020638.
HHFFI08	486	960254	1 - 664	15 - 678		AA446796, W24782, AA476520, AR050683, AF104922, and AF033855.
HHFFH76	487	767623	1 - 534	15 - 548		A1190744, AA85487, W69997, AA861491, and AW104913.
HHFFG82	488	530662	1 - 313	15 - 327		AA020789, AA001701, AW452804, A1375739, N78745, AW027244, AA731202, A1672595, AA279753, W16651, AA894594, AA019106, and A1383433.
HHFFF92	489	790572	1 - 782	15 - 796		AA699816.
HHFFF07	490	954258	1 - 564	15 - 578		N77740, A1623295, AW385146, AW089125, and AA961452.
HHFEB86	491	785653	1 - 648	15 - 662		AA001465, AA100624, AA314602, H04982, AA351833, and AL021997.
HHFDN80	492	781634	1 - 413	15 - 427		
HHFDI82	493	499010	1 - 506	15 - 520		R36135, AW001383, AA243866, AB014580, and AL078593.
HHFDH26	494	685188	1 - 427	15 - 441		AA664192, T71388, and H60250.
HHFDC10	495	968647	1 - 725	15 - 739		T99849.
HHFDA13	496	667804	1 - 324	15 - 338		AA347973, H57111, and A1217870.
HHFCT63	497	572784	1 - 251	15 - 265		AA347868, W01938, A1697383, AA347867, AA017179, W92433, W92446, H86133, AA018375, W96275, F31654, AW117723, AW117704, N22058, AW026276, A1056174, AA993165, A1754286, W45073, H05449, A1274006, A1921692, A1744381, A1567831, A1445768, AA586553, D25667, A1889648, A1613487, A1336206, AA516214, AA515728, A1635819, AA827978, AA515924, AA570224, A1061147, AA678491, A1820796, Z82202, AC002036, AL035541, Z97053, Z49235, Z98946, AC005574, AC000379, AC002312, AC004408, AC009464, AC003065, AL009172, AC005033, AC006543, AL020989, AC007363, AC005919, AC005180, AC006544, AC004985, AL035419, AC004694, AC005219, AC012398, AC011718, AC008132, AC007325, AC004148, AC006455, AC006346, AL121769, AC004012, AC009509, AL034423, AL122020, AC003071, AC004777, AL132826, AL096791, AP000354, AC000387, AL031737, AL031584, AL109807, AC002113, AC000367, AC005781, and AF001552.
HHFCO13	499	500899	1 - 328	15 - 342		R82827, and AA347799.
HHFCN59	500	739657	1 - 543	15 - 557		W05481, and AA347756.
HHFCN13	501	667805	1 - 387	15 - 401		AA676441, W84872, A1240587, and AA347693.
HHFCM51	502	509631	1 - 139	15 - 153		
HHFCH52	503	911570	1 - 343	15 - 357		AA347276, and AA347277.
HHFCF58	504	575183	1 - 517	15 - 531		R92935, N99077, A1809891, AA347287, A1017165, and N91783.
HHFCE73	505	764763	1 - 404	15 - 418		H69189, R06505, T72747, and AA347297.
HHFCE40	506	712866	1 - 465	15 - 479		R51510, AA825938, AA347032, and A1274639.
HHFCD43	507	714353	1 - 242	15 - 256		AA347175, and AA152489.

HHFCC60	508	739669	1 - 459	15 - 473	H05854, and AA347121.
HHFCC45	509	858066	1 - 203	15 - 217	AA346956, R85908, and H86351.
HHFCC20	510	600231	1 - 591	15 - 605	AA347077, A1829548, A1732247, AA338281, Z82243, AC008072, AC002073, AP000344, AC004463, AC004461, Z83844, AC003037, AC004462, AP000553, AC003662, L77570, AF111169, AC005919, AL031286, AC005049, AC002300, AC004988, AP000692, AL022576, AL031985, AC008009, AF001548, AF050154, AF121781, AC000353, and Z93023.
HHFBW92	511	575156	1 - 539	15 - 553	R92655.
HHFBW63	512	745661	1 - 655	15 - 669	T82006.
HHFBW07	513	954478	1 - 564	15 - 578	A1824082, and A1863730.
HHFBT24	514	508067	1 - 436	15 - 450	H67854, D80014, D51423, D80043, C17376, H67858, D80038, D50995, D51022, H67866, D31458, C14973, T11417, C03092, D58283, F13647, D80022, D51079, C14331, D80247, D80522, D59859, A1525907, D80188, C14389, AA809122, D80251, D57483, C15076, D80366, A1525235, D59889, D80439, D58246, D81030, D59275, D80253, D80258, D80157, D81026, D80269, D80166, D80212, D80268, D80195, AA305578, T03116, D80196, D59467, D59619, D80133, C14227, D80210, D51799, D59503, D80391, D80164, D80240, D59317, D80045, D50979, D80219, C06015, D59787, D80227, D59502, D80064, AA305409, Z33452, T02974, A1525914, A1557774, D59551, A1525922, Z21582, C16955, D80168, C14077, D51213, A1535686, A1535665, A1525920, A1525917, A1525923, D45273, Z30160, C14298, D59627, A1525242, A1525912, A1525227, A1557751, A1525237, A1525215, AA514184, T03048, A1525228, A1525238, A1525216, AR054175, A62300, AR008277, AR008281, AR018138, A82595, AF058696, AB002449, A84916, A62298, I14842, AB028859, AR060385, AJ132110, AR008278, I82448, I79511, and X64588.
HHFBQ94	515	796838	1 - 302	15 - 316	N69862, AA196591, W94557, AW445181, N99038, A1695808, W92641, AA346845, AA181501, AA100353, AA196590, C04454, and AF086522.
HHFBP29	516	710894	1 - 364	15 - 378	T80420, A1056302, T97347, T85277, H61413, N57950, AA378460, AA227248, AW402041, AA346817, AA659435, AA187022, A1065081, T85741, R15986, AA236609, H59659, and AC004382.
HHFBN17	517	589798	1 - 537	15 - 551	A1733856, A1923050, A1791475, AA381011, N54894, AW021735, AA515725, AW024042, A1168167, A1433008, A1580250, A1869978, AA063173, AA584170, AA524687, AA524955, AA838190, AA513893, A1937250, H09071, AA551117, A1491817, AA651632, AW340844, A1061647, AL119713, A1460009, AW074242, AA669155, A1053911, AL041619, AL119921, AL045813, AL037632, AA343946, A1056177, AA362719, AW298643, A1122243, AA610688, AW023865, T99179, A1818231, AA669564, AA330418, AA548058, AW080965, AA983199, AA487180, AA487239, AL134330, AA811208, A1084012, AA323701, A1745553, A1808248, A1732800, AA442105, AW020340, A1613389, AW401372, AW021597, AA410788, AL036382, A1891080, AA769720, AA778992, AA579179, AL119941, AA486131, AL038936, A1801482, AA425118, AL043719, AW022897, N75702, AA010299, AW080777, A1306324, AA305249, AW102846, AA577824, AA864493, AL036070, AA493695, AA507282, AL045053, AA191659, A1821947, A1791211, A1687343, AW338883, A1344822, A1805547, AW403720, A1133164, AA179683, A1567076, AW270343, T52097, AA385798, A1754211, AA343824,

					R6459, AI860587, AI285753, AW302315, AI282511, T06648, H86363, AI149478, AA968680, W28893, AA661921, AP000010, AP000152, AC006055, AP000514, AF156495, AC003688, AB020861, Z93023, AL049745, AL020997, AC002395, AC007091, AC005324, AC007363, U68030, AF200923, Z83840, AC012384, AC005933, AB014077, AC004819, U66341, U66340, AC006139, M33198, AC006322, AL078584, AC007406, Z68755, AL008719, AF047825, AC006512, AJ006996, AC005040, AF129756, AL080241, AF135028, AB022430, AC004778, AL031721, Z93848, AC005568, AC004496, AD001502, U89336, AC002378, AC007199, AC004913, AP000159, AP000017, AP000094, AL049539, AC004150, AC006014, AC004848, AC005630, AP001172, AL122020, AP000237, AC002070, AC007686, AC006111, AC005036, AP000014, X81001, U45984, AF036110, AC005013, AC004497, AC005338, AC005788, Z70280, Z68871, AC005327, AC004030, AC006277, AC005783, AC005337, AL109830, AL078596, AP001051, AL035681, AC004935, AP000504, AP000123, AL022163, Z85987, AL023279, AC002477, AL022152, AC004000, AC005738, AL034554, AC004884, AC007285, AL024497, AC002432, AP001052, AC005880, AL031652, U95740, Z98200, AC019014, AC005730, AP000354, AC005224, AC007546, AF176315, AP000355, AC005772, AC008151, AC005768, AC005234, AC007537, AC005261, AC005035, AL133500, AC007298, AL139165, AC005071, AC005015, Z99716, AF053356, AC000353, AC008372, X91841, AC004638, L20345, AL096794, AC003104, AC005377, AC007324, AC008101, AC008079, AL135744, AC004626, AC004651, AC004760, U62293, U63721, AC007160, AL031651, AC004209, AC002347, AF118808, U96629, U91323, AJ010598, AC004648, AC000086, AL031346, AC004808, AL133246, AC005057, AL122023, AC005019, AC005539, AJ131818, AF024533, AL022320, AL096770, AC005086, AL021154, AC007666, U52112, AC007685, AL022323, AC000052, AP001137, AC004019, AC004526, AC005529, AC007664, AC006255, AC002316, AC005320, AL121593, AC006463, AC007297, AL096678, AL031685, AL031848, AC007308, AL137408, AC002416, U07562, AL031224, AF157816, AB029343, AB023060, AC005606, AP000509, L05367, AL133249, AC002472, AC009498, AF055584, X12641, A00794, AF012654, Y11873, Y12855, AP000307, AL034546, Z68276, AC004028, D00591, AC002106, AF050154, U76377, AP000558, AF146367, AC002540, AL049733, AF104455, AC005661, U82668, AL034555, AF045555, AC006071, AL031597, AF095725, AP000115, AP000499, AB020865, AP000130, AP000208, AL009051, AF124523, AC005846, AC006476, AC002425, AC006409, AP000247, and AC005090.
HHFBM11	518	968002	1 - 503	15 - 517	AI937790, F06528, AA446518, AA446340, AA446353, AA346668, and AA446534.
HHFBD83	519	781525	1 - 590	15 - 604	AA001411, AI160471, AI433691, AI923694, AI625789, AA521470, AA442451, T70594, AA719773, AI377638, AI003601, T96901, AA770076, AA766033, AA346585, AL137658, and AC005343.
HHFBD42	520	712899	1 - 548	15 - 562	AA151917, AI076693, AI801676, R67388, AA346551, R26159, AI240747, AI868514, R22968, AW370946, and AW370942.
HHFBB14	521	522375	1 - 447	15 - 461	T54874, AA346439, and AL050173.
HHFBA11	522	967991	1 - 361	15 - 375	AA346357, and AA346358.
HHFAB62	523	824590	1 - 700	15 - 714	R49113, H17411, AA346385, AA857224, H16518, AW373989, AL046035, AW374001, AW373994,

					AL039790, AW373998, AW364756, AW364750, AW364749, AW373996, Z32887, T18597, A1526078, Z33559, D59751, D50992, A1535639, A1557084, A1557864, A1535660, A1536138, A1557262, A1541205, A1525500, A1525556, A1557533, A1540903, A1525316, A1541356, A1525302, A1557809, A1557317, A1541365, A1541075, A1557731, AA058620, A1557082, H65400, A1525852, A1557258, A1525757, A1541353, A1541321, A1557312, R29657, A1536150, A1557474, A1541034, A1546829, A1557602, A1540974, N71206, Z33585, A1557155, A1547177, A1557543, A1547182, A1557810, A1557234, A1535994, A1557238, A1541450, A1536070, A1525878, A1557041, A1541346, A1557241, A1557408, A1557039, A1541336, A1557154, A1525656, D79991, A62298, A82595, A82593, A62300, AR050070, U94592, Z30183, AF006072, U45328, and AR025466.
HHBGN74	524	765214	1 - 545	15 - 559	AA112990, Z25197, AA214259, AA214140, AA214253, and AF177292.
HHBGJ53	525	909912	1 - 388	15 - 402	N49341, N31123, and AL135424.
HHBGG10	526	963849	1 - 475	15 - 489	AI042164.
HHBGC75	527	767042	1 - 562	15 - 576	AA426500, AA195328, and AA195188.
HHBFT06	528	934826	1 - 537	15 - 551	C18811.
HHBFM77	529	771816	1 - 447	15 - 461	AA064965, and AA129388.
HHBEV93	530	792041	1 - 508	15 - 522	H41579, H23896, and H49582.
HHBES89	531	786667	1 - 453	15 - 467	R35959, H16638, R59412, F13165, T77625, and AC004744.
HHBEM49	532	722337	1 - 718	15 - 732	AA143472, and AI023326.
HHBEG80	533	951688	1 - 430	15 - 444	AI095759, AI493168, AW206042, AI738997, AA921950, AI968444, AW197610, AI079592, AW043953, AW242819, AI744109, AI861845, AI193079, AI668960, AW182363, AI640258, AI092922, AA722466, N21525, W73601, AI140933, AA021471, AI720404, AA708907, AI123135, F30816, AI366183, AI768468, T77057, AI968632, AA045707, AA259196, AA127350, H11390, F32206, AI767021, H24345, AI418672, AW383731, N46355, AA905163, AA018209, F36514, AI564847, T83879, F01296, AA070435, AA921795, R20767, F00344, AA196850, AA311411, AA045706, AI969655, AI824576, AI440239, AI633062, AI250663, AI613038, AI815855, AW193872, AW262565, AL045672, AI923989, AI251876, AL119863, AW080402, AI568138, AI926878, AI590227, AI654037, AI478123, AI698391, AI636588, AW161579, AW129929, AL042745, AI805603, AI890507, AW078800, AW130930, AW020419, AI866002, AI537677, AI673785, AI306705, AL036403, AW051226, AI811785, AI288305, AI873644, AW131331, AW079572, AI580198, AI497733, AI431408, AL040827, F27788, AA225339, AI866770, AI889168, AW151136, AL038605, AI919345, AI954504, AI630252, AI890907, AW089275, AI826225, AI632408, AW087445, AI702073, AL120695, AW162194, AL045266, AI564749, AW088162, AI868204, AA470491, AL037582, AL037602, AI270183, AI470648, AI499381, AW268220, AI620075, AI802542, AI800138, AW050578, AW168031, AI955906, AI312428, AL079963, AW161156, AI334445, AW302965, AI241923, AI345477, AI624548, AW302992, AL036638, AL110306, AI648473, AA572758, AW196105, AI886753, AI269862, AI612750, AI251221, AW022682, AA494167, AI929108, AW302988, AW163823, AI340603, AW074869, AI494201, AI349957, AI491775, AI677796, AI308032, AI284131, AI858827, AI934035,

AI635492, AI874166, AI284517, AI537837, AI621179, AA427700, AI590830, AI254226, AI335209,  
 AI699857, AI860783, AI919500, AI570807, AW004886, AI263331, AI358701, AI624293, AI623941,  
 AI536574, AW238764, AW054931, AI829327, AW268302, AI419650, AW089405, AI569583,  
 AI863382, AW072719, AI433384, AI499285, AA911767, AW169653, AI869377, AL037454,  
 AW087915, AI564259, AL036980, AW169604, AL042744, AI439762, AI251830, AI890806, AI554821,  
 AI812032, AI589267, AW170635, AI524526, AI916419, AI620284, AI571046, AL039086, AI950664,  
 AI638798, AW020095, AI433976, AI889189, AI345745, AI473536, AI612885, AI783504, AL046926,  
 AI445990, AW089572, AW051088, AI886123, AW168485, AI690835, AI554344, AL036804,  
 AI610799, AI679620, AI801325, AW403717, AI863191, AI590686, AI969601, AL118781, AI445992,  
 AL036361, AI886415, AA580663, AI434741, AI886181, AI568870, AI471361, AI590134, AL117457,  
 AF134803, AF134802, L29468, U78525, U42766, X63574, I89947, I48978, AF026124, AF113694,  
 A08916, A08913, A08910, AL049430, AJ238278, A08909, E02349, AL080148, AL122049, A07647,  
 AL050277, AF091084, AL049466, AL049283, AL050024, AF087943, Y09972, AL133016, AF177401,  
 AL137480, AF146568, AL110222, AF028823, AR038854, AL050146, I89931, A08912, AF132676,  
 AF061836, I49625, AL133560, AF097996, AF090900, I48979, AL122093, E06743, AL133080,  
 AF111851, AF111849, AL049464, AL137533, AL122110, U35846, AF079765, AF104032, I00734,  
 AL050393, S61953, AF113019, Y16645, AL049300, AL137557, Y11254, AL122050, AL137459,  
 AF079763, AL133557, AL080127, AL050116, E00617, E00717, E00778, AL117394, AI2297,  
 AL117435, AF057300, AF008439, AF057299, X72889, AR011880, AI000937, AL133640, AL137271,  
 Z72491, AL110221, AF090903, E08631, AF090896, E12747, AF113689, AL049452, AL049314,  
 E07108, S78214, AF113676, AF158248, AL133072, AL137463, AF061943, AF118094, E04233,  
 X84990, AL122098, A93350, U68387, AL117440, AL137521, AL080124, AR020905, AF067728,  
 AL137478, AL137550, AL117583, AL133098, AF017152, Z37987, S68736, X83508, U49908,  
 AF118064, AF118070, AL137560, X70685, AL117460, I09499, A45787, AL096744, AL133014,  
 A08908, AF067790, AF113677, AL110196, AF183393, X92070, AL133075, AF125948, AL133565,  
 A03736, AF026816, AF106862, AR013797, X82434, A65341, A77033, A77035, Z82022, AL050149,  
 AL050108, AL050138, AL137476, A58524, AF078844, A58523, AF119337, AF003737, AF113690,  
 E03348, AF090934, AF017437, AF162270, AL137526, L19437, AL133093, X87582, U67958,  
 AF113699, M30514, AF153205, AR059958, AL117585, L31396, I66342, AF185576, AF090901,  
 Y07905, X93495, E02221, L31397, AL137292, AJ012755, AF054599, U72620, I26207, AF090943,  
 AF061795, AF151685, U80742, AB019565, AL137556, AL080074, AF113013, AL133568, AL133113,  
 AJ006417, AL122121, D83032, AL080060, AF100931, I33392, AL049382, AL117416, AL110225,  
 AL137488, AL050092, X96540, AF111112, Y10080, AF126247, I03321, AL049938, Y11587,  
 AL080159, AL133665, E15569, AL122118, AL080137, AL137527, AL133606, AL110280, A93016,  
 A90832, U00763, I42402, U58996, L30117, AL137273, X65873, AL080158, X53587, AR038969,  
 AF118090, AJ242859, AF113691, AL080086, AL133104, AL133067, I09360, AR000496, U39656,  
 AL137538, U96683, Y14314, AF125949, AL133077, AL117432, AF081197, and AL122123.



HHBEG72	534	761150	1 - 550	15 - 564	H49710, and AL047185.
HEMHA53	535	728297	1 - 593	15 - 607	AA010194.
HEMGX57	536	872083	1 - 594	15 - 608	AA485012, AL138096, AA743445, AC004542, AC005358, AC007056, AL021368, U93037, AB014608, AC004526, AC005209, AC004953, AL121595, AL021397, AP000692, AC005253, AC005387, AC004148, AC005089, AC005519, U91322, AC005730, AC004774, AC005539, AC003006, AL023653, AC000090, AC005326, AC005800, AC006960, Z83820, AC002302, AC007487, Z15025, AC004237, Y14768, AP000505, AL031276, Z95152, AF129756, AP000962, Z82176, AC003101, AC003669, AL109628, AC006050, AL117354, Z99571, AL031597, AC005940, AC005486, AL035530, AP000500, and AP000555.
HEMGT27	537	851065	1 - 542	15 - 556	AA213376, AA713538, AA713930, AC002395, AC010170, AC006515, AC000355, D87675, and AC005057.
HEMGL56	538	767669	1 - 327	15 - 341	AA122010, W40525, AA173833, AA622443, AA334425, AF042166, AF043045, and AL137574.
HEMFN30	539	692818	1 - 1033	15 - 1047	AA099585, AA099645, AW271704, H88677, H88726, AF086025, and AF190639.
HEMFF16	540	576539	1 - 128	15 - 142	
HEMEF34	541	596812	1 - 1243	15 - 1257	AA653240, AA812058, AA569667, AA644090, AA714224, AL755214, AL037714, AL754567, AL525532, AL754105, AA488903, AW103406, AW023111, AA535216, AA524616, AA829036, AL380617, AA701080, AL279417, AL366555, AL912401, AL358712, T05118, AW328331, AA704393, AL754170, AA290878, AL079734, AL284543, AL635440, AA601278, AW237905, AL289277, AL669421, AA019973, AA013168, AL250552, AA169245, AL792521, AL120141, AW303098, AL251284, AL251034, AL251203, AW271977, AL043351, AL049709, AA654778, AL793172, AL793209, AA054085, AA640685, AL275982, AL369580, AL792499, H07953, AA559241, AA747757, AL254770, AL114557, AL043105, AL135377, AL452836, AL613389, F00564, AL612142, AA578621, T74524, AA420546, AA572813, AC005089, Z95115, AC005225, AC007227, AC005899, AC003982, AC002302, AC007193, AL121603, AP000692, AC006023, AC004882, AC002565, U91323, Z98742, AC007225, AL139054, AC005015, AC004408, AC004851, Z97054, AL034420, AC005081, AC002073, AL009181, AF053356, AL033521, AC005529, AL049576, AC005193, AL031255, AC002364, AL049776, AL080243, AL031311, AC002350, AC005520, AC004821, AL022318, AC002400, Z84469, AC004813, U91321, AL022316, AP000193, AC004491, AC005280, AC015853, Z93023, AC005800, AC005332, AC016830, AC002316, Z83844, AC007055, AC005875, AC002115, AC002477, AL049856, AC002425, AC005207, AP000117, AC009247, AC006449, AC005519, AC005082, AC006530, AC003007, AL050318, AP000354, AC006285, AC000134, AC005071, U62317, AC005971, AD000092, AC004099, AF129756, AL049697, AC002544, U91326, AL021407, AC007036, AC004000, AL022326, AL132777, AL117694, AF196779, AL078581, AF045555, AC003684, AL035588, AC006538, AL133246, AL021397, AL133448, AC004771, AC004231, AP000689, AC006121, Z85986, AL022165, AC004596, AC010205, AC003036, AC005412, AC004148, AC009248, AC004883, U82828, AC005261, AC007051, AC007308, AC005011, AC006942, AL024498, L78833, AL031848, AL050321, AL049694, AC005409, AC016025, AC004019, AC002312, AC004967, AL022320, AC006312, AC005940, AC006241, Z84466, AL251973,

HEMA03	542	921922	1 - 403	15 - 417	AC007917, AL020997, AL023807, AC005632, AL031432, AC016831, AC002492, AC004922, AL096701, AC005399, Y18000, AC002347, AC006480, AC002996, AC007934, AC002511, AP000556, AC007406, AC006139, AL121825, AC005037, Z83838, AC003665, AL031289, AP000501, AC005291, AP000557, AC006430, AC007919, U95740, AC005823, AC007388, AC002470, AC004098, AF038458, Z93017, AL031291, AC007201, AC005231, AC004797, AC005041, AC005480, AC004508, AL031283, AL021391, AL008582, AL035400, AL121852, AL035587, AL050332, AC005914, AC007011, AC004750, AL021154, Z93096, AC007371, L44140, AP000552, AF003626, AP000347, AC002430, AP000248, AL049569, AL031670, AL008725, AC005837, AC005592, AC004876, AL122020, AP000704, Z83826, AC006441, AC005088, AP000114, AP000046, AC004383, AL121652, AL034555, AL049839, AC005488, AL021366, AC000052, AP000355, AP000514, AL133163, AP000247, AC004933, AL035684, AL109758, AL135744, AC005274, AC004815, AL035249, AC007536, AC007384, AL109827, AC000025, AC005695, AF109907, AL031588, AC005696, AL034379, U47924, AL034549, AC005726, AL121655, AL049712, AC006536, AC005484, AL049780, AC006130, AB023049, AC003037, AC007687, AC005736, AC004854, AC006026, AC006014, and Z99716.
HEMDX96	543	935963	1 - 833	15 - 847	W90730.
HEMCV44	544	574321	1 - 418	15 - 432	AA463972, AA464108, A1080269, AA528133, W04732, AA912293, AA903826, W79633, AA557293, A1912574, A1906793, W25281, R47804, A1129889, AA365267, A1081962, W24623, AA404469, W19157, A1189367, A1159933, A1273685, AA301944, N70568, A1291841, AA911261, A1350318, A1541205, A1557426, A1541056, and AF111170.
HEMCK53	545	728424	1 - 495	15 - 509	R10409, and R09783.
HEMC159	546	739551	1 - 407	15 - 421	W03547, and AC002117.
HEMC141	547	712614	1 - 938	15 - 952	A1797078, AA129381, A1701463, A1808119, AA129318, A1309212, AA970643, N47001, A1263617, AA633720, and AB028993.
HEMCC38	548	707453	1 - 372	15 - 386	H61491, and AA699633.
HEMBU26	549	684928	1 - 305	15 - 319	R32090.
HEMBT61	550	939957	1 - 449	15 - 463	N86549, AW369713, and AB002301.
HEMAL61	551	851106	1 - 520	15 - 534	T85402.
HEMAA63	552	745498	1 - 320	15 - 334	AA058694, AA015929, AA012840, AA054133, and AC007279.
HELU74	553	765696	1 - 516	15 - 530	A1567076, AW419262, AW303196, AW274349, A1281881, AA720702, AW327868, A1270117, AA533725, AW301350, AW193265, AA649642, AA446657, AA747276, A1963720, A1350211, AL046409, AW193432, A1613280, A1085719, A1471481, AA847499, A1198376, AA581903, AA973803, AA828704, A1619997, A1688846, AA649705, A1653886, AA525824, A1590958, A1610920, A1375710, W79504, AA623002, A1623696, AW275432, A1801600, AW072923, A1590689, A1061334, R77905, AW406162, AA652057, AA491814, F36273, AA357937, AA350859, AA714453, A1431303.
HELHD20	554	668881	1 - 421	15 - 435	

					AA494163, AA493708, AA715355, F25199, AA773886, AI133164, AW276817, AA825357, AW265170, T15977, AI783494, AA469451, AA502866, AI143242, AA535661, T07451, AI754658, AW249224, AI305766, AW338417, AW088846, AW410400, AW057877, AA177061, AI761471, F19012, AL042420, AI334435, AC005971, AF077058, AL096701, Z99716, AL049766, AC004931, AL031286, D83989, AC004033, AL034549, AC004955, AC004808, AC005969, AC002302, AL023799, AL031542, X55926, X54181, X54178, I51997, AL096703, AC005399, AL109654, U18394, Z86061, AC007537, U18391, U18392, X55925, U57005, AC007386, X54180, U57009, X54179, AL031668, AC004448, X54175, U18393, AC005730, U67801, AC005690, X55931, U57008, AC005722, AL035687, AL022318, AC000085, AC006324, AC005846, AC005291, AC007283, AC002452, X54176, AP000362, AC005553, U57007, AF196969, AC006210, AC002542, U18390, AC003104, AL133396, AC012384, AC004526, U18396, AC007450, AC002564, X55932, AF117829, AC006996, AC004453, AC005951, AC004690, AC006581, AC006367, AC007388, Z69714, AC005881, U67825, X54177, AL121871, AL034351, AB020862, AC004673, AC004745, AC006328, AC004552, Z94801, AC005019, AL035695, AL031073, AL049833, AL034547, AC004072, U02532, AC007934, AC010197, AF015153, AC005919, AC004948, AL022311, AL109653, AC007546, AC004227, U18400, AL050308, AL049539, AP000561, AC009891, AC007385, AL109963, AP000548, AC007225, AL133353, S77605, AL031643, AF015151, AF222684, AC004984, U73465, Z81364, AL121658, AF064866, Z84474, AL031577, AC005409, AJ011930, AL049835, AC011311, AC004945, AC009363, AC002038, Z92545, AL008721, AC004909, AC002301, AC006043, AC004492, Z93928, AC005342, AL133241, AC006241, AC007676, AC007344, AC005820, AC007285, AC006374, AC005610, AC004992, AC004796, AC004890, AC007436, AL049591, AC005387, AC005082, AC004626, AL031848, AL031681, AL031774, AF015157, AC008064, AC006057, Z82976, AL023574, AP000688, AC007358, AC005839, AF188024, AL133289, AC007731, AL136018, AC007227, AC005500, AC004782, S70707, AC005703, AC005183, AL031229, AC006333, Z68289, AC000094, AC004531, Z82201, AP000557, AC004021, AC006571, AC006478, AL109628, AF015149, AL031390, AL049647, AC007370, AC004993, Y18000, AL121578, AC004386, AL021069, AC004629, Z84721, AL008723, AL109801, U57004, AL049869, AC010072, AC004916, AC004864, AC007207, AF165926, AL035079, AC007262, AP000705, AL049759, AC005033, AC011604, AL031287, and AF015148.
HELHC59	555	769404	1 - 505	15 - 519	N72137, N24885, N25625, AI128843, AI126506, AI160533, AI200037, AW450603, N35103, N26904, AW020616, AI695490, AI802647, A729493, AA605122, AW024969, AA806507, AA743134, AI217597, AI312534, AI219599, N25653, AA729125, AA568681, H86995, N26781, AI829099, AA648514, AA568193, AI032141, R73137, R26304, AW298649, AA768761, N20053, AW236299, AW453038, AW452862, AI141901, AA988539, AI361669, AI674252, AI039557, AI299683, AI984739, AI452444, W52017, AA543074, N34223, N28403, T25828, N36303, AI362330, AA296449, AW074182, AI805445, W52651, and AA653691.
HELGY42	556	713019	1 - 640	15 - 654	H23853, and H46507.
HELGY02	557	948302	1 - 510	15 - 524	AI939550.

HELGW31	558	610003	1 - 1645	15 - 1659	C14389, D80268, AW177440, AW177501, AW177511, AW352117, D81026, D59502, AI905856, AW178893, T03269, C14014, AA305578, AW179328, AW366296, AW360811, AW375405, AA514188, D58283, D59859, D80022, C14331, D80166, D80195, D80193, D59927, D59467, D51423, D59619, D80210, D51799, D80391, D80164, D59275, D80240, D80253, D80043, D59787, D80227, AW378532, D81030, D80212, D80196, D80188, D80219, AW176467, C15076, D80269, D80038, D59610, D57483, D80366, AA305409, C14429, D51022, D50979, D50995, D59889, AW178762, D80024, AW377671, D80378, AW178775, AW360844, AW360817, D80241, D51060, AW352158, AW375406, D80248, AW378534, AW179332, AW377672, AW179023, AW178905, D80134, D80045, D80132, D51097, AW352170, D58253, AW352171, D80522, AW377676, AW177731, AW178907, AW179019, AW179024, D80251, AA514186, C75259, D80133, AW178906, AW177505, AW179020, AW178909, AW177456, AW179329, AW178980, AW177733, AW378528, AW178908, AW178754, AW179018, AW179004, AW178914, AW178911, AW367967, AW352174, D80302, AW178774, AW177723, D80439, D80247, T48593, AI535850, AW178983, D51103, AW367950, C14975, AW178986, D45260, AI525913, Y17188, X82626, A84916, A67220, D89785, A62300, A62298, A78862, D34614, D26022, D88547, AJ132110, AR018138, X67155, A25909, AF058696, AR008278, Y12724, AR025207, AB028859, AB012117, A94995, A85396, D88507, AR066482, A44171, A85477, AR008443, I19525, A86792, I18367, X93549, I50126, I50132, I50128, I50133, AR066488, A82595, AR066490, AR016514, D50010, D13509, AR060138, A45456, A26615, AR052274, Y09669, AR060385, AB002449, AR066487, A43192, A43190, AR038669, A30438, AR008408, U79457, AF135125, AR060133, and AR008382.
HELGV36	559	597120	1 - 359	15 - 373	H79363, R79034, Z26324, and AC007199.
HELQ55	560	732223	1 - 346	15 - 360	AA410214, W24694, and AI220186.
HELK36	561	925698	1 - 776	15 - 790	
HELGG21	562	671071	1 - 463	15 - 477	W01622.
HELGD47	563	851143	1 - 442	15 - 456	AA496758.
HELQ55	564	732224	1 - 449	15 - 463	
HELFN75	565	658681	1 - 621	15 - 635	R59270, F11339, and AB020864.
HELFI35	566	506277	1 - 307	15 - 321	
HELFA38	567	851146	1 - 391	15 - 405	AC005736, AC005412, AC004491, AP000689, AB003151, AC004796, AC004019, AC004967, AC004913, Z83822, AF185589, AL121655, AC006120, AL031594, AF047825, and AC005291.
HELEZ81	568	571340	1 - 322	15 - 336	AA355367, AA452887, AA581247, AA501794, AW303142, AI253987, AI077941, AA904275, AW302081, AA557790, AL079734, AA666295, AI802804, AI249688, AW151247, AA642809, AL049712, I34294, AL031258, AC005839, AC005512, AC005520, AP000694, AC006449, AC009516, AL021707, AC005944, AL035405, AC004148, AC004253, Z98051, AL050332, AC002544, Z93023, Z73420, AC006276, AB023049, AC004033, AC005261, AL080242, AF109907, AC005399, AP000305, Z85986, AC007216, AL096712, L47234, AL049760, AL022326, AC005777, AL133246, AL031686, AP000557, AC004812, AC002395, AC005578, AL031680, AC005920, AC002470, AP000152,

Variable	Mean	SD	Min	Max
Age	34.5	10.5	20	55
Gender	0.5	0.5	0	1
Marital status	0.5	0.5	0	1
Education	12.5	1.5	10	15
Income	15.5	5.5	10	25
Health status	0.5	0.5	0	1
Smoking status	0.5	0.5	0	1
Alcohol consumption	0.5	0.5	0	1
Exercise frequency	0.5	0.5	0	1
Stress level	0.5	0.5	0	1
Depression score	0.5	0.5	0	1
Life satisfaction	0.5	0.5	0	1
Quality of life	0.5	0.5	0	1
Healthcare utilization	0.5	0.5	0	1
Health insurance status	0.5	0.5	0	1
Healthcare access	0.5	0.5	0	1
Healthcare cost	0.5	0.5	0	1
Healthcare quality	0.5	0.5	0	1
Healthcare satisfaction	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.5	0	1
Healthcare cost (continued)	0.5	0.5	0	1
Healthcare quality (continued)	0.5	0.5	0	1
Healthcare satisfaction (continued)	0.5	0.5	0	1
Healthcare utilization (continued)	0.5	0.5	0	1
Health insurance status (continued)	0.5	0.5	0	1
Healthcare access (continued)	0.5	0.		

HELET68	569	800029	1 - 446	15 - 460	AF053356, AC006241, AC006509, AF165926, AC004876, AC007664, AC005695, AC002314, AC004655, AC005529, AL031431, AP000502, Z93017, Z83844, AC005900, AC005288, AC005256, AJ003147, AC004796, AF196779, AC002059, AC002133, AC004962, AL109827, AC002054, AC004991, AC005971, AC004491, AC004522, AC004099, AP000045, AC002316, AC009247, AC005800, AC016027, AL031281, AL031005, AL049570, AP000512, AP000047, AL035587, AL109623, AC006014, AC005255, AL034420, AC007201, AC002115, U91326, Z97630, AC004678, AL008726, AC016830, U47924, AL049869, AL109952, AC007546, AL031662, AC002126, AC004804, AC006236, AC007226, AC006538, AC004686, AL022315, AC007387, AL132712, AC002107, AF111168, AC005476, AL034423, AC004883, AC006480, U95742, AL122020, AL021154, AL021977, AC003101, AL031311, AP000252, AL022320, AL050318, AC004760, AC004966, AP000691, AP000563, AP000692, AF001548, AC005015, AC005581, AC004832, Z82176, AP000300, Z84480, AC004878, AL031735, AL133448, AC008040, AC004531, AF001549, AC005057, AF207550, AC006285, AB001523, AL050341, AC005940, AC004449, AC007565, AC007308, AC005231, AC005082, AL031427, AC005089, Z98941, AL049856, AF030453, AC005071, AC005048, AC008044, AC004851, AL121652, AC002300, AC005829, Z95331, and Z68870.
HELET68	569	800029	1 - 446	15 - 460	R96611, AA019463, AA018536, AC006001, AJ250042, and AJ001119.
HELEO42	570	579016	1 - 556	15 - 570	H00519, AI671775, AA689296, AW082267, AW340166, AI671779, AI298659, AA251240, AI189305, AI695034, AW085290, AI290925, AA977259, AA806531, AA836353, AA809326, AW086155, AA811618, AA731135, AF027156, and Z57575.
HELEH76	571	506674	1 - 418	15 - 432	AI813783, AW028943, AI872000, AA429481, AI146266, AW249720, AA496309, H71659, F31867, AW304536, AA583579, AW194077, W24312, AA758131, AI207476, AA297698, AA531596, AW003612, AI918419, AA654874, AA767884, AI358712, AA729512, AI299882, R74433, AI198718, AA745302, AA058312, AA558404, AA904211, F31951, AI866971, AA487300, C14330, T39217, AA225358, AA584183, AI310464, AA225890, AA341865, AA299422, AW381847, AW381904, F16409, AA713735, R93919, F26072, AA343810, AA937809, AA513884, T54464, AI365625, AA729253, AA808982, AI653999, AA516214, AI309979, AA525293, T39225, AW023111, AI915081, AW237905, AA584493, AI873627, AA179663, AW003595, AA634926, AA018777, AA508036, AA583568, AI198948, AI565084, AL037714, H67064, AA525407, AA644090, H15295, AI053784, AA654482, AA515728, W52705, AA744423, AA320105, AL045476, AA298573, AI282629, AL045423, AA302982, F33505, AW162227, AW102811, R52008, AW275679, T05118, AW161459, J03764, AC004876, AC007225, AL133448, Z98884, AC003982, AL035405, U89337, AL049834, AL031657, U07561, AC004848, Z97056, AL022336, AC007011, AC005914, AC005244, AC004985, AL049692, AF039954, AC007686, M87889, AB018249, AC006050, AC004382, AF207550, AC000378, AC007546, AC007371, AL035659, AC005328, AC002504, AC006409, AC005030, AF041001, L78833, AC005480, AC006257, AC005340, AC005953, AL109801, AL031258, AF111168, Z99128, AP000032, AP000103, AC006312, AL031662, AL122020, AL049569, AC005484, AC002302, AC005206, U02052, AP000356,
HELEE83	572	577206	1 - 235	15 - 249	

					AC006455, AL133245, AP000512, AC005668, AL034549, AP000270, AC004106, AC005667, AC000120, AL121652, AL031053, AC006443, AC006505, AP000045, AP000113, AC004851, X81870, D87675, AC004675, AC004988, AC002350, AF112441, AC002126, AC002492, AC008116, AC002404, AC006012, AC007298, AP000555, Z99755, Z86090, AF088219, AC003012, AC005786, AL032821, AP000008, AC006511, AL049849, AL049712, AC004129, AC005529, AL096791, AP000704, AC005621, AC005102, AC004150, AC004491, AC002476, AP000553, AC006965, AC006277, L35532, AC005972, AC006237, AC007308, AC003667, AL021578, AC006013, AC005011, AC004125, AC004024, AC005237, U78027, AL031767, AL022326, U11095, AP000343, AL133243, AL034423, U95741, AC005325, AF196970, AL049539, AC005632, AL035422, AC002425, AC004662, AC004066, AC004736, AC008064, AP000109, AP000041, AL121655, AC006084, AC004216, AC005015, AC005351, AF047825, AC005544, AC005086, AC006372, AC005523, AF121782, AC005921, AL022327, AL035412, AC007283, AL133233, AC005519, AP000690, AC005082, AL133035, AC006365, Z82206, Z75746, AC004801, AC005049, AC002381, I74786, AP000501, AC006261, AC007192, AC005291, AC002511, Z97832, AC002117, AC016025, AL031283, AC005527, AC007040, AC005231, AC006509, AL035699, AL022163, AC005363, AL049911, AC006368, AC002996, AF001550, AC002312, AC002195, AC006450, D00591, AC005410, AF064105, AC005993, AL023575, AC000024, AL121748, AC007685, AC009247, AC011311, AC007226, AC002300, AC000025, AL021707, AL022721, AC002112, AC005243, AF109907, AL049557, AC005261, AC007199, AC005500, AC006449, AC006205, AC004900, AL022165, AC005088, AC004760, AC004973, AC005393, U91321, AC005952, AC003038, AL031985, Z82217, U47654, AC004771, AL133485, Z82244, AL033527, AL049759, U95742, AC007216, AL050318, AC005778, Z93930, and AC005793.
HELDT63	573	744864	1 - 697	15 - 711	AL291317, AA501614, AA937686, AA747472, AA614010, R63317, AW062724, AL053445, AL433247, AI832000, R79794, AA384039, T48872, T17269, AA434044, AL134369, AL673731, AL036283, AI538812, N62650, AA434172, AI859834, AA443727, AA122223, AI370475, AI079129, T32050, AA533408, AA227837, AI978792, AI347810, AA508873, AA857381, H29019, AI654529, H74314, AI872503, F23255, T03900, AW272763, AI053786, AI056415, AA223249, AL036805, AW193493, AI299050, AI049831, AA809189, AL135698, AA601680, AA326410, AW028908, AL120976, AI979005, AA557486, AA515224, AA715004, AA553448, AL042113, AA584489, AA046466, AA720732, AA524832, AL045077, N67816, AA342681, AW419118, AI859946, AI283090, AA507169, F17700, AW337454, AI890385, AA577732, AI168205, AA564145, AI360514, AA528480, AI343143, AA121919, AI524540, AI343113, AI440324, AI457597, AW439480, AA437402, AA485485, AI610201, AI803827, AI65213, AI696901, AL033504, AL117337, AL021407, U51281, AC005701, AF205588, AC005592, AC001228, AP000193, AP000117, AL133355, AP000050, AC002119, AF107885, AC002524, AC008372, AC005212, Z49237, AB023051, AC006965, AC008928, AF091512, AL117328, AC006571, AC006257, AC006285, AC003030, AL031733, AP000512, AP000695, AC020663, AC004791, AF051976, AP000131, AP000209, AP000248, AC004472, AC007324, AC008101, AL049780, AC007055, U07000, AC005494, AC002057, AC005914, Z84487, AC009263, AC000004,

				AL050306, AC004841, AC005378, Z81002, AF045555, AL121578, AC004878, Z85986, AC006317, Z98051, AL022353, AC005523, AC005838, AP000023, AC005081, AC006450, AC006948, Z99716, AF148461, AC016025, AC006077, AC007014, AC008079, AC005755, AC005210, AC003667, AP000509, AL109758, AL096773, AF055278, AC006211, AC005911, AB026898, AL031296, AC006372, AC005512, AL008719, AC004858, AC005192, AF109907, AL080317, L32588, AF178081, AF008243, AP000018, AP000160, AC003043, AC002367, L78833, AC011604, AC004659, AC004967, AL022401, AL022329, AC005740, AC005365, AC009501, AC007934, AC003015, AC006143, AP000302, AC004796, AC005593, Z68280, AC006141, AC005823, AP000494, AC004024, AC008394, AC004974, AC005753, AC005768, Z99129, AC006166, AL022398, AC002511, AP000504, AC006048, AL078593, AC003104, Z95116, AC008149, AC004973, AL034417, AC002980, AC006254, AC005280, AL035410, AP000046, AP000114, Z93241, AC004765, AL031673, AC002316, AC004813, AL139054, AL021706, AL133448, AC006947, AC005599, AL031255, AC003682, AC004263, AJ003147, AP000701, AL096710, AL031230, AC005951, AL021939, AC006130, Z69364, AC002301, AC005037, AC004531, AC005757, AC002539, AC004876, AC004562, Z85987, AC005759, AL031584, AC004002, L81690, AC004797, AC007795, AP000115, AL009181, AC005726, AC005832, D84394, AL031278, U85195, AC009294, AC005071, AL035089, AC005971, U80017, AC004870, AE000658, AC008154, Z84469, U95740, AC002045, AC005399, AC018767, Z73963, AC006132, L05367, AC004552, AC004526, AC005225, AL035587, AC006552, AC002432, AF111168, AL008715, AL022238, AF129756, AC007676, AL034350, AL031121, Z95331, X96421, AC006296, AP000313, AL121603, AC005519, AC005243, AC006160, AF200465, AP000563, AF037062, AC005197, AP001043, Z93341, AL050333, AL022396, AL133304, AC006442, AC005772, AL035071, AC007637, and AC002418.	
HELDL15	574	660557	1 - 533	15 - 547	AI989707, AI668636, AI668638, N80930, AI916749, N89573, AA687664, N89599, AI734212, AI734241, and AC005180.
HELDL08	575	959919	1 - 363	15 - 377	AA219475.
HELDK22	576	567310	1 - 994	15 - 1008	AI801505, AA601218, AI679294, AI679871, AI889579, AA598545, AA604843, AW105729, AL041895, AI050007, AW028950, AA604865, AI708005, AF129756, AC004805, AP000045, AF111168, U95742, AC004491, AC007216, AC005940, AC007240, AC005837, AL035681, AL035072, AP000557, AL096791, AL022336, L44140, AC002059, AC007182, AL121658, AC007226, AP000688, AC004099, AC006211, AP000555, AC002039, AC002425, AL023284, U91318, AP000553, AC004098, AC004815, L78810, AC002996, AC002301, AC005920, AC005531, AC005288, AC006241, AP000113, AC004526, AC004966, AC004841, AC004253, AC000052, AL080243, Z94056, AC005412, AC005057, AL049569, AC005377, AC006538, AC006312, AC005803, AC007055, AC005089, AC003029, AP000689, U62293, AC007666, AC005529, AL049776, AC005736, AC000134, U80017, AC004883, AC004531, AL023553, AL022165, Z93241, AC004983, AP000065, AL031295, AC005839, AJ010770, AP000010, AP000952, AC004686, AC004967, AL031670, AL022238, Z97054, AC009516, U47924, AL109628, AC004149, AC002314, AC004922, AC005255, AF030876, AC008372, Z85986, AL021397, AC007021, AL021546, AC002544, AF053356, AL049697, AC005562, AC005004, AC004084,

HELDH71	577	740198	1 - 629	15 - 643	AF038458, AL031311, AC004933, AP000505, AC002350, AC006121, AC005519, AP001053, AC004216, L48038, Z97056, AC002477, AC003684, AL022320, AC005031, AC004895, U52111, AC005015, AC006480, AC020663, AC006530, AC007011, U95743, AC005231, AC005899, Z95115, AC006449, AC005387, AL049830, AC004132, AC002492, AC005921, AC004953, AC004659, AC004887, AC004973, AL049869, AL031584, AC005822, AL133245, AC005488, AF109907, AC006064, AC005520, AL121603, AC005722, AC007676, AL049795, AF107885, AL020997, AC006285, AC005071, AC006014, AC002470, AL031291, AC002077, AL096701, AL049843, AC003101, AC008055, AF196969, U91321, AL034379, AC005049, AC006001, AC000026, AC006115, AL121653, U91326, AL035684, AF134726, AC005102, AC002395, AC003035, AL109984, AC007227, AC005274, AF067844, AC007686, AC004382, AF196972, AC005300, AC005082, AP000501, AL121825, AC010206, AL050318, AJ246003, AP000152, AC004814, AC000003, AC008040, AP000556, AC003041, AC007637, AL031283, AC002476, AP000247, AC005081, AC005332, AC005696, AC006965, Z98884, Z95152, AC004167, AC007546, AF045555, AL021393, AC000353, AC005740, AC016025, AC006947, AC016830, AC007277, AP000558, AF001552, AC016027, AC004019, AC005527, AL031985, AF205588, AC004752, AL078463, AL022316, Z82206, AP000300, AC004812, AC007376, AL035086, AC005037, AC005484, Z83844, AC009509, AC002375, Z93023, AL024498, and AL035450.
HELDG91	578	790371	1 - 757	15 - 771	H40372, AA190779, AF192065, AA315314, AF192074, H15516, R14652, R60461, AA147113, AA173721, AW028676, H11571, H65653, T85096, R96576, AA337744, R02157, AA191497, R73942, R09150, W24899, AC004537, and AF161419.
HELW51	579	531073	1 - 248	15 - 262	AI808351, N32277, AA886875, AA256444, W81274, AI056648, W81237, AI168722, AA775321, H97616, AA256318, H26750, H26847, H97380, AI241162, and N44565.
HEL30	580	691024	1 - 359	15 - 373	AR025382.
HEL36	581	655045	1 - 498	15 - 512	H29208, AA595183, and AF102777.
HELB11	582	967661	1 - 422	15 - 436	AA037425, AI741209, R81422, and AW275774.
HELB83	583	781412	1 - 531	15 - 545	AA136444, H96161, H61998, W03286, and AC010209.
HEL48	584	864515	1 - 951	15 - 965	R13326, and R20026.
					W19935, AA451818, AW402568, H96614, N25240, AW367995, Z44387, R20284, H06089, AA034188, AW247735, AI903997, R15130, F11576, R14728, AA317939, N99694, AW410153, AW409593, F05727, W16878, W21473, R15845, H46506, H20375, H50149, R73366, AA336250, AW383353, N33873, and AA379990.
HELAW26	585	684925	1 - 528	15 - 542	H49721, H78006, and R05888.
HELAQ36	586	707420	1 - 197	15 - 211	H59264.
HELAM32	587	699661	1 - 328	15 - 342	T79826, AW136576, W44672, AI863582, N77892, H62231, AW207812, AA722922, AI374730, AI632306, AW263176, AI770020, AI817074, N74527, AI948698, AA355234, AA676826, AI636019, AA063431, D25824, C21355, AW235414, AW451382, AW377510, AC016830, AC008101, AC008079, AC016027, and AC007324.



HELAH32	588	699665	1 - 490	15 - 504	W46180, AA284181, and AF086189.
HCMSY80	589	526182	1 - 264	15 - 278	
HCMSQ63	590	745584	1 - 450	15 - 464	AI679298, AI679875, AI925636, AI719454, AA235054, AI333087, AI095570, AA877427, and AL137734.
HCMK41	591	940260	1 - 597	15 - 611	AA719296, and AA758153.
HAHSB27	592	501010	1 - 405	15 - 419	
HAHFS80	593	954432	1 - 550	15 - 564	F22045, F31427, F00312, AA383663, and AA247490.
HAHFE11	594	965293	1 - 402	15 - 416	AA888984, and AL008718.
HAHEP68	595	738501	1 - 473	15 - 487	W95964, AA279579, AA357601, H05847, AI026709, AA065168, W95676, AW451661, AW382980, AA065169, AI739083, AI239672, AA729486, AA768730, AW003068, and AC006504.
HAHCU22	596	848831	1 - 199	15 - 213	N56073, AW369804, R45746, AA758653, and AC006205.
HAHCR15	597	810326	1 - 619	15 - 633	W60983, AW382330, AA400064, N83192, AA248090, N30985, AA401371, C01567, and AB014569.
HAHCL94	598	794044	1 - 387	15 - 401	T86209.
HAHBC03	599	923542	1 - 528	15 - 542	AA293827, and AA402475.
HAHAD95	600	865104	1 - 323	15 - 337	AA311201.
HAFBG30	601	693363	1 - 459	15 - 473	AA156638, AW340037, AA576325, AI090498, AI652388, AA304579, AI280372, F35830, AI623386, AI652584, F26635, AI682162, and AC005740.
HAFAY37	602	928705	1 - 516	15 - 530	AI625551, AI018611, AI004171, AW270040, AI359447, N64334, AI928764, R91517, AW205692, AI093803, AW291368, AI056157, AA304165, AW388381, AA251845, AW193685, C00746, and AI768273.
HAFAJ63	603	845452	1 - 546	15 - 560	R64458, AI657485, H66475, AA203500, R69944, AA303679, AA393390, H92410, AI908514, AW406064, AA398719, H85493, AA152049, H59673, and R37909.
HAECA04	604	932993	1 - 309	15 - 323	L44310, AA302370, AC006084, and AC004820.
HAEM82	605	781539	1 - 558	15 - 572	AA114163, AA573134, AW139049, AW204553, and AI909752.

**TABLE 4**

Code	Description	Tissue	Organ	Cell Line	Disease	Vector
AR022	a_Heart	a_Heart				
AR023	a_Liver	a_Liver				
AR024	a_mammary gland	a_mammary gland				
AR025	a_Prostate	a_Prostate				
AR026	a_small intestine	a_small intestine				
AR027	a_Stomach	a_Stomach				
AR028	Blood B cells	Blood B cells				
AR029	Blood B cells activated	Blood B cells activated				
AR030	Blood B cells resting	Blood B cells resting				
AR031	Blood T cells activated	Blood T cells activated				
AR032	Blood T cells resting	Blood T cells resting				
AR033	brain	brain				
AR034	breast	breast				
AR035	breast cancer	breast cancer				
AR036	Cell Line CAOV3	Cell Line CAOV3				
AR037	cell line PA-1	cell line PA-1				
AR038	cell line transformed	cell line transformed				
AR039	colon	colon				
AR040	colon (9808co65R)	colon (9808co65R)				
AR041	colon (9809co15)	colon (9809co15)				
AR042	colon cancer	colon cancer				
AR043	colon cancer (9808co64R)	colon cancer (9808co64R)				
AR044	colon cancer 9809co14	colon cancer 9809co14				
AR045	corn clone 5	corn clone 5				
AR046	corn clone 6	corn clone 6				
AR047	corn clone2	corn clone2				
AR048	corn clone3	corn clone3				
AR049	Corn Clone4	Corn Clone4				
AR050	Donor II B Cells 24hrs	Donor II B Cells 24hrs				
AR051	Donor II B Cells 72hrs	Donor II B Cells 72hrs				
AR052	Donor II B-Cells 24 hrs.	Donor II B-Cells 24 hrs.				
AR053	Donor II B-Cells 72hrs	Donor II B-Cells 72hrs				
AR054	Donor II Resting B Cells	Donor II Resting B Cells				
AR055	Heart	Heart				
AR056	Human Lung (clonotech)	Human Lung (clonotech)				
AR057	Human Mammary (clonotech)	Human Mammary (clonotech)				
AR058	Human Thymus	Human Thymus				

	(clonotech)	(clonotech)				
AR059	Jurkat (unstimulated)	Jurkat (unstimulated)				
AR060	Kidney	Kidney				
AR061	Liver	Liver				
AR062	Liver (Clontech)	Liver (Clontech)				
AR063	Lymphocytes chronic lymphocytic leukaemia	Lymphocytes chronic lymphocytic leukaemia				
AR064	Lymphocytes diffuse large B cell lymphoma	Lymphocytes diffuse large B cell lymphoma				
AR065	Lymphocytes follicular lymphoma	Lymphocytes follicular lymphoma				
AR066	normal breast	normal breast				
AR067	Normal Ovarian (4004901)	Normal Ovarian (4004901)				
AR068	Normal Ovary 9508G045	Normal Ovary 9508G045				
AR069	Normal Ovary 9701G208	Normal Ovary 9701G208				
AR070	Normal Ovary 9806G005	Normal Ovary 9806G005				
AR071	Ovarian Cancer	Ovarian Cancer				
AR072	Ovarian Cancer (9702G001)	Ovarian Cancer (9702G001)				
AR073	Ovarian Cancer (9707G029)	Ovarian Cancer (9707G029)				
AR074	Ovarian Cancer (9804G011)	Ovarian Cancer (9804G011)				
AR075	Ovarian Cancer (9806G019)	Ovarian Cancer (9806G019)				
AR076	Ovarian Cancer (9807G017)	Ovarian Cancer (9807G017)				
AR077	Ovarian Cancer (9809G001)	Ovarian Cancer (9809G001)				
AR078	ovarian cancer 15799	ovarian cancer 15799				
AR079	Ovarian Cancer 17717AID	Ovarian Cancer 17717AID				
AR080	Ovarian Cancer 4004664B1	Ovarian Cancer 4004664B1				
AR081	Ovarian Cancer 4005315A1	Ovarian Cancer 4005315A1				
AR082	ovarian cancer 94127303	ovarian cancer 94127303				
AR083	Ovarian Cancer 96069304	Ovarian Cancer 96069304				
AR084	Ovarian Cancer 9707G029	Ovarian Cancer 9707G029				
AR085	Ovarian Cancer 9807G045	Ovarian Cancer 9807G045				
AR086	ovarian cancer 9809G001	ovarian cancer				

		9809G001				
AR087	Ovarian Cancer 9905C032RC	Ovarian Cancer 9905C032RC				
AR088	Ovarian cancer 9907 C00 3rd	Ovarian cancer 9907 C00 3rd				
AR089	Prostate	Prostate				
AR090	Prostate (clonotech)	Prostate (clonotech)				
AR091	prostate cancer	prostate cancer				
AR092	prostate cancer #15176	prostate cancer #15176				
AR093	prostate cancer #15509	prostate cancer #15509				
AR094	prostate cancer #15673	prostate cancer #15673				
AR095	Small Intestine (Clontech)	Small Intestine (Clontech)				
AR096	Spleen	Spleen				
AR097	Thymus T cells activated	Thymus T cells activated				
AR098	Thymus T cells resting	Thymus T cells resting				
AR099	Tonsil	Tonsil				
AR100	Tonsil germinal center centroblast	Tonsil germinal center centroblast				
AR101	Tonsil germinal center B cell	Tonsil germinal center B cell				
AR102	Tonsil lymph node	Tonsil lymph node				
AR103	Tonsil memory B cell	Tonsil memory B cell				
AR104	Whole Brain	Whole Brain				
AR105	Xenograft ES-2	Xenograft ES-2				
AR106	Xenograft SW626	Xenograft SW626				
H0002	Human Adult Heart	Human Adult Heart	Heart			Uni-ZAP XR
H0019	Human Fetal Heart	Human Fetal Heart	Heart			pBluescript
H0050	Human Fetal Heart	Human Fetal Heart	Heart			Uni-ZAP XR
H0097	Human Adult Heart, subtracted	Human Adult Heart	Heart			pBluescript
H0105	Human Fetal Heart, subtracted	Human Fetal Heart	Heart			pBluescript
H0173	Human Cardiomyopathy, RNA remake	Human Cardiomyopathy	Heart		disease	Uni-ZAP XR
H0196	Human Cardiomyopathy, subtracted	Human Cardiomyopathy	Heart			Uni-ZAP XR
H0230	Human Cardiomyopathy, diff exp	Human Cardiomyopathy	Heart		disease	Uni-ZAP XR
H0233	Human Fetal Heart, Differential (Adult- Specific)	Human Fetal Heart	Heart			pBluescript
H0242	Human Fetal Heart, Differential (Fetal- Specific)	Human Fetal Heart	Heart			pBluescript
H0266	Human Microvascular Endothelial Cells, fract. A	HMEC	Vein	Cell Line		Lambda ZAP II

H0267	Human Microvascular Endothelial Cells, fract. B	HMEC	Vein	Cell Line		Lambda ZAP II
H0268	Human Umbilical Vein Endothelial Cells, fract. A	HUVE Cells	Umbilical vein	Cell Line		Lambda ZAP II
H0373	Human Heart	Human Adult Heart	Heart			pCMV Sport 1
H0403	H. Umbilical Vein Endothelial Cells, IL4 induced	HUVE Cells	Umbilical vein	Cell Line		Uni-ZAP XR
H0412	Human umbilical vein endothelial cells, IL-4 induced	HUVE Cells	Umbilical vein	Cell Line		pSport1
H0413	Human Umbilical Vein Endothelial Cells, uninduced	HUVE Cells	Umbilical vein	Cell Line		pSport1
H0433	Human Umbilical Vein Endothelial cells, frac B, re-excision	HUVE Cells	Umbilical vein	Cell Line		pBluescript
H0437	H Umbilical Vein Endothelial Cells, frac A, re-excision	HUVE Cells	Umbilical vein	Cell Line		Lambda ZAP II
H0530	Human Dermal Endothelial Cells, untreated	Human Dermal Endothelial Cells; untreated				pSport1
H0531	Human Dermal Endothelial cells, treated	Human Dermal Endothelial cells, treated with VPF				pSport1
H0599	Human Adult Heart, re-excision	Human Adult Heart	Heart			Uni-ZAP XR
H0619	Fetal Heart	Human Fetal Heart	Heart			Uni-ZAP XR
H0645	Fetal Heart, re-excision	Human Fetal Heart	Heart			Uni-ZAP XR
S0005	Heart	Heart-left ventricle	Heart			pCDNA
S0045	Endothelial cells-control	Endothelial cell	endothelial cell-lung	Cell Line		Uni-ZAP XR
S0046	Endothelial-induced	Endothelial cell	endothelial cell-lung	Cell Line		Uni-ZAP XR
T0048	Human Aortic Endothelium	Human Aortic Endothelium				pBluescript SK-
T0049	Aorta endothelial cells + TNF-a	Aorta endothelial cells				pBluescript SK-
L0002	Atrium cDNA library Human heart					
L0060	Human thymus NSTH II					
L0070	Selected chromosome 21 cDNA library					
L0142	Human placenta cDNA (TFujiwara)	placenta				
L0157	Human fetal brain (TFujiwara)		brain			
L0362	Stratagene ovarian cancer (#937219)					Bluescript SK-
L0364	NCI_CGAP_GC5	germ cell tumor				Bluescript SK-
L0367	NCI_CGAP_Sch1	Schwannoma tumor				Bluescript SK-

L0375	NCI_CGAP_Kid6	kidney tumor	kidney			Bluescript SK-
L0378	NCI_CGAP_Lu1	lung tumor	lung			Bluescript SK-
L0438	normalized infant brain cDNA	total brain	brain			lafmid BA
L0439	Soares infant brain INIB		whole brain			Lafmid BA
L0455	Human retina cDNA randomly primed sublibrary	retina	eye			lambda gt10
L0459	Adult heart, Clontech					Lambda gt11
L0471	Human fetal heart, Lambda ZAP Express					Lambda ZAP Express
L0485	STRATAGENE Human skeletal muscle cDNA library, cat. #936215.	skeletal muscle	leg muscle			Lambda ZAPII
L0529	NCI_CGAP_Pr6	prostate				pAMP10
L0530	NCI_CGAP_Pr8	prostate				pAMP10
L0532	NCI_CGAP_Thy1	thyroid				pAMP10
L0534	Chromosome 7 Fetal Brain cDNA Library	brain	brain			pAMP10
L0547	NCI_CGAP_Pr16	tumor	prostate			pAMP10
L0581	Stratagene liver (#937224)		liver			pBluescript SK
L0588	Stratagene endothelial cell 937223					pBluescript SK-
L0589	Stratagene fetal retina 937202					pBluescript SK-
L0591	Stratagene HeLa cell s3 937216					pBluescript SK-
L0592	Stratagene hNT neuron (#937233)					pBluescript SK-
L0593	Stratagene neuroepithelium (#937231)					pBluescript SK-
L0595	Stratagene NT2 neuronal precursor 937230	neuroepithelial cells	brain			pBluescript SK-
L0596	Stratagene colon (#937204)		colon			pBluescript SK-
L0598	Morton Fetal Cochlea	cochlea	ear			pBluescript SK-
L0599	Stratagene lung (#937210)		lung			pBluescript SK-
L0601	Stratagene pancreas (#937208)		pancreas			pBluescript SK-
L0602	Pancreatic Islet	pancreatic islet	pancreas			pBluescript SK-
L0603	Stratagene placenta (#937225)		placenta			pBluescript SK-
L0604	Stratagene muscle 937209	muscle	skeletal muscle			pBluescript SK-
L0605	Stratagene fetal spleen (#937205)	fetal spleen	spleen			pBluescript SK-
L0608	Stratagene lung carcinoma 937218	lung carcinoma	lung	NCI-H69		pBluescript SK-
L0617	Chromosome 22 exon					pBluescriptIIKS +
L0622	HM1					pcDNAII (Invitrogen)

L0623	HM3	pectoral muscle (after mastectomy)				pcDNAII (Invitrogen)
L0637	NCI_CGAP_Brn53	three pooled meningiomas	brain			pCMV-SPORT6
L0638	NCI_CGAP_Brn35	tumor, 5 pooled (see description)	brain			pCMV-SPORT6
L0646	NCI_CGAP_Co14	moderately- differentiated adenocarcinoma	colon			pCMV-SPORT6
L0651	NCI_CGAP_Kid8	renal cell tumor	kidney			pCMV-SPORT6
L0657	NCI_CGAP_Ov23	tumor, 5 pooled (see description)	ovary			pCMV-SPORT6
L0659	NCI_CGAP_Pan1	adenocarcinoma	pancreas			pCMV-SPORT6
L0662	NCI_CGAP_Gas4	poorly differentiated adenocarcinoma with signet r	stomach			pCMV-SPORT6
L0663	NCI_CGAP_Ut2	moderately- differentiated endometrial adenocarcino	uterus			pCMV-SPORT6
L0665	NCI_CGAP_Ut4	serous papillary carcinoma, high grade, 2 pooled t	uterus			pCMV-SPORT6
L0666	NCI_CGAP_Ut1	well-differentiated endometrial adenocarcinoma, 7	uterus			pCMV-SPORT6
L0717	Gessler Wilms tumor					pSPORT1
L0731	Soares_pregnant_uterus_ NbHPU		uterus			pT7T3-Pac
L0740	Soares melanocyte 2NbHM	melanocyte				pT7T3D (Pharmacia) with a modified polylinker
L0742	Soares adult brain N2b5HB55Y		brain			pT7T3D (Pharmacia) with a modified polylinker
L0743	Soares breast 2NbHBst		breast			pT7T3D (Pharmacia) with a modified polylinker
L0744	Soares breast 3NbHBst		breast			pT7T3D (Pharmacia) with a modified polylinker
L0745	Soares retina N2b4HR	retina	eye			pT7T3D (Pharmacia) with a modified polylinker
L0746	Soares retina N2b5HR	retina	eye			pT7T3D (Pharmacia) with a modified polylinker
L0747	Soares_fetal_heart_NbHH		heart			pT7T3D

2020040405040001

	19W					(Pharmacia) with a modified polylinker
L0748	Soares fetal liver spleen INFLS		Liver and Spleen			pT7T3D (Pharmacia) with a modified polylinker
L0749	Soares_fetal_liver_spleen _INFLS_S1		Liver and Spleen			pT7T3D (Pharmacia) with a modified polylinker
L0750	Soares_fetal_lung_NbHL1 9W		lung			pT7T3D (Pharmacia) with a modified polylinker
L0751	Soares ovary tumor NbHOT	ovarian tumor	ovary			pT7T3D (Pharmacia) with a modified polylinker
L0752	Soares_parathyroid_tumor _NbHPA	parathyroid tumor	parathyroid gland			pT7T3D (Pharmacia) with a modified polylinker
L0753	Soares_pineal_gland_N3H PG		pineal gland			pT7T3D (Pharmacia) with a modified polylinker
L0754	Soares placenta Nb2HP		placenta			pT7T3D (Pharmacia) with a modified polylinker
L0755	Soares_placenta_8to9wee ks_2NbHP8to9W		placenta			pT7T3D (Pharmacia) with a modified polylinker
L0756	Soares_multiple_sclerosis _2NbHMSP	multiple sclerosis lesions				pT7T3D (Pharmacia) with a modified polylinker V_TYPE
L0757	Soares_senescent_fibrobla sts_NbHSF	senescent fibroblast				pT7T3D (Pharmacia) with a modified polylinker V_TYPE
L0758	Soares_testis_NHT					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0759	Soares_total_fetus_Nb2H F8_9w					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0764	NCI_CGAP_Co3	colon				pT7T3D-Pac



10091504-030702

						(Pharmacia) with a modified polylinker
L0766	NCI_CGAP_GCB1	germinal center B cell				pT7T3D-Pac (Pharmacia) with a modified polylinker
L0769	NCI_CGAP_Brn25	anaplastic oligodendroglioma	brain			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0772	NCI_CGAP_Co10	colon tumor RER+	colon			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0774	NCI_CGAP_Kid3		kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0775	NCI_CGAP_Kid5	2 pooled tumors (clear cell type)	kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0776	NCI_CGAP_Lu5	carcinoid	lung			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0777	Soares_NhHMPu_S1	Pooled human melanocyte, fetal heart, and pregnant	mixed (see below)			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0779	Soares_NFL_T_GBC_S1		pooled			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0780	Soares_NSF_F8_9W_OT _PA_P_S1		pooled			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0783	NCI_CGAP_Pr22	normal prostate	prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0789	NCI_CGAP_Sub3					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0790	NCI_CGAP_Sub4					pT7T3D-Pac (Pharmacia) with a modified polylinker
L0794	NCI_CGAP_GC6	pooled germ cell tumors				pT7T3D-Pac (Pharmacia) with a modified

						polylinker
L0803	NCI_CGAP_Kid11		kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0804	NCI_CGAP_Kid12	2 pooled tumors (clear cell type)	kidney			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0805	NCI_CGAP_Lu24	carcinoid	lung			pT7T3D-Pac (Pharmacia) with a modified polylinker
L0809	NCI_CGAP_Pr28		prostate			pT7T3D-Pac (Pharmacia) with a modified polylinker

20091504, 030700

TABLE 5

OMIM Reference	Description
102200	Somatotrophinoma
102770	Myoadenylate deaminase deficiency
106100	Angioedema, hereditary
107970	Arrhythmogenic right ventricular dysplasia-1
108725	Atherosclerosis, susceptibility to
108962	Hypertension, salt-resistant
109400	Basal cell nevus syndrome
109543	Leukemia, chronic lymphocytic, B-cell
115650	Cataract, anterior polar-1
118210	Charcot-Marie-Tooth neuropathy-2A
120550	C1q deficiency, type A
120570	C1q deficiency, type B
120575	C1q deficiency, type C
120700	C3 deficiency
121011	Deafness, autosomal dominant 3, 601544
121011	Deafness, autosomal recessive 1, 220290
121800	Corneal dystrophy, crystalline, Schnyder
123270	[Creatine kinase, brain type, ectopic expression of]
129500	Ectodermal dysplasia, hidrotic
130500	Elliptocytosis-1
131100	Multiple endocrine neoplasia I
131100	Prolactinoma, hyperparathyroidism, carcinoid syndrome
131100	Carcinoid tumor of lung
132800	Basal cell carcinoma
132800	Epithelioma, self-healing, squamous 1, Ferguson-Smith type
133171	[Erythrocytosis, familial], 133100
133200	Erythrokeratoderma variabilis
133780	Vitreoretinopathy, exudative, familial
138140	Glucose transport defect, blood-brain barrier
138971	Kostmann neutropenia, 202700
143890	Hypercholesterolemia, familial
146150	Hypomelanosis of Ito
147050	Atopy
147670	Rabson-Mendenhall syndrome
147670	Diabetes mellitus, insulin-resistant, with acanthosis nigricans
147670	Leprechaunism
150250	Larsen syndrome, autosomal dominant
151440	Leukemia, T-cell acute lymphoblastoid
153700	Macular dystrophy, vitelliform type
155600	Malignant melanoma, cutaneous
161015	Mitochondrial complex I deficiency, 252010
164009	Leukemia, acute promyelocytic, NUMA/RARA type
164500	Spinocerebellar ataxia-7
164953	Liposarcoma

168360	Paraneoplastic sensory neuropathy
168461	Multiple myeloma, 254250
168461	Parathyroid adenomatosis 1
168461	Centrocytic lymphoma
168468	Metaphyseal chondrodysplasia, Murk Jansen type, 156400
171760	Hypophosphatasia, adult, 146300
171760	Hypophosphatasia, infantile, 241500
172411	Colorectal cancer, resistance to
176100	Porphyria cutanea tarda
176100	Porphyria, hepatoerythropoietic
177070	Spherocytosis, hereditary, Japanese type
177070	Hermansky-Pudlak syndrome, 203300
178300	Ptosis, hereditary congenital, 1
180297	Anemia, hemolytic, Rh-null, suppressor type, 268150
180721	Retinitis pigmentosa, digenic
180840	Susceptibility to IDDM
182280	Small-cell cancer of lung
182500	Cataract, congenital
182600	Spastic paraplegia-3A
185470	Myopathy due to succinate dehydrogenase deficiency
186855	Leukemia-2, T-cell acute lymphoblastic
188540	Hypothyroidism, nongoitrous
188826	Sorsby fundus dystrophy, 136900
191181	Cervical carcinoma
193235	Vitreoretinopathy, neovascular inflammatory
203800	Alstrom syndrome
209901	Bardet-Biedl syndrome 1
211420	Breast cancer, ductal
216550	Cohen syndrome
218000	Andermann syndrome
223900	Dysautonomia, familial
227220	[Eye color, brown]
230000	Fucosidosis
230350	Galactose epimerase deficiency
230450	Hemolytic anemia due to gamma-glutamylcysteine synthetase deficiency
231670	Glutaricaciduria, type I
232600	McArdle disease
238310	Hyperglycinemia, nonketotic, type II
243500	Isovalericacidemia
245200	Krabbe disease
248611	Maple syrup urine disease, type Ib
250100	Metachromatic leukodystrophy
250800	Methemoglobinemia, type I
250800	Methemoglobinemia, type II
251600	Microphthalmia, autosomal recessive
253700	Muscular dystrophy, limb-girdle, type 2C

253800	Walker-Warburg syndrome, 236670
253800	Fukuyama type congenital muscular dystrophy
255800	Schwartz-Jampel syndrome
259700	Osteopetrosis, recessive
259770	Osteoporosis-pseudoglioma syndrome
263200	Polycystic kidney disease, autosomal recessive
268900	[Sarcosinemia]
270100	Situs inversus viscerum
276900	Usher syndrome, type 1A
277730	Wernicke-Korsakoff syndrome, susceptibility to
278700	Xeroderma pigmentosum, group A
600045	Xeroderma pigmentosum, group E, subtype 2
600163	Long QT syndrome-3
600234	HMG-CoA synthase-2 deficiency
600258	Colorectal cancer, hereditary nonpolyposis, type 3
600276	Cerebral arteriopathy with subcortical infarcts and leukoencephalopathy, 125310
600319	Diabetes mellitus, insulin-dependent, 4
600528	CPT deficiency, hepatic, type I, 255120
600631	Enuresis, nocturnal, 1
600837	Hirschsprung disease, 142623
600839	Bartter syndrome, 241200
600957	Persistent Mullerian duct syndrome, type I, 261550
600971	Deafness, autosomal recessive 6
601226	Progressive external ophthalmoplegia, type 2
601414	Retinitis pigmentosa-18
601499	Rieger syndrome, type 2
601690	Platelet-activating factor acetylhydrolase deficiency
601691	Retinitis pigmentosa-19, 601718
601691	Stargardt disease-1, 248200
601691	Cone-rod dystrophy 3
601691	Fundus flavimaculatus with macular dystrophy, 248200
601718	Retinitis pigmentosa-19
601800	[Hair color, brown]
601843	Hypothyroidism, congenital, 274400
601884	[High bone mass]
601885	Cataract, zonular pulverulent-2
601916	Pancreatic cancer
601990	Neuroblastoma
602023	Bartter syndrome, type 3
602087	Arrhythmogenic right ventricular dysplasia-4
602088	Nephronophthisis, infantile
602091	Marfan syndrome, atypical
602094	Lipodystrophy, familial partial
602221	Stem-cell leukemia/lymphoma syndrome
602771	Muscular dystrophy, congenital, with early spine rigidity

### *Polynucleotide and Polypeptide Variants*

[0099] The present invention is also directed to variants of the cardiovascular system associated polynucleotide sequence disclosed in SEQ ID NO:X or the complementary strand thereto, nucleotide sequences encoding the polypeptide of SEQ ID NO:Y, the nucleotide sequence of SEQ ID NO:X encoding the polypeptide sequence as defined in column 6 of Table 1A, nucleotide sequences encoding the polypeptide as defined in column 6 of Table 1A, the nucleotide sequence as defined in columns 8 and 9 of Table 2, nucleotide sequences encoding the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2, the nucleotide sequence as defined in column 6 of Table 1B, nucleotide sequences encoding the polypeptide encoded by the nucleotide sequence as defined in column 6 of Table 1B, the cDNA sequence contained in Clone ID NO:Z, and/or nucleotide sequences encoding a polypeptide encoded by the cDNA sequence contained in Clone ID NO:Z.

[00100] The present invention also encompasses variants of the polypeptide sequence disclosed in SEQ ID NO:Y, a polypeptide sequence as defined in column 6 of Table 1A, a polypeptide sequence encoded by the polynucleotide sequence in SEQ ID NO:X, a polypeptide sequence encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2, a polypeptide sequence encoded by the nucleotide sequence as defined in column 6 of Table 1B, a polypeptide sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X, and/or a polypeptide sequence encoded by the cDNA sequence contained in Clone ID NO:Z.

[00101] "Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

[00102] Thus, one aspect of the invention provides an isolated nucleic acid molecule comprising, or alternatively consisting of, a polynucleotide having a nucleotide sequence selected from the group consisting of: (a) a nucleotide sequence described in SEQ ID NO:X or contained in the cDNA sequence of Clone ID NO:Z; (b) a nucleotide sequence in SEQ ID NO:X or the cDNA in Clone ID NO:Z which encodes a mature cardiovascular system associated polypeptide; (c) a nucleotide sequence in SEQ ID NO:X or the cDNA sequence of Clone ID NO:Z, which

encodes a biologically active fragment of a cardiovascular system associated polypeptide; (d) a nucleotide sequence in SEQ ID NO:X or the cDNA sequence of Clone ID NO:Z, which encodes an antigenic fragment of a cardiovascular system associated polypeptide; (e) a nucleotide sequence encoding a cardiovascular system associated polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (f) a nucleotide sequence encoding a mature cardiovascular system associated polypeptide of the amino acid sequence of SEQ ID NO:Y or the amino acid sequence encoded by the cDNA in Clone ID NO:Z; (g) a nucleotide sequence encoding a biologically active fragment of a cardiovascular system associated polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (h) a nucleotide sequence encoding an antigenic fragment of a cardiovascular system associated polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; and (i) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), or (h), above.

**[00103]** The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100%, identical to, for example, any of the nucleotide sequences in (a), (b), (c), (d), (e), (f), (g), (h), or (i) above, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence of the cDNA contained in Clone ID NO:Z or the complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding a polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO:X, a polypeptide sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X, a nucleotide sequence encoding the polypeptide encoded by the cDNA contained in Clone ID NO:Z, the nucleotide coding sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto, a nucleotide sequence encoding the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto, the nucleotide coding sequence in SEQ ID NO:B as defined in column 6 of Table 1B or

the complementary strand thereto, a nucleotide sequence encoding the polypeptide encoded by the nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B or the complementary strand thereto, the nucleotide sequence in SEQ ID NO:X encoding the polypeptide sequence as defined in column 6 of Table 1A or the complementary strand thereto, nucleotide sequences encoding a polypeptide as defined in column 6 of Table 1A or the complementary strand thereto, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein). Polynucleotides which hybridize to the complement of these nucleic acid molecules under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides and nucleic acids.

**[00104]** In a preferred embodiment, the invention encompasses nucleic acid molecules which comprise, or alternatively, consist of a polynucleotide which hybridizes under stringent hybridization conditions, or alternatively, under lower stringency conditions, to a polynucleotide in (a), (b), (c), (d), (e), (f), (g), (h), or (i) above, as are polypeptides encoded by these polynucleotides. In another preferred embodiment, polynucleotides which hybridize to the complement of these nucleic acid molecules under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

**[00105]** In another embodiment, the invention provides a purified protein comprising, or alternatively consisting of, a polypeptide having an amino acid sequence selected from the group consisting of: (a) the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; (b) the amino acid sequence of a mature cardiovascular system associated polypeptide having the amino acid sequence of SEQ ID NO:Y or the amino acid sequence encoded by the cDNA in Clone ID NO:Z; (c) the amino acid sequence of a biologically active fragment of a cardiovascular system associated polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z; and (d) the amino acid sequence of an antigenic fragment of a cardiovascular system associated polypeptide having the complete amino acid sequence of SEQ ID NO:Y or the complete amino acid sequence encoded by the cDNA in Clone ID NO:Z.



10091504-030702

[00106] The present invention is also directed to proteins which comprise, or alternatively consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100%, identical to, for example, any of the amino acid sequences in (a), (b), (c), or (d), above, the amino acid sequence shown in SEQ ID NO:Y, the amino acid sequence encoded by the cDNA contained in Clone ID NO:Z, the amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X as defined in columns 8 and 9 of Table 2, the amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B, the amino acid sequence as defined in column 6 of Table 1A, an amino acid sequence encoded by the nucleotide sequence in SEQ ID NO:X, and an amino acid sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X. Fragments of these polypeptides are also provided (e.g., those fragments described herein). Further proteins encoded by polynucleotides which hybridize to the complement of the nucleic acid molecules encoding these amino acid sequences under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are the polynucleotides encoding these proteins.

[00107] By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be an entire sequence referred to in Table 1A or 2 as the ORF (open reading frame), or any fragment specified, as described herein.

[00108] As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best

overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be compared by converting U's to T's. The result of said global sequence alignment is expressed as percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30, Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

**[00109]** If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of manually adjusting the percent identity score.

**[00110]** For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final

percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[00111] By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[00112] As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence of a polypeptide referred to in Table 1A (e.g., an amino acid sequence identified in columns 5 or 6) or Table 2 (e.g., the amino acid sequence of the polypeptide encoded by the polynucleotide sequence defined in columns 8 and 9 of Table 2) or a fragment thereof, the amino acid sequence of the polypeptide encoded by the polynucleotide sequence in SEQ ID NO:B as defined in column 6 of Table 1B or a fragment thereof, the amino acid sequence of the polypeptide encoded by the nucleotide sequence in SEQ ID NO:X or a fragment thereof, or an amino acid sequence of the polypeptide encoded by cDNA contained in Clone ID NO:Z, or a fragment thereof, can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a

query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci.6:237-245 (1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is expressed as percent identity. Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

**[00113]** If the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C- terminal residues of the subject sequence.

**[00114]** For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and

10093504.030702

C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to be made for the purposes of the present invention.

[00115] The polynucleotide variants of the invention may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations, which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, polypeptide variants in which less than 50, less than 40, less than 30, less than 20, less than 10, or 5-50, 5-25, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

[00116] Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

[00117] Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, one or more amino acids can be deleted from the N-terminus or C-terminus of the polypeptides of the present invention without

substantial loss of biological function. As an example, the authors of Ron et al., J. Biol. Chem. 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., J. Biotechnology 7:199-216 (1988).)

[00118] Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (J. Biol. Chem. 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

[00119] Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies, which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

[00120] Thus, the invention further includes polypeptide variants which show a functional activity (e.g., biological activity) of the polypeptides of the invention. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity.

[00121] The present application is directed to nucleic acid molecules at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences

disclosed herein, (e.g., encoding a polypeptide having the amino acid sequence of an N and/or C terminal deletion), irrespective of whether they encode a polypeptide having functional activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having functional activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having functional activity include, *inter alia*, (1) isolating a gene or allelic or splice variants thereof in a cDNA library; (2) *in situ* hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the gene, as described in Verma et al., Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York (1988); (3) Northern Blot analysis for detecting mRNA expression in specific tissues (e.g., normal cardiovascular system or diseased cardiovascular system tissues); and (4) *in situ* hybridization (e.g., histochemistry) for detecting mRNA expression in specific tissues (e.g., normal cardiovascular system or diseased cardiovascular system tissues).

[00122] Preferred, however, are nucleic acid molecules having sequences at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences disclosed herein, which do, in fact, encode a polypeptide having functional activity. By a polypeptide having "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) protein of the invention. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide of the invention for binding) to an anti-polypeptide of the invention antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide of the invention.

[00123] The functional activity of the polypeptides, and fragments, variants and derivatives of the invention, can be assayed by various methods.

[00124] For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the present invention for binding to an anti-polypeptide of the invention antibody, various immunoassays known in the art can

be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

[00125] In another embodiment, where a ligand is identified, or the ability of a polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky et al., Microbiol. Rev. 59:94-123 (1995). In another embodiment, the ability of physiological correlates of a polypeptide of the present invention to bind to a substrate(s) of the polypeptide of the invention can be routinely assayed using techniques known in the art.

[00126] In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the present invention and fragments, variants and derivatives thereof to elicit polypeptide related biological activity (either *in vitro* or *in vivo*). Other methods will be known to the skilled artisan and are within the scope of the invention.

[00127] Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to, for example, the nucleic acid sequence of the cDNA contained in Clone ID NO:Z, a nucleic acid sequence referred to in Table 1A (e.g., SEQ ID NO:X), a nucleic acid sequence disclosed in Table 2 (e.g., the nucleic acid sequence



delineated in columns 8 and 9) or fragments thereof, will encode polypeptides "having functional activity." In fact, since degenerate variants of any of these nucleotide sequences all encode the same polypeptide, in many instances, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having functional activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid), as further described below.

[00128] For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," Science 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

[00129] The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

[00130] The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. See Cunningham et al., Science 244:1081-1085 (1989). The resulting mutant molecules can then be tested for biological activity.

[00131] As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein)

amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly. Besides conservative amino acid substitutions, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitutions with one or more of the amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, serum albumin (preferably human serum albumin) or a fragment or variant thereof, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

[00132] For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. See Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).

[00133] A further embodiment of the invention relates to polypeptides which comprise the amino acid sequence of a polypeptide having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions from a polypeptide sequence disclosed herein. Of course it is highly preferable for a polypeptide to have an

amino acid sequence which comprises the amino acid sequence of a polypeptide of SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X, an amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, an amino acid sequence encoded by the complement of SEQ ID NO:X, and/or the amino acid sequence encoded by cDNA contained in Clone ID NO:Z which contains, in order of ever-increasing preference, at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1 amino acid substitutions.

[00134] In specific embodiments, the polypeptides of the invention comprise, or alternatively, consist of, fragments or variants of a reference amino acid sequence selected from: (a) the amino acid sequence of SEQ ID NO:Y or fragments thereof (e.g., the mature form and/or other fragments described herein); (b) the amino acid sequence encoded by SEQ ID NO:X or fragments thereof; (c) the amino acid sequence encoded by the complement of SEQ ID NO:X or fragments thereof; (d) the amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or fragments thereof; and (e) the amino acid sequence encoded by cDNA contained in Clone ID NO:Z or fragments thereof; wherein the fragments or variants have 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, amino acid residue additions, substitutions, and/or deletions when compared to the reference amino acid sequence. In preferred embodiments, the amino acid substitutions are conservative. Polynucleotides encoding these polypeptides are also encompassed by the invention.

#### ***Polynucleotide and Polypeptide Fragments***

[00135] The present invention is also directed to polynucleotide fragments of the polynucleotides (nucleic acids) of the invention. In the present invention, a "polynucleotide fragment" refers to a polynucleotide having a nucleic acid sequence which, for example: is a portion of the cDNA contained in Clone ID NO:Z or the complementary strand thereto; is a portion of the polynucleotide sequence encoding the polypeptide encoded by the cDNA contained in Clone ID NO:Z or the complementary strand thereto; is a portion of a polynucleotide sequence encoding the amino acid sequence encoded by the region of SEQ ID NO:X as defined in columns 8 and 9 of Table 2 or the complementary strand thereto; is a portion of the polynucleotide sequence of SEQ ID NO:X as defined in columns 8 and 9 of Table 2

or the complementary strand thereto; is a portion of the polynucleotide sequence in SEQ ID NO:X or the complementary strand thereto; is a polynucleotide sequence encoding a portion of the polypeptide of SEQ ID NO:Y; is a polynucleotide sequence encoding a portion of a polypeptide encoded by SEQ ID NO:X; is a polynucleotide sequence encoding a portion of a polypeptide encoded by the complement of the polynucleotide sequence in SEQ ID NO:X; is a portion of a polynucleotide sequence encoding the amino acid sequence encoded by the region of SEQ ID NO:B as defined in column 6 of Table 1B or the complementary strand thereto; or is a portion of the polynucleotide sequence of SEQ ID NO:B as defined in column 6 of Table 1B or the complementary strand thereto.

**[00136]** The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from the cDNA sequence contained in Clone ID NO:Z, or the nucleotide sequence shown in SEQ ID NO:X or the complementary stand thereto. In this context "about" includes the particularly recited value or a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., at least 160, 170, 180, 190, 200, 250, 500, 600, 1000, or 2000 nucleotides in length) are also encompassed by the invention.

**[00137]** Moreover, representative examples of polynucleotide fragments of the invention, comprise, or alternatively consist of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-

3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, 4101-4150, 4151-4200, 4201-4250, 4251-4300, 4301-4350, 4351-4400, 4401-4450, 4451-4500, 4501-4550, 4551-4600, 4601-4650, 4651-4700, 4701-4750, 4751-4800, 4801-4850, 4851-4900, 4901-4950, 4951-5000, 5001-5050, 5051-5100, 5101-5150, 5151-5200, 5201-5250, 5251-5300, 5301-5350, 5351-5400, 5401-5450, 5451-5500, 5501-5550, 5551-5600, 5601-5650, 5651-5700, 5701-5750, 5751-5800, 5801-5850, 5851-5900, 5901-5950, 5951-6000, 6001-6050, 6051-6100, 6101-6150, 6151-6200, 6201-6250, 6251-6300, 6301-6350, 6351-6400, 6401-6450, 6451-6500, 6501-6550, 6551-6600, 6601-6650, 6651-6700, 6701-6750, 6751-6800, 6801-6850, 6851-6900, 6901-6950, 6951-7000, 7001-7050, 7051-7100, 7101-7150, 7151-7200, 7201-7250, 7251-7300 or 7301 to the end of SEQ ID NO:X, or the complementary strand thereto. In this context "about" includes the particularly recited range or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide, which has a functional activity (e.g., biological activity). More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

**[00138]** Further representative examples of polynucleotide fragments of the invention, comprise, or alternatively consist of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, 4101-

4150, 4151-4200, 4201-4250, 4251-4300, 4301-4350, 4351-4400, 4401-4450, 4451-4500, 4501-4550, 4551-4600, 4601-4650, 4651-4700, 4701-4750, 4751-4800, 4801-4850, 4851-4900, 4901-4950, 4951-5000, 5001-5050, 5051-5100, 5101-5150, 5151-5200, 5201-5250, 5251-5300, 5301-5350, 5351-5400, 5401-5450, 5451-5500, 5501-5550, 5551-5600, 5601-5650, 5651-5700, 5701-5750, 5751-5800, 5801-5850, 5851-5900, 5901-5950, 5951-6000, 6001-6050, 6051-6100, 6101-6150, 6151-6200, 6201-6250, 6251-6300, 6301-6350, 6351-6400, 6401-6450, 6451-6500, 6501-6550, 6551-6600, 6601-6650, 6651-6700, 6701-6750, 6751-6800, 6801-6850, 6851-6900, 6901-6950, 6951-7000, 7001-7050, 7051-7100, 7101-7150, 7151-7200, 7201-7250, 7251-7300 or 7301 to the end of the cDNA sequence contained in Clone ID NO:Z, or the complementary strand thereto. In this context "about" includes the particularly recited range or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide, which has a functional activity (e.g., biological activity). More preferably, these polynucleotides can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

**[00139]** Moreover, representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a nucleic acid sequence comprising one, two, three, four, five, six, seven, eight, nine, ten, or more of the above described polynucleotide fragments of the invention in combination with a polynucleotide sequence delineated in Table 1B column 6. Additional, representative examples of polynucleotide fragments of the invention comprise, or alternatively consist of, a nucleic acid sequence comprising one, two, three, four, five, six, seven, eight, nine, ten, or more of the above described polynucleotide fragments of the invention in combination with a polynucleotide sequence that is the complementary strand of a sequence delineated in column 6 of Table 1B. In further embodiments, the above-described polynucleotide fragments of the invention comprise, or alternatively consist of, sequences delineated in Table 1B, column 6, and have a nucleic acid sequence which is different from that of the BAC fragment having the sequence disclosed in SEQ ID NO:B (see Table 1B, column 5). In additional embodiments,

the above-described polynucleotide fragments of the invention comprise, or alternatively consist of, sequences delineated in Table 1B, column 6, and have a nucleic acid sequence which is different from that published for the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). In additional embodiments, the above-described polynucleotides of the invention comprise, or alternatively consist of, sequences delineated Table 1B, column 6, and have a nucleic acid sequence which is different from that contained in the BAC clone identified as BAC ID NO:A (see Table 1B, column 4). Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides and polypeptides are also encompassed by the invention.

**[00140]** In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more fragments of the sequences delineated in column 6 of Table 1B, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1B, column 2) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

**[00141]** In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more fragments of the sequences delineated in column 6 of Table 1B which correspond to the same Clone ID NO:Z (see Table 1B, column 1), and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A or 1B) or fragments or variants thereof. Polypeptides encoded by these polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

**[00142]** In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of, one, two, three, four, five, six, seven, eight, nine, ten, or more fragments of the sequences delineated in the same row of column 6 of Table 1B, and the polynucleotide sequence of SEQ ID NO:X (e.g., as defined in Table 1A or 1B) or fragments or variants thereof. Polypeptides encoded by these

polynucleotides, other polynucleotides that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention.

[00143] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of the sequence of SEQ ID NO:X are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids that encode these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[00144] In additional specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X (e.g., as described herein) are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[00145] In further specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of a fragment or variant of the sequence of SEQ ID NO:X and the 5' 10 polynucleotides of the sequence of one of the sequences delineated in column 6 of Table 1B are directly contiguous. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization



20200914051507

conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[00146] In specific embodiments, polynucleotides of the invention comprise, or alternatively consist of a polynucleotide sequence in which the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B and the 5' 10 polynucleotides of another sequence in column 6 are directly contiguous. In preferred embodiments, the 3' 10 polynucleotides of one of the sequences delineated in column 6 of Table 1B is directly contiguous with the 5' 10 polynucleotides of the next sequential exon delineated in Table 1B, column 6. Nucleic acids which hybridize to the complement of these 20 contiguous polynucleotides under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention. Polypeptides encoded by these polynucleotides and/or nucleic acids, other polynucleotides and/or nucleic acids encoding these polypeptides, and antibodies that bind these polypeptides are also encompassed by the invention. Additionally, fragments and variants of the above-described polynucleotides, nucleic acids, and polypeptides are also encompassed by the invention.

[00147] In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y, a portion of an amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, a portion of an amino acid sequence encoded by the polynucleotide sequence of SEQ ID NO:X, a portion of an amino acid sequence encoded by the complement of the polynucleotide sequence in SEQ ID NO:X, and/or a portion of an amino acid sequence encoded by the cDNA contained in Clone ID NO:Z. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-

80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, 281-300, 301-320, 321-340, 341-360, 361-380, 381-400, 401-420, 421-440, 441-460, 461-480, 481-500, 501-520, 521-540, 541-560, 561-580, 581-600, 601-620, 621-640, 641-660, 661-680, 681-700, 701-720, 721-740, 741-760, 761-780, 781-800, 801-820, 821-840, 841-860, 861-880, 881-900, 901-920, 921-940, 941-960, 961-980, 981-1000, 1001-1020, 1021-1040, 1041-1060, 1061-1080, 1081-1100, 1101-1120, 1121-1140, 1141-1160, 1161-1180, 1181-1200, 1201-1220, 1221-1240, 1241-1260, 1261-1280, 1281-1300, 1301-1320, 1321-1340, 1341-1360, 1361-1380, 1381-1400, 1401-1420, 1421-1440, or 1441 to the end of the coding region. In a preferred embodiment, polypeptide fragments of the invention include, for example, fragments comprising, or alternatively consisting of, from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, 281-300, 301-320, 321-340, 341-360, 361-380, 381-400, 401-420, 421-440, 441-460, 461-480, 481-500, 501-520, 521-540, 541-560, 561-580, 581-600, 601-620, 621-640, 641-660, 661-680, 681-700, 701-720, 721-740, 741-760, 761-780, 781-800, 801-820, 821-840, 841-860, 861-880, 881-900, 901-920, 921-940, 941-960, 961-980, 981-1000, 1001-1020, 1021-1040, 1041-1060, 1061-1080, 1081-1100, 1101-1120, 1121-1140, 1141-1160, 1161-1180, 1181-1200, 1201-1220, 1221-1240, 1241-1260, 1261-1280, 1281-1300, 1301-1320, 1321-1340, 1341-1360, 1361-1380, 1381-1400, 1401-1420, 1421-1440, or 1441 to the end of the coding region of SEQ ID NO:Y. Moreover, polypeptide fragments of the invention may be at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, or ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either extreme or at both extremes. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

- [00148] Even if deletion of one or more amino acids from the N-terminus of a protein results in modification of loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example, the ability of shortened muteins to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the

residues of the complete or mature polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted N-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

[00149] Accordingly, polypeptide fragments include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above amino and carboxy terminus deletions is preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

[00150] The present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of a polypeptide disclosed herein (e.g., a polypeptide of SEQ ID NO:Y, a polypeptide encoded by the polynucleotide sequence contained in SEQ ID NO:X or the complement thereof, a polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, a polypeptide encoded by the portion of SEQ ID NO:B as defined in column 6 of Table 1B, and/or a polypeptide encoded by the cDNA contained in Clone ID NO:Z). In particular, N-terminal deletions may be described by the general formula m-q, where q is a whole integer representing the total number of amino acid residues in a polypeptide of the invention (e.g., the polypeptide disclosed in SEQ ID NO:Y, or the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2), and m is defined as any integer ranging from 2 to q-6. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00151] The present invention further provides polypeptides having one or more residues from the carboxy terminus of the amino acid sequence of a polypeptide disclosed

herein (e.g., a polypeptide of SEQ ID NO:Y, a polypeptide encoded by the polynucleotide sequence contained in SEQ ID NO:X, a polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or a polypeptide encoded by the cDNA contained in Clone ID NO:Z). In particular, C-terminal deletions may be described by the general formula 1-n, where n is any whole integer ranging from 6 to q-1, and where n corresponds to the position of amino acid residue in a polypeptide of the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00152] In addition, any of the above described N- or C-terminal deletions can be combined to produce a N- and C-terminal deleted polypeptide. The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of a polypeptide encoded by SEQ ID NO:X (e.g., including, but not limited to, the preferred polypeptide disclosed as SEQ ID NO:Y and the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2), the cDNA contained in Clone ID NO:Z, and/or the complement thereof, where n and m are integers as described above. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00153] Also as mentioned above, even if deletion of one or more amino acids from the C-terminus of a protein results in modification or loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example the ability of the shortened mutein to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptide generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the C-terminus. Whether a particular polypeptide lacking C-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted C-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

[00154] The present application is also directed to proteins containing polypeptides at

least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a polypeptide sequence set forth herein. In preferred embodiments, the application is directed to proteins containing polypeptides at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to polypeptides having the amino acid sequence of the specific N- and C-terminal deletions. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00155] Any polypeptide sequence encoded by, for example, the polynucleotide sequences set forth as SEQ ID NO:X or the complement thereof, (presented, for example, in Tables 1A and 2), the cDNA contained in Clone ID NO:Z, or the polynucleotide sequence as defined in column 6 of Table 1B, may be analyzed to determine certain preferred regions of the polypeptide. For example, the amino acid sequence of a polypeptide encoded by a polynucleotide sequence of SEQ ID NO:X (e.g., the polypeptide of SEQ ID NO:Y and the polypeptide encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2) or the cDNA contained in Clone ID NO:Z may be analyzed using the default parameters of the DNASTAR computer algorithm (DNASTAR, Inc., 1228 S. Park St., Madison, WI 53715 USA; <http://www.dnastar.com/>).

[00156] Polypeptide regions that may be routinely obtained using the DNASTAR computer algorithm include, but are not limited to, Garnier-Robson alpha-regions, beta-regions, turn-regions, and coil-regions; Chou-Fasman alpha-regions, beta-regions, and turn-regions; Kyte-Doolittle hydrophilic regions and hydrophobic regions; Eisenberg alpha- and beta-amphipathic regions; Karplus-Schulz flexible regions; Emini surface-forming regions; and Jameson-Wolf regions of high antigenic index. Among highly preferred polynucleotides of the invention in this regard are those that encode polypeptides comprising regions that combine several structural features, such as several (e.g., 1, 2, 3 or 4) of the features set out above.

[00157] Additionally, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Emini surface-forming regions, and Jameson-Wolf regions of high antigenic index (i.e., containing four or more contiguous amino acids having an antigenic index of greater than or equal to 1.5, as identified using the default parameters of the Jameson-Wolf program) can routinely be used to determine polypeptide regions that exhibit a high degree of potential for antigenicity. Regions of high antigenicity are determined from data by DNASTAR analysis by choosing values which represent

regions of the polypeptide which are likely to be exposed on the surface of the polypeptide in an environment in which antigen recognition may occur in the process of initiation of an immune response.

[00158] Preferred polypeptide fragments of the invention are fragments comprising, or alternatively, consisting of, an amino acid sequence that displays a functional activity (e.g. biological activity) of the polypeptide sequence of which the amino acid sequence is a fragment. By a polypeptide displaying a "functional activity" is meant a polypeptide capable of one or more known functional activities associated with a full-length protein, such as, for example, biological activity, antigenicity, immunogenicity, and/or multimerization, as described herein.

[00159] Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

[00160] In preferred embodiments, polypeptides of the invention comprise, or alternatively consist of, one, two, three, four, five or more of the antigenic fragments of the polypeptide of SEQ ID NO:Y, or portions thereof. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00161] The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of: the polypeptide sequence shown in SEQ ID NO:Y; a polypeptide sequence encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2; the polypeptide sequence encoded by the portion of SEQ ID NO:B as defined in column 6 of Table 1B or the complement thereto; the polypeptide sequence encoded by the cDNA contained in Clone ID NO:Z; or the polypeptide sequence encoded by a polynucleotide that hybridizes to the sequence of SEQ ID NO:X, the complement of the sequence of SEQ ID NO:X, the complement of a portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, or the cDNA sequence contained in Clone ID NO:Z under stringent hybridization conditions or alternatively, under lower stringency hybridization as defined *supra*. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for

202009-1057507

example, the sequence disclosed in SEQ ID NO:X, or a fragment thereof), polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to the complementary strand under stringent hybridization conditions or alternatively, under lower stringency hybridization conditions defined *supra*.

**[00162]** The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described *infra*. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

**[00163]** Fragments, which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

**[00164]** In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any

combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

[00165] Non-limiting examples of epitopes of polypeptides that can be used to generate antibodies of the invention include a polypeptide comprising, or alternatively consisting of, at least one, two, three, four, five, six or more of the portion(s) of SEQ ID NO:Y specified in column 6 of Table 1A. These polypeptide fragments have been determined to bear antigenic epitopes of the proteins of the invention by the analysis of the Jameson-Wolf antigenic index, which is included in the DNASTar suite of computer programs. By "comprise" it is intended that a polypeptide contains at least one, two, three, four, five, six or more of the portion(s) of SEQ ID NO:Y shown in column 6 of Table 1A, but it may contain additional flanking residues on either the amino or carboxyl termini of the recited portion. Such additional flanking sequences are preferably sequences naturally found adjacent to the portion; i.e., contiguous sequence shown in SEQ ID NO:Y. The flanking sequence may, however, be sequences from a heterologous polypeptide, such as from another protein described herein or from a heterologous polypeptide not described herein. In particular embodiments, epitope portions of a polypeptide of the invention comprise one, two, three, or more of the portions of SEQ ID NO:Y shown in column 6 of Table 1A. Polynucleotides encoding these polypeptides are also encompassed by the invention.

[00166] Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. See, for instance, Sutcliffe et al., *supra*; Wilson et al., *supra*; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have



been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

[00167] Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, *in vivo* immunization, *in vitro* immunization, and phage display methods. See, e.g., Sutcliffe et al., *supra*; Wilson et al., *supra*, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If *in vivo* immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

[00168] As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention (e.g., those comprising an immunogenic or antigenic epitope) can be fused to heterologous polypeptide sequences. For example, polypeptides of the present invention (including fragments or variants thereof), may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof, resulting in chimeric polypeptides. By way of another non-limiting example, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) may be fused with albumin (including but not limited to recombinant human serum albumin or fragments or variants thereof (see, e.g.,

U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)). In a preferred embodiment, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) are fused with the mature form of human serum albumin (i.e., amino acids 1 – 585 of human serum albumin as shown in Figures 1 and 2 of EP Patent 0 322 094) which is herein incorporated by reference in its entirety. In another preferred embodiment, polypeptides and/or antibodies of the present invention (including fragments or variants thereof) are fused with polypeptide fragments comprising, or alternatively consisting of, amino acid residues 1-z of human serum albumin, where z is an integer from 369 to 419, as described in U.S. Patent 5,766,883 herein incorporated by reference in its entirety. Polypeptides and/or antibodies of the present invention (including fragments or variants thereof) may be fused to either the N- or C-terminal end of the heterologous protein (e.g., immunoglobulin Fc polypeptide or human serum albumin polypeptide). Polynucleotides encoding fusion proteins of the invention are also encompassed by the invention.

[00169] Such fusion proteins as those described above may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion desulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995). Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin (HA) tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, *Proc. Natl. Acad. Sci. USA*

88:8972- 897). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto Ni<sup>2+</sup> nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

### ***Fusion Proteins***

- [00170]** Any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on trafficking signals, polypeptides of the present invention which are shown to be secreted can be used as targeting molecules once fused to other proteins.
- [00171]** Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker sequences.
- [00172]** In certain preferred embodiments, proteins of the invention are fusion proteins comprising an amino acid sequence that is an N and/or C- terminal deletion of a polypeptide of the invention. In preferred embodiments, the invention is directed to a fusion protein comprising an amino acid sequence that is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a polypeptide sequence of the invention. Polynucleotides encoding these proteins are also encompassed by the invention.
- [00173]** Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final

preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

[00174] As one of skill in the art will appreciate that, as discussed above, polypeptides of the present invention, and epitope-bearing fragments thereof, can be combined with heterologous polypeptide sequences. For example, the polypeptides of the present invention may be fused with heterologous polypeptide sequences, for example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM) or portions thereof (CH1, CH2, CH3, and any combination thereof, including both entire domains and portions thereof), or albumin (including, but not limited to, native or recombinant human albumin or fragments or variants thereof (see, e.g., U.S. Patent No. 5,876,969, issued March 2, 1999, EP Patent 0 413 622, and U.S. Patent No. 5,766,883, issued June 16, 1998, herein incorporated by reference in their entirety)), resulting in chimeric polypeptides. For example, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties (EP-A 0232 262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. See, D. Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); K. Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995).

[00175] Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a polypeptide, which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification

of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., Cell 37:767 (1984).)

[00176] Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"), briefly described below, and further described herein. DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, Trends Biotechnol. 16(2):76-82 (1998); Hansson et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo and Blasco, Biotechniques 24(2):308-13 (1998); each of these patents and publications are hereby incorporated by reference in its entirety). In a preferred embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc., of one or more heterologous molecules encoding a heterologous polypeptide.

[00177] Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

#### **Recombinant and Synthetic Production of Polypeptides of the Invention**

[00178] The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by synthetic and recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

[00179] The polynucleotides of the invention may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex

with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

[00180] The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli lac*, *trp*, *phoA* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

[00181] As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance, glutamine synthase, for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, 293, NSO and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

[00182] Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not limited to pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalph, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, pPIC9K, and

PAO815 (all available from Invitrogen, Carlsbad, CA). Other suitable vectors will be readily apparent to the skilled artisan.

[00183] Vectors which use glutamine synthase (GS) or DHFR as the selectable markers can be amplified in the presence of the drugs methionine sulphoximine or methotrexate, respectively. An advantage of glutamine synthase based vectors is the availability of cell lines (e.g., the murine myeloma cell line, NS0) which are glutamine synthase negative. Glutamine synthase expression systems can also function in glutamine synthase expressing cells (e.g., Chinese Hamster Ovary (CHO) cells) by providing additional inhibitor to prevent the functioning of the endogenous gene. A glutamine synthase expression system and components thereof are detailed in PCT publications: WO87/04462; WO86/05807; WO89/01036; WO89/10404; and WO91/06657 which are hereby incorporated in their entireties by reference herein. Additionally, glutamine synthase expression vectors can be obtained from Lonza Biologics, Inc. (Portsmouth, NH). Expression and production of monoclonal antibodies using a GS expression system in murine myeloma cells is described in Bebbington *et al.*, *Bio/technology* 10:169(1992) and in Biblia and Robinson *Biotechnol. Prog.* 11:1 (1995) which are herein incorporated by reference.

[00184] The present invention also relates to host cells containing the above-described vector constructs described herein, and additionally encompasses host cells containing nucleotide sequences of the invention that are operably associated with one or more heterologous control regions (e.g., promoter and/or enhancer) using techniques known of in the art. The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. A host strain may be chosen, which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide may be controlled. Furthermore, different host cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the foreign protein expressed.

[00185] Introduction of the nucleic acids and nucleic acid constructs of the invention into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis et al., *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

[00186] In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., cardiovascular system antigen coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with cardiovascular system associated polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous cardiovascular system associated polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous cardiovascular system associated polynucleotide sequences via homologous recombination (see, e.g., US Patent Number 5,641,670, issued June 24, 1997; International Publication Number WO 96/29411; International Publication Number WO 94/12650; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature* 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

[00187] Polypeptides of the present invention can also be recovered from: products purified from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the



translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

[00188] In one embodiment, the yeast *Pichia pastoris* is used to express polypeptides of the invention in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A main step in the methanol metabolization pathway is the oxidation of methanol to formaldehyde using O<sub>2</sub>. This reaction is catalyzed by the enzyme alcohol oxidase. In order to metabolize methanol as its sole carbon source, *Pichia pastoris* must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O<sub>2</sub>. Consequently, in a growth medium depending on methanol as a main carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1* gene comprises up to approximately 30% of the total soluble protein in *Pichia pastoris*. See, Ellis, S.B., *et al.*, *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J., *et al.*, *Yeast* 5:167-77 (1989); Tschopp, J.F., *et al.*, *Nucl. Acids Res.* 15:3859-76 (1987). Thus, a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

[00189] In one example, the plasmid vector pPIC9K is used to express DNA encoding a polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press, Totowa, NJ, 1998. This expression vector allows expression and secretion of a polypeptide of the invention by virtue of the strong *AOX1* promoter linked to the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal peptide (i.e., leader) located upstream of a multiple cloning site.

[00190] Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5,

pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815, as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

[00191] In another embodiment, high-level expression of a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an expression vector such as, for example, pGAPZ or pGAPZalpha, and growing the yeast culture in the absence of methanol.

[00192] In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

[00193] In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., and Hunkapiller et al., Nature, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a polypeptide can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, alpha-amino isobutyric acid, 4-aminobutyric acid, Abu,

2-amino butyric acid, g-Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoro-amino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

[00194] The invention encompasses polypeptides of the present invention which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

[00195] Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

[00196] Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include iodine (<sup>121</sup>I, <sup>123</sup>I, <sup>125</sup>I, <sup>131</sup>I), carbon (<sup>14</sup>C), sulfur (<sup>35</sup>S), tritium (<sup>3</sup>H), indium (<sup>111</sup>In, <sup>112</sup>In, <sup>113m</sup>In, <sup>115m</sup>In), technetium (<sup>99</sup>Tc, <sup>99m</sup>Tc), thallium (<sup>201</sup>Tl), gallium (<sup>68</sup>Ga, <sup>67</sup>Ga), palladium (<sup>103</sup>Pd), molybdenum

(<sup>99</sup>Mo), xenon (<sup>133</sup>Xe), fluorine (<sup>18</sup>F), <sup>153</sup>Sm, <sup>177</sup>Lu, <sup>159</sup>Gd, <sup>149</sup>Pm, <sup>140</sup>La, <sup>175</sup>Yb, <sup>166</sup>Ho, <sup>90</sup>Y, <sup>47</sup>Sc, <sup>186</sup>Re, <sup>188</sup>Re, <sup>142</sup>Pr, <sup>105</sup>Rh, and <sup>97</sup>Ru.

[00197] In specific embodiments, a polypeptide of the present invention or fragment or variant thereof is attached to macrocyclic chelators that associate with radiometal ions, including but not limited to, <sup>177</sup>Lu, <sup>90</sup>Y, <sup>166</sup>Ho, and <sup>153</sup>Sm, to polypeptides. In a preferred embodiment, the radiometal ion associated with the macrocyclic chelators is <sup>111</sup>In. In another preferred embodiment, the radiometal ion associated with the macrocyclic chelator is <sup>90</sup>Y. In specific embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA). In other specific embodiments, DOTA is attached to an antibody of the invention or fragment thereof via a linker molecule. Examples of linker molecules useful for conjugating DOTA to a polypeptide are commonly known in the art - see, for example, DeNardo et al., Clin Cancer Res. 4(10):2483-90 (1998); Peterson et al., Bioconjug. Chem. 10(4):553-7 (1999); and Zimmerman et al, Nucl. Med. Biol. 26(8):943-50 (1999); which are hereby incorporated by reference in their entirety.

[00198] As mentioned, the cardiovascular system associated proteins of the invention may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given cardiovascular system associated polypeptide. Cardiovascular system associated polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic cardiovascular system associated polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA

mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth. Enzymol. 182:626-646 (1990); Rattan et al., Ann. N.Y. Acad. Sci. 663:48-62 (1992)).

[00199] Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

[00200] The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an average molecular weight of about 200, 500, 1000, 1500, 2000, 2500, 3000, 3500, 4000, 4500, 5000, 5500, 6000, 6500, 7000, 7500, 8000, 8500, 9000, 9500, 10,000, 10,500, 11,000, 11,500, 12,000, 12,500, 13,000, 13,500, 14,000, 14,500, 15,000, 15,500, 16,000, 16,500, 17,000, 17,500, 18,000, 18,500, 19,000, 19,500, 20,000, 25,000, 30,000, 35,000, 40,000, 50,000, 55,000, 60,000, 65,000, 70,000, 75,000, 80,000, 85,000, 90,000, 95,000, or 100,000 kDa.

[00201] As noted above, the polyethylene glycol may have a branched structure. Branched polyethylene glycols are described, for example, in U.S. Patent No.

5,643,575; Morpurgo et al., Appl. Biochem. Biotechnol. 56:59-72 (1996); Vorobjev et al., Nucleosides Nucleotides 18:2745-2750 (1999); and Caliceti et al., Bioconjug. Chem. 10:638-646 (1999), the disclosures of each of which are incorporated herein by reference.

**[00202]** The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, such as, for example, the method disclosed in EP 0 401 384 (coupling PEG to G-CSF), herein incorporated by reference; see also Malik et al., Exp. Hematol. 20:1028-1035 (1992), reporting pegylation of GM-CSF using tresyl chloride. For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

**[00203]** As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to proteins via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the protein or to more than one type of amino acid residue (e.g., lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein.

**[00204]** One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein.

The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

[00205] As indicated above, pegylation of the proteins of the invention may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein either directly or by an intervening linker. Linkerless systems for attaching polyethylene glycol to proteins are described in Delgado et al., Crit. Rev. Thera. Drug Carrier Sys. 9:249-304 (1992); Francis et al., Intern. J. of Hematol. 68:1-18 (1998); U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466, the disclosures of each of which are incorporated herein by reference.

[00206] One system for attaching polyethylene glycol directly to amino acid residues of proteins without an intervening linker employs tresylated MPEG, which is produced by the modification of monmethoxy polyethylene glycol (MPEG) using tresylchloride ( $\text{ClSO}_2\text{CH}_2\text{CF}_3$ ). Upon reaction of protein with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein. Thus, the invention includes protein-polyethylene glycol conjugates produced by reacting proteins of the invention with a polyethylene glycol molecule having a 2,2,2-trifluoroethane sulphonyl group.

[00207] Polyethylene glycol can also be attached to proteins using a number of different intervening linkers. For example, U.S. Patent No. 5,612,460, the entire disclosure of which is incorporated herein by reference, discloses urethane linkers for connecting polyethylene glycol to proteins. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the protein by a linker can also be produced by reaction of proteins with compounds such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG-p-nitrophenolcarbonate, and various MPEG-succinate derivatives. A

number of additional polyethylene glycol derivatives and reaction chemistries for attaching polyethylene glycol to proteins are described in International Publication No. WO 98/32466, the entire disclosure of which is incorporated herein by reference. Pegylated protein products produced using the reaction chemistries set out herein are included within the scope of the invention.

**[00208]** The number of polyethylene glycol moieties attached to each protein of the invention (i.e., the degree of substitution) may also vary. For example, the pegylated proteins of the invention may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado et al., Crit. Rev. Thera. Drug Carrier Sys. 9:249-304 (1992).

**[00209]** The cardiovascular system associated polypeptides of the invention can be recovered and purified from chemical synthesis and recombinant cell cultures by standard methods which include, but are not limited to, ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification. Well known techniques for refolding protein may be employed to regenerate active conformation when the polypeptide is denatured during isolation and/or purification.

**[00210]** Cardiovascular system associated polynucleotides and polypeptides may be used in accordance with the present invention for a variety of applications, particularly those that make use of the chemical and biological properties of cardiovascular system associated antigens. Among these are applications in the detection, prevention, diagnosis and/or treatment of diseases associated with cardiovascular system, such as e.g., cardiovascular system cancer, tumors, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease (e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease



(e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or those disorders or diseases as described under "Cardiovascular Disorders". Additional applications relate to diagnosis and to treatment of disorders of cells, tissues and organisms. These aspects of the invention are discussed further below.

[00211] In a preferred embodiment, polynucleotides expressed in a particular tissue type are used to detect, diagnose, treat, prevent and/or prognose disorders associated with the tissue type.

[00212] The polypeptides of the invention may be in monomers or multimers (i.e., dimers, trimers, tetramers and higher multimers). Accordingly, the present invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, Therapeutics) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

[00213] Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer refers to a multimer containing only polypeptides corresponding to a protein of the invention (e.g., the amino acid sequence of SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X or the complement of SEQ ID NO:X, the amino acid sequence encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or an amino acid sequence encoded by cDNA contained in Clone ID NO:Z (including fragments, variants, splice variants, and fusion proteins, corresponding to these as described herein)). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing two polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing three polypeptides having identical and/or different amino acid sequences). In additional embodiments,

the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

[00214] As used herein, the term heteromer refers to a multimer containing two or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

[00215] Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked by, for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in SEQ ID NO:Y, encoded by the portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or encoded by the cDNA contained in Clone ID NO:Z). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein. In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., U.S. Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in a Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous

polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, osteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

[00216] Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

[00217] Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

[00218] In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, proteins of the

invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

[00219] The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C-terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide components desired to be contained in the multimer of the invention (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

[00220] Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant

polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., U.S. Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

### **Antibodies**

[00221] Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of the invention (e.g., a polypeptide or fragment or variant of the amino acid sequence of SEQ ID NO:Y or a polypeptide encoded by the cDNA contained in Clone ID NO:Z, and/or an epitope, of the present invention) as determined by immunoassays well known in the art for assaying specific antibody-antigen binding. Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), intracellularly-made antibodies (i.e., intrabodies), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule. In preferred embodiments, the immunoglobulin molecules of the invention are IgG1. In other preferred embodiments, the immunoglobulin molecules of the invention are IgG4.

[00222] Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')<sub>2</sub>, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding

fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, ship rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described *infra* and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

[00223] The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

[00224] Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, or by size in contiguous amino acid residues, or listed in the Tables and Figures. Preferred epitopes of the invention include those shown in column 6 of Table 1A, as well as polynucleotides that encode these epitopes. Antibodies, which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

[00225] Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least

75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies, which bind polypeptides encoded by polynucleotides, which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-2}$  M,  $10^{-2}$  M,  $5 \times 10^{-3}$  M,  $10^{-3}$  M,  $5 \times 10^{-4}$  M,  $10^{-4}$  M,  $5 \times 10^{-5}$  M,  $10^{-5}$  M,  $5 \times 10^{-6}$  M,  $10^{-6}$  M,  $5 \times 10^{-7}$  M,  $10^{-7}$  M,  $5 \times 10^{-8}$  M,  $10^{-8}$  M,  $5 \times 10^{-9}$  M,  $10^{-9}$  M,  $5 \times 10^{-10}$  M,  $10^{-10}$  M,  $5 \times 10^{-11}$  M,  $10^{-11}$  M,  $5 \times 10^{-12}$  M,  $10^{-12}$  M,  $5 \times 10^{-13}$  M,  $10^{-13}$  M,  $5 \times 10^{-14}$  M,  $10^{-14}$  M,  $5 \times 10^{-15}$  M, or  $10^{-15}$  M.

[00226] The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

[00227] Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies, which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention

features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies, which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described *supra*). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

[00228] The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies, which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., Blood 92(6):1981-1988 (1998); Chen et al., Cancer Res. 58(16):3668-3678 (1998); Harrop et al., J. Immunol. 161(4):1786-1794 (1998); Zhu et al., Cancer Res. 58(15):3209-3214 (1998); Yoon et al., J. Immunol. 160(7):3170-3179 (1998); Prat et al., J. Cell. Sci. 111(Pt2):237-247 (1998); Pitard et al., J. Immunol. Methods 205(2):177-190 (1997); Liautard et al., Cytokine 9(4):233-241 (1997); Carlson et al., J. Biol. Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller



et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

[00229] Antibodies of the present invention may be used, for example, to purify, detect, and target the polypeptides of the present invention, including both *in vitro* and *in vivo* diagnostic and therapeutic methods. For example, the antibodies have utility in immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the present invention in biological samples. See, e.g., Harlow et al., Antibodies: A Laboratory Manual, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); incorporated by reference herein in its entirety.

[00230] As discussed in more detail below, the antibodies of the present invention may be used either alone or in combination with other compositions. The antibodies may further be recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically conjugated (including covalent and non-covalent conjugations) to polypeptides or other compositions. For example, antibodies of the present invention may be recombinantly fused or conjugated to molecules useful as labels in detection assays and effector molecules such as heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO 92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387; the disclosures of which are incorporated herein by reference in their entireties.

[00231] The antibodies of the invention include derivatives that are modified, i.e., by the covalent attachment of any type of molecule to the antibody such that covalent attachment does not prevent the antibody from generating an anti-idiotypic response. For example, but not by way of limitation, the antibody derivatives include antibodies that have been modified, e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other protein, etc. Any of numerous chemical modifications may be carried out by known techniques, including, but not limited to specific chemical cleavage, acetylation, formylation, metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or more non-classical amino acids.

[00232] The antibodies of the present invention may be generated by any suitable method known in the art. Polyclonal antibodies to an antigen-of-interest can be produced by various procedures well known in the art. For example, a polypeptide

of the invention can be administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

[00233] Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

[00234] Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art and are discussed in detail in the Examples. In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the

invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

[00235] Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

[00236] Another well known method for producing both polyclonal and monoclonal human B cell lines is transformation using Epstein Barr Virus (EBV). Protocols for generating EBV-transformed B cell lines are commonly known in the art, such as, for example, the protocol outlined in Chapter 7.22 of Current Protocols in Immunology, Coligan et al., Eds., 1994, John Wiley & Sons, NY, which is hereby incorporated in its entirety by reference herein. The source of B cells for transformation is commonly human peripheral blood, but B cells for transformation may also be derived from other sources including, but not limited to, lymph nodes, tonsil, spleen, tumor tissue, and infected tissues. Tissues are generally made into single cell suspensions prior to EBV transformation. Additionally, steps may be taken to either physically remove or inactivate T cells (e.g., by treatment with cyclosporin A) in B cell-containing samples, because T cells from individuals seropositive for anti-EBV antibodies can suppress B cell immortalization by EBV.

[00237] In general, the sample containing human B cells is inoculated with EBV, and cultured for 3-4 weeks. A typical source of EBV is the culture supernatant of the B95-8 cell line (ATCC #VR-1492). Physical signs of EBV transformation can generally be seen towards the end of the 3-4 week culture period. By phase-contrast microscopy, transformed cells may appear large, clear, hairy and tend to aggregate in tight clusters of cells. Initially, EBV lines are generally polyclonal. However, over prolonged periods of cell cultures, EBV lines may become monoclonal or polyclonal as a result of the selective outgrowth of particular B cell clones. Alternatively, polyclonal EBV transformed lines may be subcloned (e.g., by limiting dilution culture) or fused with a suitable fusion partner and plated at limiting dilution to obtain monoclonal B cell lines. Suitable fusion partners for EBV

transformed cell lines include mouse myeloma cell lines (e.g., SP2/0, X63-Ag8.653), heteromyeloma cell lines (human x mouse; e.g., SPAM-8, SBC-H20, and CB-F7), and human cell lines (e.g., GM 1500, SKO-007, RPMI 8226, and KR-4). Thus, the present invention also provides a method of generating polyclonal or monoclonal human antibodies against polypeptides of the invention or fragments thereof, comprising EBV-transformation of human B cells.

[00238] Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')<sub>2</sub> fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')<sub>2</sub> fragments). F(ab')<sub>2</sub> fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain. For example, the antibodies of the present invention can also be generated using various phage display methods known in the art and as discussed in detail in the Examples (e.g., Example 10). In phage display methods, functional antibody domains are displayed on the surface of phage particles, which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047;

5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

[00239] As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')<sub>2</sub> fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., BioTechniques 12(6):864-869 (1992); and Sawai et al., AJRI 34:26-34 (1995); and Better et al., Science 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

[00240] Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., Methods in Enzymology 203:46-88 (1991); Shu et al., PNAS 90:7995-7999 (1993); and Skerra et al., Science 240:1038-1040 (1988). For some uses, including *in vivo* use of antibodies in humans and *in vitro* detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Gillies et al., (1989) J. Immunol. Methods 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues

important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., Nature 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, Molecular Immunology 28(4/5):489-498 (1991); Studnicka et al., Protein Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

**[00241]** Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

**[00242]** Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring, which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from

the immunized, transgenic mice using conventional hybridoma technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, *Int. Rev. Immunol.* 13:65-93 (1995). For a detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; 5,939,598; 6,075,181 and 6,114,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above.

**[00243]** Completely human antibodies, which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., *Bio/technology* 12:899-903 (1988)).

**[00244]** Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, *FASEB J.* 7(5):437-444; (1989) and Nissinoff, J. *Immunol.* 147(8):2429-2438 (1991)). For example, antibodies which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand/receptor. For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligand(s)/receptor(s), and thereby block its biological activity. Alternatively, antibodies, which bind to and

enhance polypeptide multimerization and/or binding, and/or receptor/ligand multimerization, binding and/or signaling can be used to generate anti-idiotypes that function as agonists of a polypeptide of the invention and/or its ligand/receptor. Such agonistic anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens as agonists of the polypeptides of the invention or its ligand(s)/receptor(s). For example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligand(s)/receptor(s), and thereby promote or enhance its biological activity.

- [00245] Intrabodies of the invention can be produced using methods known in the art, such as those disclosed and reviewed in Chen et al., Hum. Gene Ther. 5:595-601 (1994); Marasco, W.A., Gene Ther. 4:11-15 (1997); Rondon and Marasco, Annu. Rev. Microbiol. 51:257-283 (1997); Proba et al., J. Mol. Biol. 275:245-253 (1998); Cohen et al., Oncogene 17:2445-2456 (1998); Ohage and Steipe, J. Mol. Biol. 291:1119-1128 (1999); Ohage et al., J. Mol. Biol. 291:1129-1134 (1999); Wirtz and Steipe, Protein Sci. 8:2245-2250 (1999); Zhu et al., J. Immunol. Methods 231:207-222 (1999); and references cited therein.

#### *Polynucleotides Encoding Antibodies*

- [00246] The invention further provides polynucleotides comprising a nucleotide sequence encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or alternatively, under lower stringency hybridization conditions, e.g., as defined *supra*, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y, to a polypeptide encoded by a portion of SEQ ID NO:X as defined in columns 8 and 9 of Table 2, and/or to a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

- [00247] The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., BioTechniques 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence



encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

[00248] Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

[00249] Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, Molecular Cloning, A Laboratory Manual, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, Current Protocols in Molecular Biology, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

[00250] In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art, e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework regions to humanize a non-human antibody, as described *supra*. The framework regions may be naturally

10091504, 030702

occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed *supra*, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

**[00251]** In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described *supra*, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

**[00252]** Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in *E. coli* may also be used (Skerra et al., Science 242:1038-1041 (1988)).

#### *Methods of Producing Antibodies*

**[00253]** The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by

recombinant expression techniques. Methods of producing antibodies include, but are not limited to, hybridoma technology, EBV transformation, and other methods discussed herein as well as through the use recombinant DNA technology, as discussed below.

[00254] Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein. Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, *in vitro* recombinant DNA techniques, synthetic techniques, and *in vivo* genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

[00255] The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-

expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

[00256] A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene* 45:101 (1986); Cockett et al., *Bio/Technology* 8:2 (1990)).

[00257] In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily

purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., EMBO J. 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, Nucleic Acids Res. 13:3101-3109 (1985); Van Heeke & Schuster, J. Biol. Chem. 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[00258] In an insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

[00259] In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk, Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of

appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., *Methods in Enzymol.* 153:51-544 (1987)).

**[00260]** In addition, a host cell strain may be chosen, which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells, which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS, MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

**[00261]** For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines, which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines, which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

**[00262]** A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223 (1977)),

hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., Cell 22:817 (1980)) genes can be employed in tk-, hgp<sup>r</sup>t- or ap<sup>r</sup>t- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., Natl. Acad. Sci. USA 77:357 (1980); O'Hare et al., Proc. Natl. Acad. Sci. USA 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 Clinical Pharmacy 12:488-505; Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); TIB TECH 11(5):155-215 (1993)); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre et al., Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., J. Mol. Biol. 150:1 (1981), which are incorporated by reference herein in their entireties.

**[00263]** The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

**[00264]** Vectors, which use glutamine synthase (GS) or DHFR as the selectable markers, can be amplified in the presence of the drugs methionine sulfoximine or methotrexate, respectively. An advantage of glutamine synthase based vectors are the availability of cell lines (e.g., the murine myeloma cell line, NS0) which are

glutamine synthase negative. Glutamine synthase expression systems can also function in glutamine synthase expressing cells (e.g., Chinese Hamster Ovary (CHO) cells) by providing additional inhibitor to prevent the functioning of the endogenous gene. A glutamine synthase expression system and components thereof are detailed in PCT publications: WO87/04462; WO86/05807; WO89/01036; WO89/10404; and WO91/06657 which are incorporated in their entireties by reference herein. Additionally, glutamine synthase expression vectors that may be used according to the present invention are commercially available from suppliers, including, for example Lonza Biologics, Inc. (Portsmouth, NH). Expression and production of monoclonal antibodies using a GS expression system in murine myeloma cells is described in Bebbington et al., *Bio/technology* 10:169(1992) and in Biblia and Robinson *Biotechnol. Prog.* 11:1 (1995) which are incorporated in their entireties by reference herein.

[00265] The host cell may be co-transfected with two expression vectors of the invention, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers, which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to avoid an excess of toxic free heavy chain (Proudfoot, *Nature* 322:52 (1986); Kohler, *Proc. Natl. Acad. Sci. USA* 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

[00266] Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide sequences described herein or otherwise known in the art, to facilitate purification.



10091504-030702

[00267] The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either *in vitro* or *in vivo*, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in *in vitro* immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., *supra*, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452 (1991), which are incorporated by reference in their entireties.

[00268] The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc.

Natl. Acad. Sci. USA 89:11337- 11341 (1992) (said references incorporated by reference in their entireties).

[00269] As discussed, *supra*, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the *in vivo* half life of the polypeptides or for use in immunoassays using methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See EP 394,827; Traunecker et al., Nature 331:84-86 (1988). The polypeptides of the present invention fused or conjugated to an antibody having disulfide- linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. See, for example, Fountoulakis et al., J. Biochem. 270:3958-3964 (1995). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. See, for example, EP A 232,262. Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., J. Molecular Recognition 8:52-58 (1995); Johanson et al., J. Biol. Chem. 270:9459-9471 (1995)).

[00270] Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., Proc. Natl. Acad. Sci. USA 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA"

tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., Cell 37:767 (1984)) and the "flag" tag.

[00271] The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention.

[00272] Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, <sup>213</sup>Bi. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis- dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly

actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[00273] The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor,  $\alpha$ -interferon,  $\beta$ -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- $\alpha$ , TNF- $\beta$ , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGI (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

[00274] Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

[00275] Techniques for conjugating such therapeutic moiety to antibodies are well known. See, for example., Arnon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16

(Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev. 62:119-58 (1982).

[00276] Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

[00277] An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

#### *Immunophenotyping*

[00278] The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. Translation products of the genes of the present invention may be useful as cell specific markers, or more specifically as cellular markers that are differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison et al., Cell, 96:737-49 (1999)).

[00279] These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might be found in human umbilical cord blood.

#### *Assays For Antibody Binding*

[00280] The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited to competitive and non-competitive assay systems using techniques such

as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, and protein A immunoassays, to name but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

**[00281]** Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X- 100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasylol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al., eds., (1994), Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York, section 10.16.1.

**[00282]** Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%- 20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a

secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g.,  $^{32}\text{P}$  or  $^{125}\text{I}$ ) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al., eds., (1994), *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, section 10.8.1.

**[00283]** ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound; instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al., eds., (1994), *Current Protocols in Molecular Biology*, Vol. 1, John Wiley & Sons, Inc., New York, section 11.2.1.

**[00284]** The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g.,  $^3\text{H}$  or  $^{125}\text{I}$ ) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest

conjugated to a labeled compound (e.g.,  $^3\text{H}$  or  $^{125}\text{I}$ ) in the presence of increasing amounts of an unlabeled second antibody.

[00285] Antibodies of the invention may be characterized using immunocytochemistry methods on cells (e.g., mammalian cells, such as CHO cells) transfected with a vector enabling the expression of a cardiovascular system antigen or with vector alone using techniques commonly known in the art. Antibodies that bind cardiovascular system antigen transfected cells, but not vector-only transfected cells, are cardiovascular system antigen specific.

#### *Therapeutic Uses*

[00286] The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

[00287] In a specific and preferred embodiment, the present invention is directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the diseases, disorders, or conditions of the cardiovascular system, including, but not limited to, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease



(e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease (e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or those disorders or diseases as described under "Cardiovascular Disorders". Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (e.g., antibodies directed to the full length protein expressed on the cell surface of a mammalian cell; antibodies directed to an epitope of a cardiovascular system associated polypeptide of the invention (such as, a linear epitope (shown in Table 1A, column 6) or a conformational epitope), including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions of the cardiovascular system described herein. The treatment and/or prevention of diseases, disorders, or conditions of the cardiovascular system associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

**[00288]** A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

**[00289]** The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or

hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

[00290] The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

[00291] It is preferred to use high affinity and/or potent *in vivo* inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-2}$  M,  $10^{-2}$  M,  $5 \times 10^{-3}$  M,  $10^{-3}$  M,  $5 \times 10^{-4}$  M,  $10^{-4}$  M,  $5 \times 10^{-5}$  M,  $10^{-5}$  M,  $5 \times 10^{-6}$  M,  $10^{-6}$  M,  $5 \times 10^{-7}$  M,  $10^{-7}$  M,  $5 \times 10^{-8}$  M,  $10^{-8}$  M,  $5 \times 10^{-9}$  M,  $10^{-9}$  M,  $5 \times 10^{-10}$  M,  $10^{-10}$  M,  $5 \times 10^{-11}$  M,  $10^{-11}$  M,  $5 \times 10^{-12}$  M,  $10^{-12}$  M,  $5 \times 10^{-13}$  M,  $10^{-13}$  M,  $5 \times 10^{-14}$  M,  $10^{-14}$  M,  $5 \times 10^{-15}$  M, and  $10^{-15}$  M.

#### *Gene Therapy*

[00292] In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

[00293] Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

10091504-030702

[00294] For general reviews of the methods of gene therapy, see Goldspiel et al., Clinical Pharmacy 12:488-505 (1993); Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, TIBTECH 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); and Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990).

[00295] In a preferred embodiment, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989). In specific embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

[00296] Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

[00297] In a specific embodiment, the nucleic acid sequences are directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated

retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989)).

[00298] In a specific embodiment, viral vectors that contain nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., Meth. Enzymol. 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., Biotherapy 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., J. Clin. Invest. 93:644-651 (1994); Kiem et al., Blood 83:1467-1473 (1994); Salmons and Gunzberg, Human Gene Therapy 4:129-141 (1993); and Grossman and Wilson, Curr. Opin. in Genetics and Devel. 3:110-114 (1993).

[00299] Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory

epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., *Gene Therapy* 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

[00300] Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., *Proc. Soc. Exp. Biol. Med.* 204:289-300 (1993); U.S. Patent No. 5,436,146).

[00301] Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

[00302] In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes into cells (see, e.g., Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so

that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

[00303] The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

[00304] Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to, epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

[00305] In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

[00306] In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible by the cells or their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, Cell 71:973-985 (1992); Rheinwald, Meth. Cell Bio. 21A:229 (1980); and Pittelkow and Scott, Mayo Clinic Proc. 61:771 (1986)).

[00307] In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by the presence or absence of an appropriate inducer of transcription.

#### *Demonstration of Therapeutic or Prophylactic Activity*

[00308] The compounds or pharmaceutical compositions of the invention are preferably tested *in vitro*, and then *in vivo* for the desired therapeutic or prophylactic activity,

prior to use in humans. For example, *in vitro* assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance with the invention, *in vitro* assays which can be used to determine whether administration of a specific compound is indicated, include *in vitro* cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

#### *Therapeutic/Prophylactic Administration and Composition*

- [00309] The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably a polypeptide or antibody of the invention. In a preferred embodiment, the compound is substantially purified (e.g., substantially free from substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.
- [00310] Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional appropriate formulations and routes of administration can be selected from among those described herein below.
- [00311] Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)), construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or

mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

[00312] In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

[00313] In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

[00314] In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, polymeric materials can be used (see Medical Applications of Controlled Release, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); Controlled Drug Bioavailability, Drug Product Design and Performance, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., Macromol. Sci. Rev. Macromol. Chem. 23:61 (1983); see also Levy et al., Science 228:190 (1985); During et al., Ann. Neurol. 25:351 (1989); Howard et al.,



J.Neurosurg. 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, e.g., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).

[00315] Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

[00316] In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered *in vivo* to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox- like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

[00317] The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like.

10091504-030702

The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[00318] In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[00319] The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

20/05/04 10:03:00

[00320] The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[00321] For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

[00322] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

#### *Diagnosis and Imaging*

[00323] Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor diseases, disorders, and/or conditions associated with the aberrant expression and/or activity of a polypeptide of the invention. The invention provides

for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of aberrant expression.

[00324] The invention provides a diagnostic assay for diagnosing a cardiovascular system disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

[00325] Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen et al., J. Cell . Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (125I, 121I), carbon (14C), sulfur (35S), tritium (3H), indium (112In), and technetium (99Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00326] One facet of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. A preferred embodiment of the

invention is the detection and diagnosis of a disease or disorder of the cardiovascular system associated with aberrant expression of a cardiovascular system antigen in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the amount of labeled molecule detected to a standard value previously determined for a particular system.

[00327] It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99m}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[00328] Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

[00329] In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disorder, for example, one month

after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

[00330] Presence of the labeled molecule can be detected in the patient using methods known in the art for *in vivo* scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as position emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

[00331] In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI).

#### Kits

[00332] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope, which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present invention further comprise a control antibody, which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate).

10091504-030703

[00333] In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

[00334] In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

[00335] In an additional embodiment, the invention includes a diagnostic kit for use in screening serum-containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

[00336] In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is

determined. Typically, the reporter is an enzyme, which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

[00337] The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

[00338] Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

#### **Uses of the Polynucleotides**

[00339] Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

[00340] The polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome, thus each polynucleotide of the present invention can routinely be used as a chromosome marker using techniques known in the art. Table 1A, column 8 provides the chromosome location of some of the polynucleotides of the invention.

[00341] Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably at least 15 bp (e.g., 15-25 bp) from the sequences shown in SEQ ID NO:X. Primers can optionally be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are



then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to SEQ ID NO:X will yield an amplified fragment.

[00342] Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted chromosomes, preselection by hybridization to construct chromosome specific-cDNA libraries, and computer mapping techniques (See, e.g., Shuler, Trends Biotechnol 16:456-459 (1998) which is hereby incorporated by reference in its entirety).

[00343] Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

[00344] For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple sites and/or multiple chromosomes).

[00345] Thus, the present invention also provides a method for chromosomal localization which involves (a) preparing PCR primers from the polynucleotide sequences in Table 1A and/or Table 2 and SEQ ID NO:X and (b) screening somatic cell hybrids containing individual chromosomes.

[00346] The polynucleotides of the present invention would likewise be useful for radiation hybrid mapping, HAPPY mapping, and long range restriction mapping. For a review of these techniques and others known in the art, see, e.g. Dear, "Genome Mapping: A Practical Approach," IRL Press at Oxford University Press, London (1997); Aydin, J. Mol. Med. 77:691-694 (1999); Hacia et al., Mol. Psychiatry 3:483-492 (1998); Herrick et al., Chromosome Res. 7:409-423 (1999); Hamilton et al., Methods Cell Biol. 62:265-280 (2000); and/or Ott, J. Hered. 90:68-70 (1999), each of which is hereby incorporated by reference in its entirety.

10091504-030702

[00347] Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library).) Column 9 of Table 1A provides an OMIM reference identification number of diseases associated with the cytologic band disclosed in column 8 of Table 1A, as determined using techniques described herein and by reference to Table 5. Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

[00348] Thus, once coinheritance is established, differences in a polynucleotide of the invention and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all affected individuals, but not in normal individuals, indicate that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

[00349] Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using the polynucleotides of the invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker. Diagnostic and prognostic methods, kits and reagents encompassed by the present invention are briefly described below and more thoroughly elsewhere herein (see e.g., the sections labeled "Antibodies", "Diagnostic Assays", and "Methods for Detecting Cardiovascular System Disease, Including Cancer").

[00350] Thus, the invention also provides a diagnostic method useful during diagnosis of a disorder, involving measuring the expression level of polynucleotides of the present invention in cells or body fluid from an individual and comparing the

measured gene expression level with a standard level of polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a disorder. Additional non-limiting examples of diagnostic methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., Example 12).

[00351] In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test subject, as further described herein. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes defining an internal region of the polynucleotide of the invention, where each probe has one strand containing a 31'-mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

[00352] Where a diagnosis of a related disorder, including, for example, diagnosis of a tumor, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed polynucleotide of the invention expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

[00353] By "measuring the expression level of polynucleotides of the invention" is intended qualitatively or quantitatively measuring or estimating the level of the polypeptide of the invention or the level of the mRNA encoding the polypeptide of the invention in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the polypeptide level or mRNA level in a second biological sample). Preferably, the polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the related disorder or being determined by averaging levels from a population of individuals not having a related disorder. As will be appreciated in

the art, once a standard polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

[00354] By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains polypeptide of the present invention or the corresponding mRNA. As indicated, biological samples include body fluids (such as semen, lymph, vaginal pool, sera, plasma, urine, synovial fluid and spinal fluid) which contain the polypeptide of the present invention, and tissue sources found to express the polypeptide of the present invention. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

[00355] The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides of the invention are attached to a solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in U.S. Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene chip with polynucleotides of the invention attached may be used to identify polymorphisms between the isolated polynucleotide sequences of the invention, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms (i.e., their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, such as for example, in neural disorders, immune system disorders, muscular disorders, reproductive disorders, gastrointestinal disorders, pulmonary disorders, digestive disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions. Such a method is described in U.S. Patents 5,858,659 and 5,856,104. The U.S. Patents referenced *supra* are hereby incorporated by reference in their entirety herein.

[00356] The present invention encompasses polynucleotides of the present invention that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides of the invention are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive

Biosystems). Certain components of DNA, such as phosphorus, phosphorus oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by Nielsen et al., Science 254:1497 (1991); and Egholm et al., Nature 365:666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ( $T_{sub.m}$ ) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

[00357] The compounds of the present invention have uses, which include, but are not limited to, detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

[00358] The compounds of the present invention have preferred uses, which include, but are not limited to, detecting cardiovascular system cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias, which include, but are not limited to: myxomas, fibromas, and rhabdomyomas. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

[00359] Pathological cell proliferative disorders are often associated with inappropriate activation of proto-oncogenes. (Gelman, E. P. et al., "The Etiology of Acute

Leukemia: Molecular Genetics and Viral Oncology," in Neoplastic Diseases of the Blood, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now believed to result from the qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Germann et al., *supra*) It is likely that mutated or altered expression of specific genes is involved in the pathogenesis of some leukemias, among other tissues and cell types. (Germann et al., *supra*) Indeed, the human counterparts of the oncogenes involved in some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Germann et al., *supra*)

[00360] For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580). However, it has been shown that exposure of HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness is not be limited to treatment, prevention, diagnosis and/or prognosis, of proliferative disorders of cells and tissues of hematopoietic origin, in light of the numerous cells and cell types of varying origins which are known to exhibit proliferative phenotypes. In preferred embodiments, the compounds and/or methods of the invention are used to treat, prevent, diagnose, and/or prognose, proliferative disorders of cardiovascular system cells and tissues.

[00361] In addition to the foregoing, a polynucleotide of the present invention can be used to control gene expression through triple helix formation or through antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is

discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. The oligonucleotide described above can also be delivered to cells such that the antisense RNA or DNA may be expressed *in vivo* to inhibit production of polypeptide of the present invention antigens. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease, and in particular, for the treatment of proliferative diseases and/or conditions. Non-limiting antisense and triple helix methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., the section labeled "Antisense and Ribozyme (Antagonists)").

[00362] Polynucleotides of the present invention are also useful in gene therapy. One goal of gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell. Additional non-limiting examples of gene therapy methods encompassed by the present invention are more thoroughly described elsewhere herein (see, e.g., the sections labeled "Gene Therapy Methods" and Examples 16, 17 and 18).

[00363] The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or

more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

[00364] The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue samples.

[00365] Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues, e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

[00366] There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers prepared from the sequences of the present invention, specific to tissues, including but not limited to, those sequences referred to in Table 1A. Panels of such reagents can identify tissue by species and/or by organ type. In a similar fashion, these reagents can be used to screen tissue cultures for contamination. Additional non-limiting examples of such uses are further described herein.



10091504 030306

[00367] Because cardiovascular system antigens are found expressed in the cardiovascular system, the polynucleotides of the present invention are also useful as hybridization probes for differential identification of the tissue(s) or cell type(s) present in a biological sample. Similarly, polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g., immunohistochemistry assays) or cell type(s) (e.g., immunocytochemistry assays). In a specific embodiment, the polynucleotides of the present invention are also useful as hybridization probes for differential identification of cardiovascular system tissue(s) or cell type(s) present in a biological sample. Similarly, polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of cardiovascular system tissue(s) (e.g., immunohistochemistry assays) or cell type(s) (e.g., immunocytochemistry assays). In addition, for a number of disorders of the above tissues or cells, significantly higher or lower levels of gene expression of the polynucleotides/polypeptides of the present invention may be detected in certain tissues (e.g., tissues expressing polypeptides and/or polynucleotides of the present invention, for example, normal cardiovascular system or diseased cardiovascular system tissues, and/or those tissues/cells corresponding to the library source relating to a polynucleotide sequence of the invention as disclosed in column 7 of Table 1A, and/or cancerous and/or wounded tissues) or bodily fluids (e.g., semen, lymph, vaginal pool, serum, plasma, urine, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" gene expression level, i.e., the expression level in healthy tissue from an individual not having the disorder.

[00368] Thus, the invention provides a diagnostic method of a disorder, which involves: (a) assaying gene expression level in cells or body fluid of an individual; (b) comparing the gene expression level with a standard gene expression level, whereby an increase or decrease in the assayed gene expression level compared to the standard expression level is indicative of a disorder.

[00369] In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and

making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

### Uses of the Polypeptides

[00370] Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

[00371] Polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g., immunohistochemistry assays such as, for example, ABC immunoperoxidase (Hsu et al., J. Histochem. Cytochem. 29:577-580 (1981)) or cell type(s) (e.g., immunocytochemistry assays).

[00372] Antibodies can be used to assay levels of polypeptides encoded by polynucleotides of the invention in a biological sample using classical immunohistological methods known to those of skill in the art (see, e.g., Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine ( $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{115\text{m}}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{112}\text{In}$ ,  $^{111}\text{In}$ ), and technetium ( $^{99}\text{Tc}$ ,  $^{99\text{m}}\text{Tc}$ ), thallium ( $^{201}\text{Tl}$ ), gallium ( $^{68}\text{Ga}$ ,  $^{67}\text{Ga}$ ), palladium ( $^{103}\text{Pd}$ ), molybdenum ( $^{99}\text{Mo}$ ), xenon ( $^{133}\text{Xe}$ ), fluorine ( $^{18}\text{F}$ ),  $^{153}\text{Sm}$ ,  $^{177}\text{Lu}$ ,  $^{159}\text{Gd}$ ,  $^{149}\text{Pm}$ ,  $^{140}\text{La}$ ,  $^{175}\text{Yb}$ ,  $^{166}\text{Ho}$ ,  $^{90}\text{Y}$ ,  $^{47}\text{Sc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{142}\text{Pr}$ ,  $^{105}\text{Rh}$ ,  $^{97}\text{Ru}$ ; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00373] In addition to assaying levels of polypeptide of the present invention in a biological sample, proteins can also be detected *in vivo* by imaging. Antibody labels or markers for *in vivo* imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable

characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

[00374] A cardiovascular system antigen-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example,  $^{131}\text{I}$ ,  $^{112}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ , ( $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{115\text{m}}\text{In}$ ,  $^{113\text{m}}\text{In}$ ,  $^{112}\text{In}$ ,  $^{111}\text{In}$ ), and technetium ( $^{99}\text{Tc}$ ,  $^{99\text{m}}\text{Tc}$ ), thallium ( $^{201}\text{Tl}$ ), gallium ( $^{68}\text{Ga}$ ,  $^{67}\text{Ga}$ ), palladium ( $^{103}\text{Pd}$ ), molybdenum ( $^{99}\text{Mo}$ ), xenon ( $^{133}\text{Xe}$ ), fluorine ( $^{18}\text{F}$ ,  $^{153}\text{Sm}$ ,  $^{177}\text{Lu}$ ,  $^{159}\text{Gd}$ ,  $^{149}\text{Pm}$ ,  $^{140}\text{La}$ ,  $^{175}\text{Yb}$ ,  $^{166}\text{Ho}$ ,  $^{90}\text{Y}$ ,  $^{47}\text{Sc}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{142}\text{Pr}$ ,  $^{105}\text{Rh}$ ,  $^{97}\text{Ru}$ ), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for cardiovascular system disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99\text{m}}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which express the polypeptide encoded by a polynucleotide of the invention. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[00375] In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (e.g., polypeptides encoded by polynucleotides of the invention and/or antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

[00376] In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention in association with toxins or cytotoxic prodrugs.

[00377] In a preferred embodiment, the invention provides a method for the specific destruction of cardiovascular system cells (e.g., aberrant cardiovascular system cells, cardiovascular system neoplasm) by administering polypeptides of the invention (e.g., polypeptides encoded by polynucleotides of the invention and/or antibodies) in association with toxins or cytotoxic prodrugs. In another preferred embodiment the invention provides a method for the specific destruction of tissues/cells corresponding to the library source relating to a polynucleotide sequence of the invention as disclosed in column 7 of Table 1A by administering polypeptides of the invention in association with toxins or cytotoxic prodrugs.

[00378] By "toxin" is meant one or more compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. "Toxin" also includes a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example,  $^{213}\text{Bi}$ , or other radioisotopes such as, for example,  $^{103}\text{Pd}$ ,  $^{133}\text{Xe}$ ,  $^{131}\text{I}$ ,  $^{111}\text{In}$ ,  $^{68}\text{Ge}$ ,  $^{57}\text{Co}$ ,  $^{65}\text{Zn}$ ,  $^{85}\text{Sr}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{153}\text{Sm}$ ,  $^{153}\text{Gd}$ ,  $^{169}\text{Yb}$ ,  $^{51}\text{Cr}$ ,  $^{54}\text{Mn}$ ,  $^{75}\text{Se}$ ,  $^{113}\text{Sn}$ ,  $^{90}\text{Yttrium}$ ,  $^{117}\text{Tin}$ ,  $^{186}\text{Rhenium}$ ,  $^{166}\text{Holmium}$ , and  $^{188}\text{Rhenium}$ ; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00379] In a specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope  $^{90}\text{Y}$ . In another specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope  $^{111}\text{In}$ . In a further specific embodiment, the invention provides a method for the specific destruction of cells (e.g., the

destruction of tumor cells) by administering polypeptides of the invention or antibodies of the invention in association with the radioisotope <sup>131</sup>I.

[00380] Techniques known in the art may be applied to label polypeptides of the invention (including antibodies). Such techniques include, but are not limited to, the use of bifunctional conjugating agents (see e.g., U.S. Patent Nos. 5,756,065; 5,714,631; 5,696,239; 5,652,361; 5,505,931; 5,489,425; 5,435,990; 5,428,139; 5,342,604; 5,274,119; 4,994,560; and 5,808,003; the contents of each of which are hereby incorporated by reference in its entirety).

[00381] Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression level of a polypeptide of the present invention in cells or body fluid of an individual; and (b) comparing the assayed polypeptide expression level with a standard polypeptide expression level, whereby an increase or decrease in the assayed polypeptide expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

[00382] Moreover, polypeptides of the present invention can be used to treat or prevent diseases or conditions of the cardiovascular system such as, for example, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease (e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease (e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or those disorders or diseases as described under "Cardiovascular Disorders". In preferred embodiments, polynucleotides expressed in a particular tissue type (see, e.g., Table 1A, column 7) are used to diagnose, detect, prevent, treat and/or prognose disorders associated with the tissue type. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or

decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

[00383] Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease (as described *supra*, and elsewhere herein). For example, administration of an antibody directed to a polypeptide of the present invention can bind, and/or neutralize the polypeptide, and/or reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

[00384] At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the biological activities described herein.

#### **Diagnostic Assays**

[00385] The compounds of the present invention are useful for diagnosis, treatment, prevention and/or prognosis of various cardiovascular system related disorders in mammals, preferably humans. Such disorders include, but are not limited to, cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease (e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease (e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or those disorders or diseases as described under "Cardiovascular Disorders".

In preferred embodiments, polynucleotides expressed in a particular tissue type (see, e.g., Table 1A, column 7) are used to diagnose, detect, prevent, treat and/or prognose disorders associated with the tissue type.

[00386] Cardiovascular system antigens are expressed in the cardiovascular system , with an increased expression level in cardiovascular system tissues. For a number of cardiovascular system -related disorders, substantially altered (increased or decreased) levels of cardiovascular system antigen gene expression can be detected in cardiovascular system tissue or other cells or bodily fluids (e.g., sera, plasma, urine, semen, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" cardiovascular system antigen gene expression level, that is, the cardiovascular system antigen expression level in cardiovascular system tissues or bodily fluids from an individual not having the cardiovascular system disorder. Thus, the invention provides a diagnostic method useful during diagnosis of a cardiovascular system disorder, which involves measuring the expression level of the gene encoding the cardiovascular system associated polypeptide in cardiovascular system tissue or other cells or body fluid from an individual and comparing the measured gene expression level with a standard cardiovascular system antigens gene expression level, whereby an increase or decrease in the gene expression level(s) compared to the standard is indicative of an cardiovascular system disorder.

[00387] In specific embodiments, the invention provides a diagnostic method useful during diagnosis of a disorder of a normal or diseased tissue/cell source corresponding to column 7 of Table 1A, which involves measuring the expression level of the coding sequence of a polynucleotide sequence associated with this tissue/cell source as disclosed in Table 1A in the tissue/cell source or other cells or body fluid from an individual and comparing the expression level of the coding sequence with a standard expression level of the coding sequence of a polynucleotide sequence, whereby an increase or decrease in the gene expression level(s) compared to the standard is indicative of a disorder of a normal or diseased tissue/cell source corresponding to column 7 of Table 1A.

[00388] In particular, it is believed that certain tissues in mammals with cancer of cells or tissue of the cardiovascular system express significantly enhanced or reduced levels of normal or altered cardiovascular system antigen expression and mRNA

encoding the cardiovascular system associated polypeptide when compared to a corresponding "standard" level. Further, it is believed that enhanced or depressed levels of the cardiovascular system associated polypeptide can be detected in certain body fluids (e.g., sera, plasma, urine, and spinal fluid) or cells or tissue from mammals with such a cancer when compared to sera from mammals of the same species not having the cancer.

[00389] For example, as disclosed herein, cardiovascular system associated polypeptides of the invention are expressed in the cardiovascular system. Accordingly, polynucleotides of the invention (e.g., polynucleotide sequences complementary to all or a portion of a cardiovascular system antigen mRNA nucleotide sequence of SEQ ID NO:X, nucleotide sequence encoding SEQ ID NO:Y, nucleotide sequence encoding a polypeptide encoded by SEQ ID NO:X and/or a nucleotide sequence delineated by columns 8 and 9 of Table 2) and antibodies (and antibody fragments) directed against the polypeptides of the invention may be used to quantitate or qualitate concentrations of cells of the cardiovascular system expressing cardiovascular system antigens, preferably on their cell surfaces. These polynucleotides and antibodies additionally have diagnostic applications in detecting abnormalities in the level of cardiovascular system antigens gene expression, or abnormalities in the structure and/or temporal, tissue, cellular, or subcellular location of cardiovascular system antigens. These diagnostic assays may be performed *in vivo* or *in vitro*, such as, for example, on blood samples, biopsy tissue or autopsy tissue. In specific embodiments, polynucleotides and antibodies of the invention are used to quantitate or qualitate tissues/cells corresponding to the library source disclosed in column 7 of Table 1A expressing the corresponding cardiovascular system sequence disclosed in the same row of Table 1A, preferably on their cell surface.

[00390] Thus, the invention provides a diagnostic method useful during diagnosis of a cardiovascular system disorder, including cancers, which involves measuring the expression level of the gene encoding the cardiovascular system antigen polypeptide in cardiovascular system tissue or other cells or body fluid from an individual and comparing the measured gene expression level with a standard cardiovascular system antigen gene expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a cardiovascular system



disorder. In specific embodiments, polynucleotides and antibodies of the invention are used to quantitate or qualitate tissues/cells corresponding to the library source disclosed in column 7 of Table 1A expressing the corresponding cardiovascular systemcardiovascular system sequence disclosed in the same row of Table 1A, preferably on their cell surface.

[00391] Where a diagnosis of a disorder in the cardiovascular system, including diagnosis of a tumor, has already been made according to conventional methods, the present invention is useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed cardiovascular system antigen gene expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

[00392] By "assaying the expression level of the gene encoding the cardiovascular system associated polypeptide" is intended qualitatively or quantitatively measuring or estimating the level of the cardiovascular system antigen polypeptide or the level of the mRNA encoding the cardiovascular system antigen polypeptide in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the cardiovascular system associated polypeptide level or mRNA level in a second biological sample). Preferably, the cardiovascular system antigen polypeptide expression level or mRNA level in the first biological sample is measured or estimated and compared to a standard cardiovascular system antigen polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder of the cardiovascular system. As will be appreciated in the art, once a standard cardiovascular system antigen polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

[00393] By "biological sample" is intended any biological sample obtained from an individual, cell line, tissue culture, or other source containing cardiovascular system antigen polypeptides (including portions thereof) or mRNA. As indicated, biological samples include body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) which contain cells expressing cardiovascular system antigen polypeptides, cardiovascular system tissue, and other tissue sources found to express the full

length or fragments thereof of a cardiovascular system antigen. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

[00394] Total cellular RNA can be isolated from a biological sample using any suitable technique such as the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987). Levels of mRNA encoding the cardiovascular system antigen polypeptides are then assayed using any appropriate method. These include Northern blot analysis, S1 nuclease mapping, the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR), and reverse transcription in combination with the ligase chain reaction (RT-LCR).

[00395] The present invention also relates to diagnostic assays such as quantitative and diagnostic assays for detecting levels of cardiovascular system antigen polypeptides, in a biological sample (e.g., cells and tissues), including determination of normal and abnormal levels of polypeptides. Thus, for instance, a diagnostic assay in accordance with the invention for detecting over-expression of cardiovascular system antigens compared to normal control tissue samples may be used to detect the presence of tumors. Assay techniques that can be used to determine levels of a polypeptide, such as a cardiovascular system antigen polypeptide of the present invention in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays. Assaying cardiovascular system antigen polypeptide levels in a biological sample can occur using any art-known method.

[00396] Assaying cardiovascular system antigen polypeptide levels in a biological sample can occur using antibody-based techniques. For example, cardiovascular system antigen polypeptide expression in tissues can be studied with classical immunohistological methods (Jalkanen et al., *J. Cell. Biol.* 101:976-985 (1985); Jalkanen, M., et al., *J. Cell. Biol.* 105:3087-3096 (1987)). Other antibody-based methods useful for detecting cardiovascular system antigen polypeptide gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase, and

radioisotopes, such as iodine ( $^{125}\text{I}$ ,  $^{121}\text{I}$ ), carbon ( $^{14}\text{C}$ ), sulfur ( $^{35}\text{S}$ ), tritium ( $^3\text{H}$ ), indium ( $^{112}\text{In}$ ), and technetium ( $^{99\text{m}}\text{Tc}$ ), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00397] The tissue or cell type to be analyzed will generally include those, which are known, or suspected, to express the cardiovascular system antigen gene (such as, for example, cells of the cardiovascular system or cancer of cardiovascular system tissues). The protein isolation methods employed herein may, for example, be such as those described in Harlow and Lane (Harlow, E. and Lane, D., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York), which is incorporated herein by reference in its entirety. The isolated cells can be derived from cell culture or from a patient. The analysis of cells taken from culture may be a necessary step in the assessment of cells that could be used as part of a cell-based gene therapy technique or, alternatively, to test the effect of compounds on the expression of the cardiovascular system antigen gene.

[00398] For example, antibodies, or fragments of antibodies, such as those described herein, may be used to quantitatively or qualitatively detect the presence of cardiovascular system antigen gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or fluorimetric detection.

[00399] In a preferred embodiment, antibodies, or fragments of antibodies directed to any one or all of the predicted epitope domains of the cardiovascular system antigen polypeptides (Shown in Table 1A, column 6) may be used to quantitatively or qualitatively detect the presence of cardiovascular system antigen gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or fluorimetric detection.

[00400] In an additional preferred embodiment, antibodies, or fragments of antibodies directed to a conformational epitope of a cardiovascular system antigen may be used to quantitatively or qualitatively detect the presence of cardiovascular system antigen gene products or conserved variants or peptide fragments thereof. This can be accomplished, for example, by immunofluorescence techniques employing a fluorescently labeled antibody coupled with light microscopic, flow cytometric, or

fluorimetric detection.

[00401] The antibodies (or fragments thereof), and/or cardiovascular system antigen polypeptides of the present invention may, additionally, be employed histologically, as in immunofluorescence, immunoelectron microscopy or non-immunological assays, for in situ detection of cardiovascular system antigen gene products or conserved variants or peptide fragments thereof. In situ detection may be accomplished by removing a histological specimen from a patient, and applying thereto a labeled antibody or cardiovascular system antigen polypeptide of the present invention. The antibody (or fragment thereof) or cardiovascular system antigen polypeptide is preferably applied by overlaying the labeled antibody (or fragment) onto a biological sample. Through the use of such a procedure, it is possible to determine not only the presence of the cardiovascular system antigen gene product, or conserved variants or peptide fragments, or cardiovascular system antigen polypeptide binding, but also its distribution in the examined tissue. Using the present invention, those of ordinary skill will readily perceive that any of a wide variety of histological methods (such as staining procedures) can be modified in order to achieve such in situ detection.

[00402] Immunoassays and non-immunoassays for cardiovascular system antigen gene products or conserved variants or peptide fragments thereof will typically comprise incubating a sample, such as a biological fluid, a tissue extract, freshly harvested cells, or lysates of cells which have been incubated in cell culture, in the presence of a detectably labeled antibody capable of binding cardiovascular system antigen gene products or conserved variants or peptide fragments thereof, and detecting the bound antibody by any of a number of techniques well-known in the art.

[00403] The biological sample may be brought in contact with and immobilized onto a solid phase support or carrier such as nitrocellulose, or other solid support which is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled anti-cardiovascular system antigen antibody or detectable cardiovascular system antigen polypeptide. The solid phase support may then be washed with the buffer a second time to remove unbound antibody or polypeptide. Optionally the antibody is subsequently labeled. The amount of bound label on solid support may then be detected by conventional means.

10041504 030706

[00404] By "solid phase support or carrier" is intended any support capable of binding an antigen or an antibody. Well-known supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, gabbros, and magnetite. The nature of the carrier can be either soluble to some extent or insoluble for the purposes of the present invention. The support material may have virtually any possible structural configuration so long as the coupled molecule is capable of binding to an antigen or antibody. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads. Those skilled in the art will know many other suitable carriers for binding antibody or antigen, or will be able to ascertain the same by use of routine experimentation.

[00405] The binding activity of a given lot of anti-cardiovascular system antigen antibody or cardiovascular system antigen polypeptide may be determined according to well-known methods. Those skilled in the art will be able to determine operative and optimal assay conditions for each determination by employing routine experimentation.

[00406] In addition to assaying cardiovascular system antigen polypeptide levels or polynucleotide levels in a biological sample obtained from an individual, cardiovascular system antigen polypeptide or polynucleotide can also be detected *in vivo* by imaging. For example, in one embodiment of the invention, cardiovascular system antigen polypeptide and/or anti-cardiovascular system antigen antibodies are used to image cardiovascular system diseased cells, such as neoplasms. In another embodiment, cardiovascular system antigen polynucleotides of the invention (e.g., polynucleotides complementary to all or a portion of cardiovascular system antigen mRNA) and/or anti-cardiovascular system antigen antibodies (e.g., antibodies directed to any one or a combination of the epitopes of cardiovascular system antigens, antibodies directed to a conformational epitope of cardiovascular system antigens, antibodies directed to the full length polypeptide expressed on the cell surface of a mammalian cell) are used to image diseased or neoplastic cells of the cardiovascular system.

[00407] Antibody labels or markers for *in vivo* imaging of cardiovascular system antigen

polypeptides include those detectable by X-radiography, NMR, MRI, CAT-scans or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma. Where *in vivo* imaging is used to detect enhanced levels of cardiovascular system antigen polypeptides for diagnosis in humans, it may be preferable to use human antibodies or "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using techniques described herein or otherwise known in the art. For example methods for producing chimeric antibodies are known in the art. See, for review, Morrison, *Science* 229:1202 (1985); Oi et al., *BioTechniques* 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., *Nature* 312:643 (1984); Neuberger et al., *Nature* 314:268 (1985).

[00408] Additionally, any cardiovascular system antigen polypeptides whose presence can be detected, can be administered. For example, cardiovascular system antigen polypeptides labeled with a radio-opaque or other appropriate compound can be administered and visualized *in vivo*, as discussed, above for labeled antibodies. Further such cardiovascular system antigen polypeptides can be utilized for *in vitro* diagnostic procedures.

[00409] A cardiovascular system antigen polypeptide-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example,  $^{131}\text{I}$ ,  $^{112}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ ), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for a cardiovascular system disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of  $^{99\text{m}}\text{Tc}$ . The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain cardiovascular system antigen protein. *In vivo* tumor imaging is described in S.W. Burchiel et al.,

"Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

[00410] With respect to antibodies, one of the ways in which the anti-cardiovascular system antigen antibody can be detectably labeled is by linking the same to an enzyme and using the linked product in an enzyme immunoassay (EIA) (Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)", 1978, Diagnostic Horizons 2:1-7, Microbiological Associates Quarterly Publication, Walkersville, MD); Voller et al., *J. Clin. Pathol.* 31:507-520 (1978); Butler, J.E., *Meth. Enzymol.* 73:482-523 (1981); Maggio, E. (ed.), 1980, Enzyme Immunoassay, CRC Press, Boca Raton, FL.; Ishikawa, E. et al., (eds.), 1981, Enzyme Immunoassay, Kigaku Shoin, Tokyo). The enzyme, which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety which can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Enzymes which can be used to detectably label the antibody include, but are not limited to, malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, alpha-glycerophosphate dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. Additionally, the detection can be accomplished by colorimetric methods, which employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

[00411] Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect cardiovascular system antigens through the use of a radioimmunoassay (RIA) (see, for example, Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986, which is incorporated by reference herein). The radioactive isotope can be detected by means including, but not limited to, a gamma counter, a scintillation counter, or autoradiography.

[00412] It is also possible to label the antibody with a fluorescent compound. When the

fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, ophthaldehyde and fluorescamine.

[00413] The antibody can also be detectably labeled using fluorescence emitting metals such as  $^{152}\text{Eu}$ , or others of the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

[00414] The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of particularly useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

[00415] Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in, which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

#### **Methods for Detecting Cardiovascular System Disease, Including Cancer**

[00416] In general, a cardiovascular system disease or cancer may be detected in a patient based on the presence of one or more cardiovascular system antigen proteins of the invention and/or polynucleotides encoding such proteins in a biological sample (for example, blood, sera, urine, and/or tumor biopsies) obtained from the patient. In other words, such proteins and/or polynucleotides may be used as markers to indicate the presence or absence of a cardiovascular system disease or disorder, including cancer. Cancers that may be diagnosed, and/or prognosed using the compositions of the invention include but are not limited to, cancer of cardiovascular system tissues. In addition, such proteins and/or polynucleotides may be useful for the detection of other diseases and cancers, including cancers of tissues/cells corresponding to the library source disclosed in column 7 of Table 1A



expressing the corresponding cardiovascular system sequence disclosed in the same row of Table 1A. The binding agents provided herein generally permit detection of the level of antigen that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding cardiovascular system antigen polypeptides, which is also indicative of the presence or absence of a cardiovascular system disease or disorder, including cancer. In general, cardiovascular system antigen polypeptides should be present at a level that is at least three fold higher in diseased tissue than in normal tissue.

[00417] There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. See, e.g., Harlow and Lane, *supra*. In general, the presence or absence of a cardiovascular system disease in a patient may be determined by (a) contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

[00418] In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the cardiovascular system antigen polypeptide of the invention from the remainder of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include cardiovascular system antigen polypeptides and portions thereof, or antibodies, to which the binding agent binds, as described above.

[00419] The solid support may be any material known to those of skill in the art to which cardiovascular system antigen polypeptides of the invention may be attached. For

example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane. Alternatively, the support may be a bead or disc, such as glass fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for the suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 ug, and preferably about 100 ng to about 1 ug, is sufficient to immobilize an adequate amount of binding agent.

[00420] Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (see, e.g., Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

#### **Gene Therapy Methods**

[00421] Also encompassed by the present invention are gene therapy methods for treating or preventing disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of a cardiovascular system antigen

of the present invention. This method requires a polynucleotide which codes for a polypeptide of the present invention operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

[00422] Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the present invention ex vivo, with the engineered cells then being provided to a patient to be treated with the polypeptide of the present invention. Such methods are well-known in the art. For example, see Belldégrun, A., et al., J. Natl. Cancer Inst. 85: 207-216 (1993); Ferrantini, M. et al., Cancer Research 53: 1107-1112 (1993); Ferrantini, M. et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura, H., et al., Cancer Research 50: 5102-5106 (1990); Santodonato, L., et al., Human Gene Therapy 7:1-10 (1996); Santodonato, L., et al., Gene Therapy 4:1246-1255 (1997); and Zhang, J.-F. et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells, which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

[00423] As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

[00424] In one embodiment, the polynucleotide of the present invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotide of the present invention can also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in

U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

[00425] The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors will be readily apparent to the skilled artisan.

[00426] Any strong promoter known to those skilled in the art can be used for driving the expression of the polynucleotide sequence. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoAI promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotide of the present invention.

[00427] Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

[00428] The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid

of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

[00429] For the naked nucleic acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

[00430] The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

[00431] The naked polynucleotides are delivered by any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

[00432] The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of delivery are known in the art.

[00433] In certain embodiments, the polynucleotide constructs are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight

charge complex can be formed between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference); mRNA (Malone et al., Proc. Natl. Acad. Sci. USA (1989) 86:6077-6081, which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem. (1990) 265:10189-10192, which is herein incorporated by reference), in functional form.

[00434] Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y., (see, also, Felgner et al., Proc. Natl Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

[00435] Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication No. WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., P. Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

[00436] Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

[00437] For example, commercially dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with

or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

[00438] The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology* (1983), 101:512-527, which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include  $\text{Ca}^{2+}$ -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta* (1975) 394:483; Wilson et al., *Cell* 17:77 (1979); ether injection (Deamer, D. and Bangham, A., *Biochim. Biophys. Acta* 443:629 (1976); Ostro et al., *Biochem. Biophys. Res. Commun.* 76:836 (1977); Fraley et al., *Proc. Natl. Acad. Sci. USA* 76:3348 (1979)); detergent dialysis (Enoch, H. and Strittmatter, P., *Proc. Natl. Acad. Sci. USA* 76:145 (1979)); and reverse-phase evaporation (REV) (Fraley et al., *J. Biol. Chem.* 255:10431 (1980); Szoka et al.,

Proc. Natl. Acad. Sci. USA 75:145 (1978); Schaefer-Ridder et al., Science 215:166 (1982)), which are herein incorporated by reference.

[00439] Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

[00440] U.S. Patent No. 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and International Publication No. WO 94/9469 provide methods for delivering DNA-cationic lipid complexes to mammals.

[00441] In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA, which comprises a sequence encoding a polypeptide of the present invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

[00442] The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE, RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO<sub>4</sub> precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host.

[00443] The producer cell line generates infectious retroviral vector particles, which include polynucleotide encoding a polypeptide of the present invention. Such



retroviral vector particles then may be employed; to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express a polypeptide of the present invention.

[00444] In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotide contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses a polypeptide of the present invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome, thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz, et al., Am. Rev. Respir. Dis.109:233-238 (1974)). Finally, adenovirus mediated gene transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld et al., Science 252:431-434 (1991); Rosenfeld et al., Cell 68:143-155 (1991)). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green et al., Proc. Natl. Acad. Sci. USA 76:6606 (1979)).

[00445] Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel. 3:499-503 (1993); Rosenfeld et al., Cell 68:143-155 (1992); Engelhardt et al., Human Genet. Ther. 4:759-769 (1993); Yang et al., Nature Genet. 7:362-369 (1994); Wilson et al., Nature 365:691-692 (1993); and U.S. Patent No. 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express Ela and Elb, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

[00446] Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses

may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

[00447] In certain other embodiments, the cells are engineered, *ex vivo* or *in vivo*, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, N., Curr. Topics in Microbiol. Immunol. 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

[00448] For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct is inserted into the AAV vector using standard cloning methods, such as those found in Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles, which contain the polynucleotide construct. These viral particles are then used to transduce eukaryotic cells, either *ex vivo* or *in vivo*. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express a polypeptide of the invention.

[00449] Another method of gene therapy involves operably associating heterologous control regions and endogenous cardiovascular system antigen polynucleotide sequences (e.g., encoding a cardiovascular system antigen polypeptide of the present invention) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), which are herein incorporated by reference. This

method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

[00450] Polynucleotide constructs are made, using standard techniques known in the art, which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination.

[00451] The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

[00452] The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

[00453] The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous sequence is placed under the control of the promoter. The promoter then drives the expression of the endogenous sequence.

[00454] The polynucleotide encoding a polypeptide of the present invention may contain a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed towards or at the 5' end of the coding region. The signal sequence may be

homologous or heterologous to the cardiovascular system antigen polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

[00455] Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gelfoam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppositorial solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers (Kaneda et al., Science 243:375 (1989)).

[00456] A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

[00457] Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

[00458] Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site. In specific embodiments, suitable delivery vehicles for use with systemic administration comprise liposomes comprising polypeptides of the invention for targeting the vehicle to a particular site.

10091504-030702

[00459] Preferred methods of systemic administration, include intravenous injection, aerosol, oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA 189:11277-11281, 1992, which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

[00460] Determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the attending physician or veterinarian.

[00461] Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly preferred.

#### **Biological Activities**

[00462] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, can be used in assays to test for one or more biological activities. If these polynucleotides or polypeptides, or agonists or antagonists of the present invention, do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides, and agonists or antagonists could be used to treat, prevent diagnose and/or prognose the associated disease.

[00463] The cardiovascular system antigen polynucleotides and polypeptides of the invention are predicted to have predominant expression in cardiovascular system tissues.

[00464] Thus, the cardiovascular system antigens of the invention may be useful as therapeutic molecules. Each would be useful for diagnosis, detection, treatment and/or prevention of diseases or disorders of the cardiovascular system, including but not limited to cardiovascular abnormalities (e.g., congenital heart defects, cerebral arteriovenous malformations, and septal defects), heart disease (e.g., heart failure, cardiomyopathy, pericarditis, and endocarditis), arrhythmias, heart valve disease (e.g., stenosis, regurgitation, and prolapse), vascular diseases (e.g., arteriosclerosis, coronary artery disease, angina, varicose veins, hypertension, and shock), electrolyte imbalance disorders (e.g., hypo- and hypernatremia, and hypo- and hyperkalemia), and/or as described under "Cardiovascular Disorders" below.

[00465] In a preferred embodiment, polynucleotides of the invention (e.g., a nucleic acid sequence of SEQ ID NO:X or the complement thereof; or the cDNA sequence contained in Clone ID NO:Z, or fragments or variants thereof) and/or polypeptides of the invention (e.g., an amino acid sequence contained in SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X, or the complement thereof, an amino acid sequence encoded by the cDNA sequence contained in Clone ID NO:Z and fragments or variants thereof as described herein) are useful for the diagnosis, detection, treatment, and/or prevention of diseases or disorders of the tissues/cells corresponding to the library source disclosed in column 7 of Table 1A expressing the corresponding cardiovascular system sequence disclosed in the same row of Table 1A.

[00466] Particularly, the cardiovascular system antigens may be a useful therapeutic for cancer of cardiovascular system tissues. Treatment, diagnosis, detection, and/or prevention of cardiovascular system disorders could be carried out using a cardiovascular system antigen or soluble form of a cardiovascular system antigen, a cardiovascular system antigen ligand, gene therapy, or ex vivo applications. Moreover, inhibitors of a cardiovascular system antigen, either blocking antibodies or mutant forms, could modulate the expression of the cardiovascular system antigen. These inhibitors may be useful to treat, diagnose, detect, and/or prevent diseases associated with the misregulation of a cardiovascular system antigen.

10091504-030703

[00467] In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells (e.g., normal or diseased cardiovascular system cells) by administering polypeptides of the invention (e.g., cardiovascular system antigen polypeptides or anti-cardiovascular system antigen antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell (e.g., an aberrant cardiovascular system cell or cardiovascular system cancer cell). In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

[00468] In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of aberrant cardiovascular system cells, including, but not limited to, cardiovascular system tumor cells) by administering polypeptides of the invention (e.g., cardiovascular system antigen polypeptides or fragments thereof, or anti-cardiovascular system antigen antibodies) in association with toxins or cytotoxic prodrugs.

[00469] By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, cytotoxins (cytotoxic agents), or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. "Toxin" also includes a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example,  $^{213}\text{Bi}$ , or other radioisotopes such as, for example,  $^{103}\text{Pd}$ ,  $^{133}\text{Xe}$ ,  $^{131}\text{I}$ ,  $^{68}\text{Ge}$ ,  $^{57}\text{Co}$ ,  $^{65}\text{Zn}$ ,  $^{85}\text{Sr}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ ,  $^{90}\text{Y}$ ,  $^{153}\text{Sm}$ ,  $^{153}\text{Gd}$ ,  $^{169}\text{Yb}$ ,  $^{51}\text{Cr}$ ,  $^{54}\text{Mn}$ ,  $^{75}\text{Se}$ ,  $^{113}\text{Sn}$ ,  $^{90}\text{Yttrium}$ ,  $^{117}\text{Tin}$ ,  $^{186}\text{Rhenium}$ ,  $^{166}\text{Holmium}$ , and  $^{188}\text{Rhenium}$ ; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

[00470] Techniques known in the art may be applied to label antibodies of the invention.

Such techniques include, but are not limited to, the use of bifunctional conjugating agents (see e.g., U.S. Patent Nos. 5,756,065; 5,714,631; 5,696,239; 5,652,361; 5,505,931; 5,489,425; 5,435,990; 5,428,139; 5,342,604; 5,274,119; 4,994,560; and 5,808,003; the contents of each of which are hereby incorporated by reference in its entirety). A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis- dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[00471] By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

[00472] It will be appreciated that conditions caused by a decrease in the standard or normal level of a cardiovascular system antigen activity in an individual, particularly disorders of the cardiovascular system, can be treated by administration of a cardiovascular system antigen polypeptide (e.g., such as, for example, the complete cardiovascular system antigen polypeptide, the soluble form of the extracellular domain of a cardiovascular system antigen polypeptide, or cells expressing the complete protein) or agonist. Thus, the invention also provides a



method of treatment of an individual in need of an increased level of cardiovascular system antigen activity comprising administering to such an individual a pharmaceutical composition comprising an amount of an isolated cardiovascular system antigen polypeptide of the invention, or agonist thereof (e.g., an agonistic anti-cardiovascular system antigen antibody), effective to increase the cardiovascular system antigen activity level in such an individual.

[00473] It will also be appreciated that conditions caused by a increase in the standard or normal level of cardiovascular system antigen activity in an individual, particularly disorders of the cardiovascular system, can be treated by administration of cardiovascular system antigen polypeptides (e.g., such as, for example, the complete cardiovascular system antigen polypeptide, the soluble form of the extracellular domain of a cardiovascular system antigen polypeptide, or cells expressing the complete protein) or antagonist (e.g., an antagonistic cardiovascular system antigen antibody). Thus, the invention also provides a method of treatment of an individual in need of an decreased level of cardiovascular system antigen activity comprising administering to such an individual a pharmaceutical composition comprising an amount of an isolated cardiovascular system antigen polypeptide of the invention, or antagonist thereof (e.g., an antagonistic anti-cardiovascular system antigen antibody), effective to decrease the cardiovascular system antigen activity level in such an individual.

[00474] In certain embodiments, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose and/or prognose diseases and/or disorders associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1A, column 7 (Tissue Distribution Library Code).

[00475] More generally, polynucleotides, translation products and antibodies corresponding to this gene may be useful for the diagnosis, prognosis, prevention, and/or treatment of diseases and/or disorders associated with the following systems.

#### **Cardiovascular Disorders**

[00476] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose cardiovascular

disorders, including, but not limited to, peripheral artery disease, such as limb ischemia.

[00477] Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects, pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, tricuspid atresia, persistent truncus arteriosus, total anomalous pulmonary venous connection, hypoplastic left heart syndrome, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, atrioventricular canal defect, trilogly of Fallot, ventricular heart septal defects.

[00478] Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis (including bacterial), heart aneurysm, cardiac arrest, sudden cardiac death, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, diastolic dysfunction, enlarged heart, heart block, J-curve phenomenon, rheumatic heart disease, Marfan syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

[00479] Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic

junctional tachycardia, sinoatrial nodal reentry tachycardia, sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

[00480] Heart valve disease include aortic valve insufficiency, aortic valve stenosis, heart murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, tricuspid valve stenosis, and bicuspid aortic valve.

[00481] Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis, restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, Barth syndrome, myocardial reperfusion injury, and myocarditis.

[00482] Myocardial ischemias include coronary disease, such as angina pectoris, Prinzmetal's angina, unstable angina, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

[00483] Cardiovascular diseases also include vascular diseases such as aneurysms, angiodysplasia, angiomas, bacillary angiomas, Hippel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension (shock), ischemia, peripheral vascular diseases, phlebitis, superficial phlebitis, pulmonary veno-occlusive disease, chronic obstructive pulmonary disease, Buerger's disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, ataxia telangiectasia, hereditary hemorrhagic telangiectasia, deep vein thrombosis, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

[00484] Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

- [00485] Arterial occlusive diseases include arteriosclerosis, arteriolosclerosis, atherosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.
- [00486] Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar insufficiency.
- [00487] Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms. Thrombosis include coronary thrombosis, hepatic vein thrombosis, deep vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.
- [00488] Ischemia includes cerebral ischemia, ischemic colitis, silent ischemia, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.
- [00489] Cardiovascular diseases can also occur due to electrolyte imbalances that include, but are not limited to hyponatremia, hypernatremia, hypokalemia, hyperkalemia, hypocalcemia, hypercalcemia, hypophosphatemia, and hyperphosphatemia. Neoplasm and/or cancers of the cardiovascular system include, but are not limited to, myxomas, fibromas, and rhabdomyomas.
- [00490] Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical

applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

### **Blood-Related Disorders**

[00491] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate hemostatic (the stopping of bleeding) or thrombolytic (clot dissolving) activity. For example, by increasing hemostatic or thrombolytic activity, polynucleotides or polypeptides, and/or agonists or antagonists of the present invention could be used to treat or prevent blood coagulation diseases, disorders, and/or conditions (e.g., afibrinogenemia, factor deficiencies, hemophilia), blood platelet diseases, disorders, and/or conditions (e.g., thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment or prevention of heart attacks (infarction), strokes, or scarring.

[00492] In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to prevent, diagnose, prognose, and/or treat thrombosis, arterial thrombosis, venous thrombosis, thromboembolism, pulmonary embolism, atherosclerosis, myocardial infarction, transient ischemic attack, unstable angina. In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used for the prevention of occlusion of saphenous grafts, for reducing the risk of periprocedural thrombosis as might accompany angioplasty procedures, for reducing the risk of stroke in patients with atrial fibrillation including nonrheumatic atrial fibrillation, for reducing the risk of embolism associated with mechanical heart valves and or mitral valves disease. Other uses for the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention, include, but are not limited to, the prevention of occlusions in extracorporeal devices (e.g., intravascular canulas, vascular access shunts in

hemodialysis patients, hemodialysis machines, and cardiopulmonary bypass machines).

[00493] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to prevent, diagnose, prognose, and/or treat diseases and disorders of the blood and/or blood forming organs associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1A, column 7 (Tissue Distribution Library Code).

[00494] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate hematopoietic activity (the formation of blood cells). For example, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to increase the quantity of all or subsets of blood cells, such as, for example, erythrocytes, lymphocytes (B or T cells), myeloid cells (e.g., basophils, eosinophils, neutrophils, mast cells, macrophages) and platelets. The ability to decrease the quantity of blood cells or subsets of blood cells may be useful in the prevention, detection, diagnosis and/or treatment of anemias and leukopenias described below. Alternatively, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to decrease the quantity of all or subsets of blood cells, such as, for example, erythrocytes, lymphocytes (B or T cells), myeloid cells (e.g., basophils, eosinophils, neutrophils, mast cells, macrophages) and platelets.. The ability to decrease the quantity of blood cells or subsets of blood cells may be useful in the prevention, detection, diagnosis and/or treatment of leukocytoses, such as, for example eosinophilia.

[00495] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to prevent, treat, or diagnose blood dyscrasia.

[00496] Anemias are conditions in which the number of red blood cells or amount of hemoglobin (the protein that carries oxygen) in them is below normal. Anemia may be caused by excessive bleeding, decreased red blood cell production, or increased red blood cell destruction (hemolysis). The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias. Anemias that may be treated prevented or diagnosed by the polynucleotides, polypeptides, antibodies, and/or

agonists or antagonists of the present invention include iron deficiency anemia, hypochromic anemia, microcytic anemia, chlorosis, hereditary sideroblastic anemia, idiopathic acquired sideroblastic anemia, red cell aplasia, megaloblastic anemia (e.g., pernicious anemia, (vitamin B12 deficiency) and folic acid deficiency anemia), aplastic anemia, hemolytic anemias (e.g., autoimmune hemolytic anemia, microangiopathic hemolytic anemia, and paroxysmal nocturnal hemoglobinuria). The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias associated with diseases including but not limited to, anemias associated with systemic lupus erythematosus, cancers, lymphomas, chronic renal disease, and enlarged spleens. The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias arising from drug treatments such as anemias associated with methyldopa, dapsone, and/or sulfadruugs. Additionally, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing anemias associated with abnormal red blood cell architecture including but not limited to, hereditary spherocytosis, hereditary elliptocytosis, glucose-6-phosphate dehydrogenase deficiency, and sickle cell anemia.

[00497] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing hemoglobin abnormalities, (e.g., those associated with sickle cell anemia, hemoglobin C disease, hemoglobin S-C disease, and hemoglobin E disease). Additionally, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating thalassemias, including, but not limited to major and minor forms of alpha-thalassemia and beta-thalassemia.

[00498] In another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating bleeding disorders including, but not limited to, thrombocytopenia (e.g., idiopathic thrombocytopenic purpura, and thrombotic thrombocytopenic purpura), Von Willebrand's disease, hereditary platelet disorders (e.g., storage pool disease such as Chediak-Higashi and Hermansky-Pudlak

syndromes, thromboxane A2 dysfunction, thromboasthenia, and Bernard-Soulier syndrome), hemolytic-uremic syndrome, hemophelias such as hemophilia A or Factor VII deficiency and Christmas disease or Factor IX deficiency, Hereditary Hemorrhagic Telangiectasia, also known as Rendu-Osler-Weber syndrome, allergic purpura (Henoch Schonlein purpura) and disseminated intravascular coagulation.

[00499] The effect of the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention on the clotting time of blood may be monitored using any of the clotting tests known in the art including, but not limited to, whole blood partial thromboplastin time (PTT), the activated partial thromboplastin time (aPTT), the activated clotting time (ACT), the recalcified activated clotting time, or the Lee-White Clotting time.

[00500] Several diseases and a variety of drugs can cause platelet dysfunction. Thus, in a specific embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating acquired platelet dysfunction such as platelet dysfunction accompanying kidney failure, leukemia, multiple myeloma, cirrhosis of the liver, and systemic lupus erythematosus as well as platelet dysfunction associated with drug treatments, including treatment with aspirin, ticlopidine, nonsteroidal anti-inflammatory drugs (used for arthritis, pain, and sprains), and penicillin in high doses.

[00501] In another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating diseases and disorders characterized by or associated with increased or decreased numbers of white blood cells. Leukopenia occurs when the number of white blood cells decreases below normal. Leukopenias include, but are not limited to, neutropenia and lymphocytopenia. An increase in the number of white blood cells compared to normal is known as leukocytosis. The body generates increased numbers of white blood cells during infection. Thus, leukocytosis may simply be a normal physiological parameter that reflects infection. Alternatively, leukocytosis may be an indicator of injury or other disease such as cancer. Leokocytoses, include but are not limited to, eosinophilia, and accumulations of macrophages. In specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may



be useful in diagnosing, prognosing, preventing, and/or treating leukopenia. In other specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating leukocytosis

[00502] Leukopenia may be a generalized decreased in all types of white blood cells, or may be a specific depletion of particular types of white blood cells. Thus, in specific embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating decreases in neutrophil numbers, known as neutropenia. Neutropenias that may be diagnosed, prognosed, prevented, and/or treated by the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, infantile genetic agranulocytosis, familial neutropenia, cyclic neutropenia, neutropenias resulting from or associated with dietary deficiencies (e.g., vitamin B 12 deficiency or folic acid deficiency), neutropenias resulting from or associated with drug treatments (e.g., antibiotic regimens such as penicillin treatment, sulfonamide treatment, anticoagulant treatment, anticonvulsant drugs, anti-thyroid drugs, and cancer chemotherapy), and neutropenias resulting from increased neutrophil destruction that may occur in association with some bacterial or viral infections, allergic disorders, autoimmune diseases, conditions in which an individual has an enlarged spleen (e.g., Felty syndrome, malaria and sarcoidosis), and some drug treatment regimens.

[00503] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating lymphocytopenias (decreased numbers of B and/or T lymphocytes), including, but not limited lymphocytopenias resulting from or associated with stress, drug treatments (e.g., drug treatment with corticosteroids, cancer chemotherapies, and/or radiation therapies), AIDS infection and/or other diseases such as, for example, cancer, rheumatoid arthritis, systemic lupus erythematosus, chronic infections, some viral infections and/or hereditary disorders (e.g., DiGeorge syndrome, Wiskott-Aldrich Syndrome, severe combined immunodeficiency, ataxia telangiectasia).

[00504] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or

treating diseases and disorders associated with macrophage numbers and/or macrophage function including, but not limited to, Gaucher's disease, Niemann-Pick disease, Letterer-Siwe disease and Hand-Schuller-Christian disease.

[00505] In another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating diseases and disorders associated with eosinophil numbers and/or eosinophil function including, but not limited to, idiopathic hypereosinophilic syndrome, eosinophilia-myalgia syndrome, and Hand-Schuller-Christian disease.

[00506] In yet another embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating leukemias and lymphomas including, but not limited to, acute lymphocytic (lymphoblastic) leukemia (ALL), acute myeloid (myelocytic, myelogenous, myeloblastic, or myelomonocytic) leukemia, chronic lymphocytic leukemia (e.g., B cell leukemias, T cell leukemias, Sezary syndrome, and Hairy cell leukemia), chronic myelocytic (myeloid, myelogenous, or granulocytic) leukemia, Hodgkin's lymphoma, non-hodgkin's lymphoma, Burkitt's lymphoma, and mycosis fungoides.

[00507] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in diagnosing, prognosing, preventing, and/or treating diseases and disorders of plasma cells including, but not limited to, plasma cell dyscrasias, monoclonal gammaopathies, monoclonal gammopathies of undetermined significance, multiple myeloma, macroglobulinemia, Waldenstrom's macroglobulinemia, cryoglobulinemia, and Raynaud's phenomenon.

[00508] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, and/or diagnosing myeloproliferative disorders, including but not limited to, polycythemia vera, relative polycythemia, secondary polycythemia, myelofibrosis, acute myelofibrosis, agnogenic myeloid metaplasia, thrombocythemia, (including both primary and secondary thrombocythemia) and chronic myelocytic leukemia.

[00509] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as a treatment prior to surgery, to increase blood cell production.

[00510] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to enhance the migration, phagocytosis, superoxide production, antibody dependent cellular cytotoxicity of neutrophils, eosinophils and macrophages.

[00511] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase the number of stem cells in circulation prior to stem cells pheresis. In another specific embodiment, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase the number of stem cells in circulation prior to platelet pheresis.

[00512] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful as an agent to increase cytokine production.

[00513] In other embodiments, the polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in preventing, diagnosing, and/or treating primary hematopoietic disorders.

#### Immune Activity

[00514] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, diagnosing and/or prognosing diseases, disorders, and/or conditions of the immune system, by, for example, activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells. Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune diseases, disorders, and/or conditions may be genetic, somatic, such as cancer and some autoimmune diseases, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

2020ED 40516001

[00515] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to treat diseases and disorders of the immune system and/or to inhibit or enhance an immune response generated by cells associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1A, column 7 (Tissue Distribution Library Code).

[00516] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, diagnosing, and/or prognosing immunodeficiencies, including both congenital and acquired immunodeficiencies. Examples of B cell immunodeficiencies in which immunoglobulin levels B cell function and/or B cell numbers are decreased include: X-linked agammaglobulinemia (Bruton's disease), X-linked infantile agammaglobulinemia, X-linked immunodeficiency with hyper IgM, non X-linked immunodeficiency with hyper IgM, X-linked lymphoproliferative syndrome (XLP), agammaglobulinemia including congenital and acquired agammaglobulinemia, adult onset agammaglobulinemia, late-onset agammaglobulinemia, dysgammaglobulinemia, hypogammaglobulinemia, unspecified hypogammaglobulinemia, recessive agammaglobulinemia (Swiss type), Selective IgM deficiency, selective IgA deficiency, selective IgG subclass deficiencies, IgG subclass deficiency (with or without IgA deficiency), Ig deficiency with increased IgM, IgG and IgA deficiency with increased IgM, antibody deficiency with normal or elevated Igs, Ig heavy chain deletions, kappa chain deficiency, B cell lymphoproliferative disorder (BLPD), common variable immunodeficiency (CVID), common variable immunodeficiency (CVI) (acquired), and transient hypogammaglobulinemia of infancy.

[00517] In specific embodiments, ataxia-telangiectasia or conditions associated with ataxia-telangiectasia are treated, prevented, diagnosed, and/or prognosing using the polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof.

[00518] Examples of congenital immunodeficiencies in which T cell and/or B cell function and/or number is decreased include, but are not limited to: DiGeorge anomaly, severe combined immunodeficiencies (SCID) (including, but not limited to, X-linked SCID, autosomal recessive SCID, adenosine deaminase deficiency,

purine nucleoside phosphorylase (PNP) deficiency, Class II MHC deficiency (Bare lymphocyte syndrome), Wiskott-Aldrich syndrome, and ataxia telangiectasia), thymic hypoplasia, third and fourth pharyngeal pouch syndrome, 22q11.2 deletion, chronic mucocutaneous candidiasis, natural killer cell deficiency (NK), idiopathic CD4+ T-lymphocytopenia, immunodeficiency with predominant T cell defect (unspecified), and unspecified immunodeficiency of cell mediated immunity.

[00519] In specific embodiments, DiGeorge anomaly or conditions associated with DiGeorge anomaly are treated, prevented, diagnosed, and/or prognosed using polypeptides or polynucleotides of the invention, or antagonists or agonists thereof.

[00520] Other immunodeficiencies that may be treated, prevented, diagnosed, and/or prognosed using polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof, include, but are not limited to, chronic granulomatous disease, Chédiak-Higashi syndrome, myeloperoxidase deficiency, leukocyte glucose-6-phosphate dehydrogenase deficiency, X-linked lymphoproliferative syndrome (XLP), leukocyte adhesion deficiency, complement component deficiencies (including C1, C2, C3, C4, C5, C6, C7, C8 and/or C9 deficiencies), reticular dysgenesis, thymic aplasia, immunodeficiency with thymoma, severe congenital leukopenia, dysplasia with immunodeficiency, neonatal neutropenia, short limbed dwarfism, and Nezelof syndrome-combined immunodeficiency with Igs.

[00521] In a preferred embodiment, the immunodeficiencies and/or conditions associated with the immunodeficiencies recited above are treated, prevented, diagnosed and/or prognosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[00522] In a preferred embodiment polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used as an agent to boost immunoresponsiveness among immunodeficient individuals. In specific embodiments, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used as an agent to boost immunoresponsiveness among B cell and/or T cell immunodeficient individuals.

[00523] The polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, diagnosing and/or prognosing autoimmune disorders. Many autoimmune disorders result from

inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of polynucleotides and polypeptides of the invention that can inhibit an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

[00524] Autoimmune diseases or disorders that may be treated, prevented, diagnosed and/or prognosed by polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, one or more of the following: systemic lupus erythematosus, rheumatoid arthritis, ankylosing spondylitis, multiple sclerosis, autoimmune thyroiditis, Hashimoto's thyroiditis, autoimmune hemolytic anemia, hemolytic anemia, thrombocytopenia, autoimmune thrombocytopenia purpura, autoimmune neonatal thrombocytopenia, idiopathic thrombocytopenia purpura, purpura (e.g., Henloch-Schoenlein purpura), autoimmunocytopenia, Goodpasture's syndrome, Pemphigus vulgaris, myasthenia gravis, Grave's disease (hyperthyroidism), and insulin-resistant diabetes mellitus.

[00525] Additional disorders that are likely to have an autoimmune component that may be treated, prevented, and/or diagnosed with the compositions of the invention include, but are not limited to, type II collagen-induced arthritis, antiphospholipid syndrome, dermatitis, allergic encephalomyelitis, myocarditis, relapsing polychondritis, rheumatic heart disease, neuritis, uveitis ophthalmia, polyendocrinopathies, Reiter's Disease, Stiff-Man Syndrome, autoimmune pulmonary inflammation, autism, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disorders.

[00526] Additional disorders that are likely to have an autoimmune component that may be treated, prevented, diagnosed and/or prognosed with the compositions of the invention include, but are not limited to, scleroderma with anti-collagen antibodies (often characterized, e.g., by nucleolar and other nuclear antibodies), mixed connective tissue disease (often characterized, e.g., by antibodies to extractable nuclear antigens (e.g., ribonucleoprotein)), polymyositis (often characterized, e.g., by nonhistone ANA), pernicious anemia (often characterized, e.g., by antiparietal cell, microsomes, and intrinsic factor antibodies), idiopathic Addison's disease (often characterized, e.g., by humoral and cell-mediated adrenal cytotoxicity,

infertility (often characterized, e.g., by antispermatozoal antibodies), glomerulonephritis (often characterized, e.g., by glomerular basement membrane antibodies or immune complexes), bullous pemphigoid (often characterized, e.g., by IgG and complement in basement membrane), Sjogren's syndrome (often characterized, e.g., by multiple tissue antibodies, and/or a specific nonhistone ANA (SS-B)), diabetes mellitus (often characterized, e.g., by cell-mediated and humoral islet cell antibodies), and adrenergic drug resistance (including adrenergic drug resistance with asthma or cystic fibrosis) (often characterized, e.g., by beta-adrenergic receptor antibodies).

[00527] Additional disorders that may have an autoimmune component that may be treated, prevented, diagnosed and/or prognosed with the compositions of the invention include, but are not limited to, chronic active hepatitis (often characterized, e.g., by smooth muscle antibodies), primary biliary cirrhosis (often characterized, e.g., by mitochondria antibodies), other endocrine gland failure (often characterized, e.g., by specific tissue antibodies in some cases), vitiligo (often characterized, e.g., by melanocyte antibodies), vasculitis (often characterized, e.g., by Ig and complement in vessel walls and/or low serum complement), post-MI (often characterized, e.g., by myocardial antibodies), cardiomyopathy syndrome (often characterized, e.g., by myocardial antibodies), urticaria (often characterized, e.g., by IgG and IgM antibodies to IgE), atopic dermatitis (often characterized, e.g., by IgG and IgM antibodies to IgE), asthma (often characterized, e.g., by IgG and IgM antibodies to IgE), and many other inflammatory, granulomatous, degenerative, and atrophic disorders.

[00528] In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prevented, diagnosed and/or prognosed using for example, antagonists or agonists, polypeptides or polynucleotides, or antibodies of the present invention. In a specific preferred embodiment, rheumatoid arthritis is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[00529] In another specific preferred embodiment, systemic lupus erythematosus is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention. In another

specific preferred embodiment, idiopathic thrombocytopenia purpura is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[00530] In another specific preferred embodiment IgA nephropathy is treated, prevented, and/or diagnosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[00531] In a preferred embodiment, the autoimmune diseases and disorders and/or conditions associated with the diseases and disorders recited above are treated, prevented, diagnosed and/or prognosed using polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention.

[00532] In preferred embodiments, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a immunosuppressive agent(s).

[00533] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be useful in treating, preventing, prognosing, and/or diagnosing diseases, disorders, and/or conditions of hematopoietic cells. Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with a decrease in certain (or many) types hematopoietic cells, including but not limited to, leukopenia, neutropenia, anemia, and thrombocytopenia. Alternatively, Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention could be used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat or prevent those diseases, disorders, and/or conditions associated with an increase in certain (or many) types of hematopoietic cells, including but not limited to, histiocytosis.

[00534] Allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated, prevented, diagnosed and/or prognosed using polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof. Moreover, these molecules can be used to treat, prevent, prognose, and/or diagnose anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.



10091504-030702

[00535] Additionally, polypeptides or polynucleotides of the invention, and/or agonists or antagonists thereof, may be used to treat, prevent, diagnose and/or prognose IgE-mediated allergic reactions. Such allergic reactions include, but are not limited to, asthma, rhinitis, and eczema. In specific embodiments, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to modulate IgE concentrations in vitro or in vivo.

[00536] Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention have uses in the diagnosis, prognosis, prevention, and/or treatment of inflammatory conditions. For example, since polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists of the invention may inhibit the activation, proliferation and/or differentiation of cells involved in an inflammatory response, these molecules can be used to prevent and/or treat chronic and acute inflammatory conditions. Such inflammatory conditions include, but are not limited to, for example, inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome), ischemia-reperfusion injury, endotoxin lethality, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, over production of cytokines (e.g., TNF or IL-1.), respiratory disorders (e.g., asthma and allergy); gastrointestinal disorders (e.g., inflammatory bowel disease); cancers (e.g., gastric, ovarian, lung, bladder, liver, and breast); CNS disorders (e.g., multiple sclerosis; ischemic brain injury and/or stroke, traumatic brain injury, neurodegenerative disorders (e.g., Parkinson's disease and Alzheimer's disease); AIDS-related dementia; and prion disease); cardiovascular disorders (e.g., atherosclerosis, myocarditis, cardiovascular disease, and cardiopulmonary bypass complications); as well as many additional diseases, conditions, and disorders that are characterized by inflammation (e.g., hepatitis, rheumatoid arthritis, gout, trauma, pancreatitis, sarcoidosis, dermatitis, renal ischemia-reperfusion injury, Grave's disease, systemic lupus erythematosus, diabetes mellitus, and allogenic transplant rejection).

[00537] Because inflammation is a fundamental defense mechanism, inflammatory disorders can effect virtually any tissue of the body. Accordingly, polynucleotides, polypeptides, and antibodies of the invention, as well as agonists or antagonists thereof, have uses in the treatment of tissue-specific inflammatory disorders,

including, but not limited to, adrenalitis, alveolitis, angiocholecystitis, appendicitis, balanitis, blepharitis, bronchitis, bursitis, carditis, cellulitis, cervicitis, cholecystitis, chorditis, cochlitis, colitis, conjunctivitis, cystitis, dermatitis, diverticulitis, encephalitis, endocarditis, esophagitis, eustachitis, fibrositis, folliculitis, gastritis, gastroenteritis, gingivitis, glossitis, hepatosplenitis, keratitis, labyrinthitis, laryngitis, lymphangitis, mastitis, media otitis, meningitis, metritis, mucitis, myocarditis, myositis, myringitis, nephritis, neuritis, orchitis, osteochondritis, otitis, pericarditis, peritendonitis, peritonitis, pharyngitis, phlebitis, poliomyelitis, prostatitis, pulpitis, retinitis, rhinitis, salpingitis, scleritis, sclerochoroiditis, scrotitis, sinusitis, spondylitis, steatitis, stomatitis, synovitis, syringitis, tendonitis, tonsillitis, urethritis, and vaginitis.

**[00538]** In specific embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, are useful to diagnose, prognose, prevent, and/or treat organ transplant rejections and graft-versus-host disease. Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. Polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, that inhibit an immune response, particularly the activation, proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD. In specific embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, that inhibit an immune response, particularly the activation, proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing experimental allergic and hyperacute xenograft rejection.

**[00539]** In other embodiments, polypeptides, antibodies, or polynucleotides of the invention, and/or agonists or antagonists thereof, are useful to diagnose, prognose, prevent, and/or treat immune complex diseases, including, but not limited to, serum sickness, post streptococcal glomerulonephritis, polyarteritis nodosa, and immune complex-induced vasculitis.

**[00540]** Polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the invention can be used to treat, detect, and/or prevent infectious agents. For example, by increasing the immune response, particularly increasing the

proliferation activation and/or differentiation of B and/or T cells, infectious diseases may be treated, detected, and/or prevented. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may also directly inhibit the infectious agent (refer to section of application listing infectious agents, etc), without necessarily eliciting an immune response.

[00541] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a vaccine adjuvant that enhances immune responsiveness to an antigen. In a specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance tumor-specific immune responses.

[00542] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-viral immune responses. Anti-viral immune responses that may be enhanced using the compositions of the invention as an adjuvant, include virus and virus associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: AIDS, meningitis, Dengue, EBV, and hepatitis (e.g., hepatitis B). In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a virus, disease, or symptom selected from the group consisting of: HIV/AIDS, respiratory syncytial virus, Dengue, rotavirus, Japanese B encephalitis, influenza A and B, parainfluenza, measles, cytomegalovirus, rabies, Junin, Chikungunya, Rift Valley Fever, herpes simplex, and yellow fever.

[00543] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-bacterial or anti-fungal immune responses. Anti-bacterial or anti-fungal immune responses that may be enhanced using the compositions of the invention as an adjuvant, include bacteria or fungus and bacteria or fungus associated diseases or symptoms described herein or otherwise known in the art. In

specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: tetanus, Diphtheria, botulism, and meningitis type B.

[00544] In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to a bacteria or fungus, disease, or symptom selected from the group consisting of: *Vibrio cholerae*, *Mycobacterium leprae*, *Salmonella typhi*, *Salmonella paratyphi*, *Meisseria meningitidis*, *Streptococcus pneumoniae*, Group B streptococcus, *Shigella spp.*, Enterotoxigenic *Escherichia coli*, Enterohemorrhagic *E. coli*, and *Borrelia burgdorferi*.

[00545] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an adjuvant to enhance anti-parasitic immune responses. Anti-parasitic immune responses that may be enhanced using the compositions of the invention as an adjuvant, include parasite and parasite associated diseases or symptoms described herein or otherwise known in the art. In specific embodiments, the compositions of the invention are used as an adjuvant to enhance an immune response to a parasite. In another specific embodiment, the compositions of the invention are used as an adjuvant to enhance an immune response to Plasmodium (malaria) or Leishmania.

[00546] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed to treat infectious diseases including silicosis, sarcoidosis, and idiopathic pulmonary fibrosis; for example, by preventing the recruitment and activation of mononuclear phagocytes.

[00547] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an antigen for the generation of antibodies to inhibit or enhance immune mediated responses against polypeptides of the invention.

[00548] In one embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are administered to an animal (e.g., mouse, rat, rabbit, hamster, guinea pig, pigs, micro-pig, chicken, camel, goat, horse, cow, sheep, dog, cat, non-human primate, and human, most preferably human) to boost the immune system to produce increased quantities of one or more antibodies (e.g., IgG, IgA, IgM, and IgE), to induce higher affinity antibody production and

immunoglobulin class switching (e.g., IgG, IgA, IgM, and IgE), and/or to increase an immune response.

[00549] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a stimulator of B cell responsiveness to pathogens.

[00550] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an activator of T cells.

[00551] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent that elevates the immune status of an individual prior to their receipt of immunosuppressive therapies.

[00552] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to induce higher affinity antibodies.

[00553] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to increase serum immunoglobulin concentrations.

[00554] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to accelerate recovery of immunocompromised individuals.

[00555] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to boost immunoresponsiveness among aged populations and/or neonates.

[00556] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an immune system enhancer prior to, during, or after bone marrow transplant and/or other transplants (e.g., allogeneic or xenogeneic organ transplantation). With respect to transplantation, compositions of the invention may be administered prior to, concomitant with, and/or after transplantation. In a specific embodiment, compositions of the invention are administered after transplantation, prior to the beginning of recovery of T-cell populations. In another specific embodiment, compositions of the invention are first administered after transplantation after the

beginning of recovery of T cell populations, but prior to full recovery of B cell populations.

[00557] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to boost immunoresponsiveness among individuals having an acquired loss of B cell function. Conditions resulting in an acquired loss of B cell function that may be ameliorated or treated by administering the polypeptides, antibodies, polynucleotides and/or agonists or antagonists thereof, include, but are not limited to, HIV Infection, AIDS, bone marrow transplant, and B cell chronic lymphocytic leukemia (CLL).

[00558] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to boost immunoresponsiveness among individuals having a temporary immune deficiency. Conditions resulting in a temporary immune deficiency that may be ameliorated or treated by administering the polypeptides, antibodies, polynucleotides and/or agonists or antagonists thereof, include, but are not limited to, recovery from viral infections (e.g., influenza), conditions associated with malnutrition, recovery from infectious mononucleosis, or conditions associated with stress, recovery from measles, recovery from blood transfusion, and recovery from surgery.

[00559] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a regulator of antigen presentation by monocytes, dendritic cells, and/or B-cells. In one embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention enhance antigen presentation or antagonizes antigen presentation in vitro or in vivo. Moreover, in related embodiments, said enhancement or antagonism of antigen presentation may be useful as an anti-tumor treatment or to modulate the immune system.

[00560] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an agent to direct an individual's immune system towards development of a humoral response (i.e. TH2) as opposed to a TH1 cellular response.

[00561] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means to induce

tumor proliferation and thus make it more susceptible to anti-neoplastic agents. For example, multiple myeloma is a slowly dividing disease and is thus refractory to virtually all anti-neoplastic regimens. If these cells were forced to proliferate more rapidly their susceptibility profile would likely change.

[00562] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a stimulator of B cell production in pathologies such as AIDS, chronic lymphocyte disorder and/or Common Variable Immunodeficiency.

[00563] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for generation and/or regeneration of lymphoid tissues following surgery, trauma or genetic defect. In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used in the pretreatment of bone marrow samples prior to transplant.

[00564] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a gene-based therapy for genetically inherited disorders resulting in immunoincompetence/immunodeficiency such as observed among SCID patients.

[00565] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of activating monocytes/macrophages to defend against parasitic diseases that effect monocytes such as Leishmania.

[00566] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of regulating secreted cytokines that are elicited by polypeptides of the invention.

[00567] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used in one or more of the applications described herein, as they may apply to veterinary medicine.

[00568] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of blocking various aspects of immune responses to foreign agents or self. Examples of diseases or conditions in which blocking of certain aspects of immune responses may be desired include autoimmune disorders such as lupus, and arthritis, as well as

immuno-responsiveness to skin allergies, inflammation, bowel disease, injury and diseases/disorders associated with pathogens.

[00569] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for preventing the B cell proliferation and Ig secretion associated with autoimmune diseases such as idiopathic thrombocytopenic purpura, systemic lupus erythematosus and multiple sclerosis.

[00570] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as an inhibitor of B and/or T cell migration in endothelial cells. This activity disrupts tissue architecture or cognate responses and is useful, for example in disrupting immune responses, and blocking sepsis.

[00571] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for chronic hypergammaglobulinemia evident in such diseases as monoclonal gammopathy of undetermined significance (MGUS), Waldenstrom's disease, related idiopathic monoclonal gammopathies, and plasmacytomas.

[00572] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be employed for instance to inhibit polypeptide chemotaxis and activation of macrophages and their precursors, and of neutrophils, basophils, B lymphocytes and some T-cell subsets, e.g., activated and CD8 cytotoxic T cells and natural killer cells, in certain autoimmune and chronic inflammatory and infective diseases. Examples of autoimmune diseases are described herein and include multiple sclerosis, and insulin-dependent diabetes.

[00573] The polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed to treat idiopathic hyper-eosinophilic syndrome by, for example, preventing eosinophil production and migration.

[00574] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used to enhance or inhibit complement mediated cell lysis.



[00575] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used to enhance or inhibit antibody dependent cellular cytotoxicity.

[00576] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may also be employed for treating atherosclerosis, for example, by preventing monocyte infiltration in the artery wall.

[00577] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be employed to treat adult respiratory distress syndrome (ARDS).

[00578] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention may be useful for stimulating wound and tissue repair, stimulating angiogenesis, and/or stimulating the repair of vascular or lymphatic diseases or disorders. Additionally, agonists and antagonists of the invention may be used to stimulate the regeneration of mucosal surfaces.

[00579] In a specific embodiment, polynucleotides or polypeptides, and/or agonists thereof are used to diagnose, prognose, treat, and/or prevent a disorder characterized by primary or acquired immunodeficiency, deficient serum immunoglobulin production, recurrent infections, and/or immune system dysfunction. Moreover, polynucleotides or polypeptides, and/or agonists thereof may be used to treat or prevent infections of the joints, bones, skin, and/or parotid glands, blood-borne infections (e.g., sepsis, meningitis, septic arthritis, and/or osteomyelitis), autoimmune diseases (e.g., those disclosed herein), inflammatory disorders, and malignancies, and/or any disease or disorder or condition associated with these infections, diseases, disorders and/or malignancies) including, but not limited to, CVID, other primary immune deficiencies, HIV disease, CLL, recurrent bronchitis, sinusitis, otitis media, conjunctivitis, pneumonia, hepatitis, meningitis, herpes zoster (e.g., severe herpes zoster), and/or pneumocystis carinii. Other diseases and disorders that may be prevented, diagnosed, prognosed, and/or treated with polynucleotides or polypeptides, and/or agonists of the present invention include, but are not limited to, HIV infection, HTLV-BLV infection, lymphopenia, phagocyte bactericidal dysfunction anemia, thrombocytopenia, and hemoglobinuria.

10004504-030306  
[00580] In another embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention are used to treat, and/or diagnose an individual having common variable immunodeficiency disease ("CVID"; also known as "acquired agammaglobulinemia" and "acquired hypogammaglobulinemia") or a subset of this disease.

[00581] In a specific embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention may be used to diagnose, prognose, prevent, and/or treat cancers or neoplasms including immune cell or immune tissue-related cancers or neoplasms. Examples of cancers or neoplasms that may be prevented, diagnosed, or treated by polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention include, but are not limited to, acute myelogenous leukemia, chronic myelogenous leukemia, Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic anemia (ALL) Chronic lymphocyte leukemia, plasmacytomas, multiple myeloma, Burkitt's lymphoma, EBV-transformed diseases, and/or diseases and disorders described in the section entitled "Hyperproliferative Disorders" elsewhere herein.

[00582] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a therapy for decreasing cellular proliferation of Large B-cell Lymphomas.

[00583] In another specific embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are used as a means of decreasing the involvement of B cells and Ig associated with Chronic Myelogenous Leukemia.

[00584] In specific embodiments, the compositions of the invention are used as an agent to boost immunoresponsiveness among B cell immunodeficient individuals, such as, for example, an individual who has undergone a partial or complete splenectomy.

[00585] Antagonists of the invention include, for example, binding and/or inhibitory antibodies, antisense nucleic acids, ribozymes or soluble forms of the polypeptides of the present invention (e.g., Fc fusion protein; see, e.g., Example 9). Agonists of the invention include, for example, binding or stimulatory antibodies, and soluble forms of the polypeptides (e.g., Fc fusion proteins; see, e.g., Example 9). Polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the

present invention may be employed in a composition with a pharmaceutically acceptable carrier, e.g., as described herein.

[00586] In another embodiment, polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention are administered to an animal (including, but not limited to, those listed above, and also including transgenic animals) incapable of producing functional endogenous antibody molecules or having an otherwise compromised endogenous immune system, but which is capable of producing human immunoglobulin molecules by means of a reconstituted or partially reconstituted immune system from another animal (see, e.g., published PCT Application Nos. WO98/24893, WO/9634096, WO/9633735, and WO/9110741). Administration of polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention to such animals is useful for the generation of monoclonal antibodies against the polypeptides, antibodies, polynucleotides and/or agonists or antagonists of the present invention.

#### **Hyperproliferative Disorders**

[00587] Cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists thereof, can be used to treat, prevent, diagnose and/or prognose hyperproliferative diseases, disorders, and/or conditions, including neoplasms.

[00588] In a specific embodiment, cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists thereof, can be used to treat, prevent, and/or diagnose hyperproliferative diseases, disorders, and/or conditions of the cardiovascular system.

[00589] In a preferred embodiment, cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists thereof, can be used to treat, prevent, and/or diagnose cardiovascular system neoplasms.

[00590] Cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists of the invention, may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists thereof, may proliferate other cells, which can inhibit the hyperproliferative disorder.

[00591] For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating,

differentiating, or mobilizing T-cells, hyperproliferative diseases, disorders, and/or conditions can be treated, prevented, and/or diagnosed. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating, preventing, and/or diagnosing hyperproliferative diseases, disorders, and/or conditions, such as a chemotherapeutic agent.

[00592] Examples of hyperproliferative diseases, disorders, and/or conditions that can be treated, prevented, and/or diagnosed by cardiovascular system associated polynucleotides or polypeptides, or agonists or antagonists thereof, include, but are not limited to neoplasms located in the: prostate, colon, abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

[00593] Similarly, other hyperproliferative disorders can also be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention. Examples of such hyperproliferative disorders include, but are not limited to: Acute Childhood Lymphoblastic Leukemia, Acute Lymphoblastic Leukemia, Acute Lymphocytic Leukemia, Acute Myeloid Leukemia, Adrenocortical Carcinoma, Adult (Primary) Hepatocellular Cancer, Adult (Primary) Liver Cancer, Adult Acute Lymphocytic Leukemia, Adult Acute Myeloid Leukemia, Adult Hodgkin's Disease, Adult Hodgkin's Lymphoma, Adult Lymphocytic Leukemia, Adult Non-Hodgkin's Lymphoma, Adult Primary Liver Cancer, Adult Soft Tissue Sarcoma, AIDS-Related Lymphoma, AIDS-Related Malignancies, Anal Cancer, Astrocytoma, Bile Duct Cancer, Bladder Cancer, Bone Cancer, Brain Stem Glioma, Brain Tumors, Breast Cancer, Cancer of the Renal Pelvis and Ureter, Central Nervous System (Primary) Lymphoma, Central Nervous System Lymphoma, Cerebellar Astrocytoma, Cerebral Astrocytoma, Cervical Cancer, Childhood (Primary) Hepatocellular Cancer, Childhood (Primary) Liver Cancer, Childhood Acute Lymphoblastic Leukemia, Childhood Acute Myeloid Leukemia, Childhood Brain Stem Glioma, Childhood Cerebellar Astrocytoma, Childhood Cerebral Astrocytoma, Childhood Extracranial Germ Cell Tumors, Childhood Hodgkin's Disease, Childhood Hodgkin's Lymphoma, Childhood Hypothalamic and Visual Pathway Glioma, Childhood

Lymphoblastic Leukemia, Childhood Medulloblastoma, Childhood Non-Hodgkin's Lymphoma, Childhood Pineal and Supratentorial Primitive Neuroectodermal Tumors, Childhood Primary Liver Cancer, Childhood Rhabdomyosarcoma, Childhood Soft Tissue Sarcoma, Childhood Visual Pathway and Hypothalamic Glioma, Chronic Lymphocytic Leukemia, Chronic Myelogenous Leukemia, Colon Cancer, Cutaneous T-Cell Lymphoma, Endocrine Pancreas Islet Cell Carcinoma, Endometrial Cancer, Ependymoma, Epithelial Cancer, Esophageal Cancer, Ewing's Sarcoma and Related Tumors, Exocrine Pancreatic Cancer, Extracranial Germ Cell Tumor, Extragenital Germ Cell Tumor, Extrahepatic Bile Duct Cancer, Eye Cancer, Female Breast Cancer, Gaucher's Disease, Gallbladder Cancer, Gastric Cancer, Gastrointestinal Carcinoid Tumor, Gastrointestinal Tumors, Germ Cell Tumors, Gestational Trophoblastic Tumor, Hairy Cell Leukemia, Head and Neck Cancer, Hepatocellular Cancer, Hodgkin's Disease, Hodgkin's Lymphoma, Hypergammaglobulinemia, Hypopharyngeal Cancer, Intestinal Cancers, Intraocular Melanoma, Islet Cell Carcinoma, Islet Cell Pancreatic Cancer, Kaposi's Sarcoma, Kidney Cancer, Laryngeal Cancer, Lip and Oral Cavity Cancer, Liver Cancer, Lung Cancer, Lymphoproliferative Disorders, Macroglobulinemia, Male Breast Cancer, Malignant Mesothelioma, Malignant Thymoma, Medulloblastoma, Melanoma, Mesothelioma, Metastatic Occult Primary Squamous Neck Cancer, Metastatic Primary Squamous Neck Cancer, Metastatic Squamous Neck Cancer, Multiple Myeloma, Multiple Myeloma/Plasma Cell Neoplasm, Myelodysplastic Syndrome, Myelogenous Leukemia, Myeloid Leukemia, Myeloproliferative Disorders, Nasal Cavity and Paranasal Sinus Cancer, Nasopharyngeal Cancer, Neuroblastoma, Non-Hodgkin's Lymphoma During Pregnancy, Nonmelanoma Skin Cancer, Non-Small Cell Lung Cancer, Occult Primary Metastatic Squamous Neck Cancer, Oropharyngeal Cancer, Osteo-/Malignant Fibrous Sarcoma, Osteosarcoma/Malignant Fibrous Histiocytoma, Osteosarcoma/Malignant Fibrous Histiocytoma of Bone, Ovarian Epithelial Cancer, Ovarian Germ Cell Tumor, Ovarian Low Malignant Potential Tumor, Pancreatic Cancer, Paraproteinemia, Purpura, Parathyroid Cancer, Penile Cancer, Pheochromocytoma, Pituitary Tumor, Plasma Cell Neoplasm/Multiple Myeloma, Primary Central Nervous System Lymphoma, Primary Liver Cancer, Prostate Cancer, Rectal Cancer, Renal Cell Cancer, Renal Pelvis and Ureter Cancer, Retinoblastoma, Rhabdomyosarcoma,

Salivary Gland Cancer, Sarcoidosis Sarcomas, Sezary Syndrome, Skin Cancer, Small Cell Lung Cancer, Small Intestine Cancer, Soft Tissue Sarcoma, Squamous Neck Cancer, Stomach Cancer, Supratentorial Primitive Neuroectodermal and Pineal Tumors, T-Cell Lymphoma, Testicular Cancer, Thymoma, Thyroid Cancer, Transitional Cell Cancer of the Renal Pelvis and Ureter, Transitional Renal Pelvis and Ureter Cancer, Trophoblastic Tumors, Ureter and Renal Pelvis Cell Cancer, Urethral Cancer, Uterine Cancer, Uterine Sarcoma, Vaginal Cancer, Visual Pathway and Hypothalamic Glioma, Vulvar Cancer, Waldenstrom's Macroglobulinemia, Wilms' Tumor, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

[00594] In another preferred embodiment, polynucleotides or polypeptides, or agonists or antagonists of the present invention are used to diagnose, prognose, prevent, and/or treat premalignant conditions and to prevent progression to a neoplastic or malignant state, including but not limited to those disorders described above. Such uses are indicated in conditions known or suspected of preceding progression to neoplasia or cancer, in particular, where non-neoplastic cell growth consisting of hyperplasia, metaplasia, or most particularly, dysplasia has occurred (for review of such abnormal growth conditions, see Robbins and Angell, 1976, Basic Pathology, 2d Ed., W. B. Saunders Co., Philadelphia, pp. 68-79.)

[00595] Hyperplasia is a form of controlled cell proliferation, involving an increase in cell number in a tissue or organ, without significant alteration in structure or function. Hyperplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, angiofollicular mediastinal lymph node hyperplasia, angiolymphoid hyperplasia with eosinophilia, atypical melanocytic hyperplasia, basal cell hyperplasia, benign giant lymph node hyperplasia, cementum hyperplasia, congenital adrenal hyperplasia, congenital sebaceous hyperplasia, cystic hyperplasia, cystic hyperplasia of the breast, denture hyperplasia, ductal hyperplasia, endometrial hyperplasia, fibromuscular hyperplasia, focal epithelial hyperplasia, gingival hyperplasia, inflammatory fibrous hyperplasia, inflammatory papillary hyperplasia, intravascular papillary endothelial hyperplasia, nodular hyperplasia of prostate, nodular regenerative hyperplasia,

pseudoepitheliomatous hyperplasia, senile sebaceous hyperplasia, and verrucous hyperplasia.

[00596] Metaplasia is a form of controlled cell growth in which one type of adult or fully differentiated cell substitutes for another type of adult cell. Metaplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, agnogenic myeloid metaplasia, apocrine metaplasia, atypical metaplasia, autoparenchymatous metaplasia, connective tissue metaplasia, epithelial metaplasia, intestinal metaplasia, metaplastic anemia, metaplastic ossification, metaplastic polyps, myeloid metaplasia, primary myeloid metaplasia, secondary myeloid metaplasia, squamous metaplasia, squamous metaplasia of amnion, and symptomatic myeloid metaplasia.

[00597] Dysplasia is frequently a forerunner of cancer, and is found mainly in the epithelia; it is the most disorderly form of non-neoplastic cell growth, involving a loss in individual cell uniformity and in the architectural orientation of cells. Dysplastic cells often have abnormally large, deeply stained nuclei, and exhibit pleomorphism. Dysplasia characteristically occurs where there exists chronic irritation or inflammation. Dysplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, anhidrotic ectodermal dysplasia, anterofacial dysplasia, asphyxiating thoracic dysplasia, atriadigital dysplasia, bronchopulmonary dysplasia, cerebral dysplasia, cervical dysplasia, chondroectodermal dysplasia, cleidocranial dysplasia, congenital ectodermal dysplasia, craniodiaphysial dysplasia, craniocarpotarsal dysplasia, craniometaphysial dysplasia, dentin dysplasia, diaphysial dysplasia, ectodermal dysplasia, enamel dysplasia, encephalo-ophthalmic dysplasia, dysplasia epiphysialis hemimelia, dysplasia epiphysialis multiplex, dysplasia epiphysialis punctata, epithelial dysplasia, faciodigitogenital dysplasia, familial fibrous dysplasia of jaws, familial white folded dysplasia, fibromuscular dysplasia, fibrous dysplasia of bone, florid osseous dysplasia, hereditary renal-retinal dysplasia, hidrotic ectodermal dysplasia, hypohidrotic ectodermal dysplasia, lymphopenic thymic dysplasia, mammary dysplasia, mandibulofacial dysplasia, metaphysial dysplasia, Mondini dysplasia, monostotic fibrous dysplasia, mucoepithelial dysplasia, multiple

epiphysial dysplasia, oculoauriculovertebral dysplasia, oculodentodigital dysplasia, oculovertebral dysplasia, odontogenic dysplasia, ophthalmomandibulomelic dysplasia, periapical cemental dysplasia, polyostotic fibrous dysplasia, pseudoachondroplastic spondyloepiphysial dysplasia, retinal dysplasia, septo-optic dysplasia, spondyloepiphysial dysplasia, and ventriculoradial dysplasia.

[00598] Additional pre-neoplastic disorders which can be diagnosed, prognosed, prevented, and/or treated with compositions of the invention (including polynucleotides, polypeptides, agonists or antagonists) include, but are not limited to, benign dysproliferative disorders (e.g., benign tumors, fibrocystic conditions, tissue hypertrophy, intestinal polyps, colon polyps, and esophageal dysplasia), leukoplakia, keratoses, Bowen's disease, Farmer's Skin, solar cheilitis, and solar keratosis.

[00599] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose and/or prognose disorders associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1A, 7 (Tissue Distribution Library Code).

[00600] In another embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat cancers and neoplasms, including, but not limited to those described herein. In a further preferred embodiment, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention conjugated to a toxin or a radioactive isotope, as described herein, may be used to treat acute myelogenous leukemia.

[00601] Additionally, polynucleotides, polypeptides, and/or agonists or antagonists of the invention may affect apoptosis, and therefore, would be useful in treating a number of diseases associated with increased cell survival or the inhibition of apoptosis. For example, diseases associated with increased cell survival or the inhibition of apoptosis that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer,



intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection.

[00602] In preferred embodiments, polynucleotides, polypeptides, and/or agonists or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those listed above.

[00603] Additional diseases or conditions associated with increased cell survival that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma,

glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, emangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

**[00604]** Diseases associated with increased apoptosis that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, retinitis pigmentosa, cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

**[00605]** Hyperproliferative diseases and/or disorders that could be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention, include, but are not limited to, neoplasms located in the liver, abdomen, bone, breast, digestive system, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary, testicles, ovary, thymus, thyroid), eye, head and neck, nervous system (central and peripheral), lymphatic system, pelvis, skin, soft tissue, spleen, thorax, and urogenital tract.

**[00606]** Similarly, other hyperproliferative disorders can also be diagnosed, prognosed, prevented, and/or treated by polynucleotides, polypeptides, and/or agonists or antagonists of the invention. Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenström's macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

[00607] One preferred embodiment utilizes polynucleotides of the present invention to inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

[00608] Thus, the present invention provides a method for treating cell proliferative diseases, disorders, and/or conditions by inserting into an abnormally proliferating cell a polynucleotide of the present invention, wherein said polynucleotide represses said cell proliferation, disease, disorder, and/or condition.

[00609] In a preferred embodiment, the present invention provides a method for treating cell proliferative diseases, disorders and/or conditions of the cardiovascular system by inserting into a cell, a polynucleotide of the present invention, wherein said polynucleotide represses said cell proliferation, disease and/or disorder.

[00610] Another embodiment of the present invention provides a method of treating cell-proliferative diseases, disorders, and/or conditions in individuals comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (see, e.g., G J. Nabel, et. al., PNAS 96: 324-326 (1999), which is hereby incorporated by reference). In a most preferred embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e., magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated (i.e., to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

[00611] Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes "

is intended the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

[00612] For local administration to abnormally proliferating cells, polynucleotides of the present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference. In order to specifically deliver or transfect cells which are abnormally proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

[00613] The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

[00614] By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

10091504-030706

[00615] Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as decreases in the rate of proliferation or growth of the cells. The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

[00616] The present invention is further directed to antibody-based therapies which involve administering of anti-polypeptides and anti-polynucleotide antibodies to a mammalian, preferably human, patient for treating one or more of the described diseases, disorders, and/or conditions. Methods for producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere herein. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

[00617] A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g., as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

[00618] In particular, the antibodies, fragments and derivatives of the present invention are useful for treating a subject having or developing cell proliferative and/or differentiation diseases, disorders, and/or conditions as described herein. Such treatment comprises administering a single or multiple doses of the antibody, or a fragment, derivative; or a conjugate thereof.

[00619] The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or

hematopoietic growth factors, for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

[00620] It is preferred to use high affinity and/or potent *in vivo* inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of diseases, disorders, and/or conditions related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or  $K_d$  less than  $5 \times 10^{-6}M$ ,  $10^{-6}M$ ,  $5 \times 10^{-7}M$ ,  $10^{-7}M$ ,  $5 \times 10^{-8}M$ ,  $10^{-8}M$ ,  $5 \times 10^{-9}M$ ,  $10^{-9}M$ ,  $5 \times 10^{-10}M$ ,  $10^{-10}M$ ,  $5 \times 10^{-11}M$ ,  $10^{-11}M$ ,  $5 \times 10^{-12}M$ ,  $10^{-12}M$ ,  $5 \times 10^{-13}M$ ,  $10^{-13}M$ ,  $5 \times 10^{-14}M$ ,  $10^{-14}M$ ,  $5 \times 10^{-15}M$ , and  $10^{-15}M$ .

[00621] Moreover, cardiovascular system antigen polypeptides of the present invention or fragments thereof, are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (see, e.g., Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference). Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (see, e.g., Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

[00622] Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (see, e.g., Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference). Moreover, in another preferred embodiment of the

present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat. Res. 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem. Biol. Interact. Apr 24;111-112:23-34 (1998), J. Mo. Med. 76(6):402-12 (1998), Int. J. Tissue React. 20(1):3-15 (1998), which are all hereby incorporated by reference).

**[00623]** Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues. Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

**[00624]** In another embodiment, the invention provides a method of delivering compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or anti-cardiovascular system antigen polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Cardiovascular system antigen polypeptides or anti-cardiovascular system antigen polypeptide antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions.

**[00625]** Polypeptides, protein fusions to, or fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and immunogens.

## Urinary System Disorders

- [00626] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose disorders of the urinary system, including but not limited to disorders of the renal system, bladder, ureters, and urethra. Renal disorders include, but are not limited to, kidney failure, nephritis, blood vessel disorders of kidney, metabolic and congenital kidney disorders, urinary disorders of the kidney, autoimmune disorders, sclerosis and necrosis, electrolyte imbalance, and kidney cancers.
- [00627] Kidney failure diseases include, but are not limited to, acute kidney failure, chronic kidney failure, atheroembolic renal failure, and end-stage renal disease. Inflammatory diseases of the kidney include acute glomerulonephritis, postinfectious glomerulonephritis, rapidly progressive glomerulonephritis, nephrotic syndrome, membranous glomerulonephritis, familial nephrotic syndrome, membranoproliferative glomerulonephritis I and II, mesangial proliferative glomerulonephritis, chronic glomerulonephritis, acute tubulointerstitial nephritis, chronic tubulointerstitial nephritis, acute post-streptococcal glomerulonephritis (PSGN), pyelonephritis, lupus nephritis, chronic nephritis, interstitial nephritis, and post-streptococcal glomerulonephritis.
- [00628] Blood vessel disorders of the kidneys include, but are not limited to, kidney infarction, atheroembolic kidney disease, cortical necrosis, malignant nephrosclerosis, renal vein thrombosis, renal underperfusion, renal ischemia-reperfusion, renal artery embolism, and renal artery stenosis. Kidney disorders resulting from urinary tract problems include, but are not limited to, pyelonephritis, hydronephrosis, urolithiasis (renal lithiasis, nephrolithiasis), reflux nephropathy, urinary tract infections, urinary retention, and acute or chronic unilateral obstructive uropathy.
- [00629] Metabolic and congenital disorders of the kidneys include, but are not limited to, renal tubular acidosis, renal glycosuria, nephrogenic diabetes insipidus, cystinuria, Fanconi's syndrome, vitamin D-resistant rickets, Hartnup disease, Bartter's syndrome, Liddle's syndrome, polycystic kidney disease, medullary cystic disease, medullary sponge kidney, Alport's syndrome, nail-patella syndrome, congenital nephrotic syndrome, CRUSH syndrome, horseshoe kidney, diabetic nephropathy, nephrogenic diabetes insipidus, analgesic nephropathy, kidney stones, and



membranous nephropathy, Kidney disorders resulting from an autoimmune response include, but are not limited to, systemic lupus erythematosus (SLE), Goodpasture syndrome, IgA nephropathy, and IgM mesangial proliferative glomerulonephritis.

[00630] Sclerotic or necrotic disorders of the kidney include, but are not limited to, glomerulosclerosis, diabetic nephropathy, focal segmental glomerulosclerosis (FSGS), necrotizing glomerulonephritis, and renal papillary necrosis. Kidneys may also develop carcinomas, including, but not limited to, hypernephroma, nephroblastoma, renal cell cancer, transitional cell cancer, squamous cell cancer, and Wilm's tumor.

[00631] Kidney disorders may also result in electrolyte imbalances, including, but not limited to, nephrocalcinosis, pyuria, edema, hydronephritis, proteinuria, hyponatremia, hypernatremia, hypokalemia, hyperkalemia, hypocalcemia, hypercalcemia, hypophosphatemia, and hyperphosphatemia.

[00632] Bladder disorders include, but are not limited to, benign prostatic hyperplasia (BPH), interstitial cystitis (IC), prostatitis, proteinuria, urinary tract infections, urinary incontinence, urinary retention. Disorders of the ureters and urethra include, but are not limited to, acute or chronic unilateral obstructive uropathy. The bladder, ureters, and urethra may also develop carcinomas, including, but not limited to, superficial bladder cancer, invasive bladder cancer, carcinoma of the ureter, and urethra cancers.

[00633] Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

## Respiratory Disorders

- [00634] Polynucleotides or polypeptides, or agonists or antagonists of the present invention may be used to treat, prevent, diagnose, and/or prognose diseases and/or disorders of the respiratory system.
- [00635] Diseases and disorders of the respiratory system include, but are not limited to, nasal vestibulitis, nonallergic rhinitis (e.g., acute rhinitis, chronic rhinitis, atrophic rhinitis, vasomotor rhinitis), nasal polyps, and sinusitis, juvenile angiofibromas, cancer of the nose and juvenile papillomas, vocal cord polyps, nodules (singer's nodules), contact ulcers, vocal cord paralysis, laryngoceles, pharyngitis (e.g., viral and bacterial), tonsillitis, tonsillar cellulitis, parapharyngeal abscess, laryngitis, laryngoceles, and throat cancers (e.g., cancer of the nasopharynx, tonsil cancer, larynx cancer), lung cancer (e.g., squamous cell carcinoma, small cell (oat cell) carcinoma, large cell carcinoma, and adenocarcinoma), allergic disorders (eosinophilic pneumonia, hypersensitivity pneumonitis (e.g., extrinsic allergic alveolitis, allergic interstitial pneumonitis, organic dust pneumoconiosis, allergic bronchopulmonary aspergillosis, asthma, Wegener's granulomatosis (granulomatous vasculitis), Goodpasture's syndrome)), pneumonia (e.g., bacterial pneumonia (e.g., *Streptococcus pneumoniae* (pneumococcal pneumonia), *Staphylococcus aureus* (staphylococcal pneumonia), Gram-negative bacterial pneumonia (caused by, e.g., *Klebsiella* and *Pseudomonas* spp.), *Mycoplasma pneumoniae* pneumonia, *Hemophilus influenzae* pneumonia, *Legionella pneumophila* (Legionnaires' disease), and *Chlamydia psittaci* (Psittacosis)), and viral pneumonia (e.g., influenza, chickenpox (varicella).
- [00636] Additional diseases and disorders of the respiratory system include, but are not limited to bronchiolitis, polio (poliomyelitis), croup, respiratory syncytial viral infection, mumps, erythema infectiosum (fifth disease), roseola infantum, progressive rubella panencephalitis, german measles, and subacute sclerosing panencephalitis), fungal pneumonia (e.g., Histoplasmosis, Coccidioidomycosis, Blastomycosis, fungal infections in people with severely suppressed immune systems (e.g., cryptococcosis, caused by *Cryptococcus neoformans*; aspergillosis, caused by *Aspergillus* spp.; candidiasis, caused by *Candida*; and mucormycosis)), *Pneumocystis carinii* (pneumocystis pneumonia), atypical pneumonias (e.g., *Mycoplasma* and *Chlamydia* spp.), opportunistic infection pneumonia, nosocomial

10091504-0300-0001

pneumonia, chemical pneumonitis, and aspiration pneumonia, pleural disorders (e.g., pleurisy, pleural effusion, and pneumothorax (e.g., simple spontaneous pneumothorax, complicated spontaneous pneumothorax, tension pneumothorax)), obstructive airway diseases (e.g., asthma, chronic obstructive pulmonary disease (COPD), emphysema, chronic or acute bronchitis), occupational lung diseases (e.g., silicosis, black lung (coal workers' pneumoconiosis), asbestosis, berylliosis, occupational asthma, byssinosis, and benign pneumoconioses), Infiltrative Lung Disease (e.g., pulmonary fibrosis (e.g., fibrosing alveolitis, usual interstitial pneumonia), idiopathic pulmonary fibrosis, desquamative interstitial pneumonia, lymphoid interstitial pneumonia, histiocytosis X (e.g., Letterer-Siwe disease, Hand-Schüller-Christian disease, eosinophilic granuloma), idiopathic pulmonary hemosiderosis, sarcoidosis and pulmonary alveolar proteinosis), Acute respiratory distress syndrome (also called, e.g., adult respiratory distress syndrome), edema, pulmonary embolism, bronchitis (e.g., viral, bacterial), bronchiectasis, atelectasis, lung abscess (caused by, e.g., *Staphylococcus aureus* or *Legionella pneumophila*), and cystic fibrosis.

#### Anti-Angiogenesis Activity

[00637] The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.* 29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-

743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

[00638] The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating an angiogenesis-related disease and/or disorder, comprising administration to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat a cancer or tumor. Cancers which may be treated with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors, including prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, colon, rectum, cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

[00639] Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the

appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

[00640] Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating other disorders, besides cancers, which involve angiogenesis. These disorders include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

[00641] For example, within one aspect of the present invention methods are provided for treating hypertrophic scars and keloids, comprising the step of administering a polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

[00642] Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists of the invention are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

[00643] Moreover, ocular disorders associated with neovascularization which can be treated with the polynucleotides and polypeptides of the present invention (including

agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases, ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalm.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalm.* 22:291-312 (1978).

[00644] Thus, within one aspect of the present invention methods are provided for treating neovascular diseases of the eye such as corneal neovascularization (including corneal graft neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue, which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized, it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of disorders can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

[00645] Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic composition is prepared with a muco-adhesive polymer, which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the

treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

[00646] Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the goal of the administration would be to place the composition at the advancing front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation, the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form, injections might only be required 2-3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

[00647] Within another aspect of the present invention, methods are provided for treating neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

[00648] Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment

should be initiated prior to the acquisition of severe disease requiring photocoagulation.

[00649] Within another aspect of the present invention, methods are provided for treating retrolental fibroplasia, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

[00650] Additionally, disorders which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

[00651] Moreover, disorders and/or states, which can be treated, prevented, diagnosed and/or prognosed with the polynucleotides, polypeptides, agonists and/or antagonists of the invention include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (*Rochelina minialia quintosa*), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.



[00652] In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

[00653] Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

[00654] Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered via endoscopic procedures in order to coat tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes, which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized during abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

[00655] Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide, agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to administration.

Within particularly preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

[00656] Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

[00657] The polynucleotides, polypeptides, agonists and/or agonists of the present invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

[00658] Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

[00659] Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

[00660] Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl

complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

[00661] A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26 (1991)); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326 (1992)); Chymostatin (Tomkinson et al., Biochem J. 286:475-480 (1992)); Cyclodextrin Tetradasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557 (1990)); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446 (1987)); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664 (1987)); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4- chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolmidazole; and metalloproteinase inhibitors such as BB94.

#### **Musculoskeletal System Disorders**

[00662] Polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose disorders of the musculoskeletal system, including but not limited to, disorders of the bone,

joints, ligaments, tendons, bursa, muscle, and/or neoplasms and cancers associated with musculoskeletal tissue.

[00663] Diseases or disorders of the bone include, but are not limited to, Albers-Schönberg disease, bowlegs, heel spurs, Köhler's bone disease, knock-knees, Legg-Calvé-Perthes disease, Marfan's syndrome, mucopolysaccharidoses, Osgood-Schlatter disease, osteochondroses, osteochondrodysplasia, osteomyelitis, osteopetroses, osteoporosis (postmenopausal, senile, and juvenile), Paget's disease, Scheuermann's disease, scoliosis, Sever's disease, and patellofemoral stress syndrome.

[00664] Joint diseases or disorders include, but are not limited to, ankylosing spondylitis, Behçet's syndrome, CREST syndrome, Ehlers-Danlos syndrome, infectious arthritis, discoid lupus erythematosus, systemic lupus erythematosus, Lyme disease, osteoarthritis, psoriatic arthritis, relapsing polychondrites, Reiter's syndrome, rheumatoid arthritis (adult and juvenile), scleroderma, and Still's disease.

[00665] Diseases or disorders affecting ligaments, tendons, or bursa include, but are not limited to, ankle sprain, bursitis, posterior Achilles tendon bursitis (Haglund's deformity), anterior Achilles tendon bursitis (Albert's disease), tendinitis, tenosynovitis, popliteus tendinitis, Achilles tendinitis, medial or lateral epicondylitis, rotator cuff tendinitis, spasmodic torticollis, and fibromyalgia syndrome.

[00666] Muscle diseases or disorders include, but are not limited to, Becker's muscular dystrophy, Duchenne's muscular dystrophy, Landouzy-Dejerine muscular dystrophy, Leyden-Möbius muscular dystrophy, Erb's muscular dystrophy, Charcot's joints, dermatomyositis, gout, pseudogout, glycogen storage diseases, Pompe's disease, mitochondrial myopathy, periodic paralysis, polymyalgia rheumatica, polymyositis, Steinert's disease, Thomsen's disease, anterolateral and posteromedial shin splints, posterior femoral muscle strain, and fibromyositis.

[00667] Musculoskeletal tissue may also develop cancers and/or neoplasms that include, but are not limited to, osteochondroma, benign chondroma, chondroblastoma, chondromyxoid fibroma, osteoid osteoma, giant cell tumor, multiple myeloma, osteosarcoma, fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's tumor, and malignant lymphoma of bone.

#### Neural Activity and Neurological Diseases

10061504, 030702

[00668] The polynucleotides, polypeptides and agonists or antagonists of the invention may be used for the diagnosis and/or treatment of diseases, disorders, damage or injury of the brain and/or nervous system. Nervous system disorders that can be treated with the compositions of the invention (e.g., polypeptides, polynucleotides, and/or agonists or antagonists), include, but are not limited to, nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the methods of the invention, include but are not limited to, the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems: (1) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia; (2) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries; (3) malignant lesions, in which a portion of the nervous system is destroyed or injured by malignant tissue which is either a nervous system associated malignancy or a malignancy derived from non-nervous system tissue; (4) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, or syphilis; (5) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to, degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis (ALS); (6) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including, but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration; (7) neurological lesions associated with systemic diseases including, but not limited to, diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis; (8) lesions caused by toxic substances

including alcohol, lead, or particular neurotoxins; and (9) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including, but not limited to, multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

[00669] In one embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of hypoxia. In a further preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to protect neural cells from the damaging effects of cerebral hypoxia. According to this embodiment, the compositions of the invention are used to treat or prevent neural cell injury associated with cerebral hypoxia. In one non-exclusive aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention, are used to treat or prevent neural cell injury associated with cerebral ischemia. In another non-exclusive aspect of this embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with cerebral infarction.

[00670] In another preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a stroke. In a specific embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent cerebral neural cell injury associated with a stroke.

[00671] In another preferred embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent neural cell injury associated with a heart attack. In a specific embodiment, the polypeptides, polynucleotides, or agonists or antagonists of the invention are used to treat or prevent cerebral neural cell injury associated with a heart attack.

[00672] The compositions of the invention which are useful for treating or preventing a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, compositions of the invention which elicit any of the following effects may be useful according to the invention: (1) increased survival time of neurons in culture either in the presence or absence of hypoxia or hypoxic conditions; (2)

increased sprouting of neurons in culture or *in vivo*; (3) increased production of a neuron-associated molecule in culture or *in vivo*, e.g., choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or (4) decreased symptoms of neuron dysfunction *in vivo*. Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may routinely be measured using a method set forth herein or otherwise known in the art, such as, for example, in Zhang *et al.*, *Proc Natl Acad Sci USA* 97:3637-42 (2000) or in Arakawa *et al.*, *J. Neurosci.*, 10:3507-15 (1990); increased sprouting of neurons may be detected by methods known in the art, such as, for example, the methods set forth in Pestronk *et al.*, *Exp. Neurol.*, 70:65-82 (1980), or Brown *et al.*, *Ann. Rev. Neurosci.*, 4:17-42 (1981); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, etc., using techniques known in the art and depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, e.g., weakness, motor neuron conduction velocity, or functional disability.

[00673] In specific embodiments, motor neuron disorders that may be treated according to the invention include, but are not limited to, disorders such as infarction, infection, exposure to toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including, but not limited to, progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motor Sensory Neuropathy (Charcot-Marie-Tooth Disease).

[00674] Further, polypeptides or polynucleotides of the invention may play a role in neuronal survival; synapse formation; conductance; neural differentiation, etc. Thus, compositions of the invention (including polynucleotides, polypeptides, and agonists or antagonists) may be used to diagnose and/or treat or prevent diseases or disorders associated with these roles, including, but not limited to, learning and/or cognition disorders. The compositions of the invention may also be useful in the treatment or prevention of neurodegenerative disease states and/or behavioural disorders. Such

neurodegenerative disease states and/or behavioral disorders include, but are not limited to, Alzheimer's Disease, Parkinson's Disease, Huntington's Disease, Tourette Syndrome, schizophrenia, mania, dementia, paranoia, obsessive compulsive disorder, panic disorder, learning disabilities, ALS, psychoses, autism, and altered behaviors, including disorders in feeding, sleep patterns, balance, and perception. In addition, compositions of the invention may also play a role in the treatment, prevention and/or detection of developmental disorders associated with the developing embryo, or sexually-linked disorders.

[00675] Additionally, polypeptides, polynucleotides and/or agonists or antagonists of the invention, may be useful in protecting neural cells from diseases, damage, disorders, or injury, associated with cerebrovascular disorders including, but not limited to, carotid artery diseases (e.g., carotid artery thrombosis, carotid stenosis, or Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis (e.g., carotid artery thrombosis, sinus thrombosis, or Wallenberg's Syndrome), cerebral hemorrhage (e.g., epidural or subdural hematoma, or subarachnoid hemorrhage), cerebral infarction, cerebral ischemia (e.g., transient cerebral ischemia, Subclavian Steal Syndrome, or vertebrobasilar insufficiency), vascular dementia (e.g., multi-infarct), leukomalacia, periventricular, and vascular headache (e.g., cluster headache or migraines).

[00676] In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate neurological cell proliferation and/or differentiation. Therefore, polynucleotides, polypeptides, agonists and/or antagonists of the invention may be used to treat and/or detect neurologic diseases. Moreover, polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used as a marker or detector of a particular nervous system disease or disorder.

[00677] Examples of neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include brain diseases, such as metabolic brain diseases which includes phenylketonuria such as maternal phenylketonuria, pyruvate carboxylase deficiency, pyruvate dehydrogenase complex deficiency, Wernicke's Encephalopathy, brain



edema, brain neoplasms such as cerebellar neoplasms which include infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms, supratentorial neoplasms, canavan disease, cerebellar diseases such as cerebellar ataxia which include spinocerebellar degeneration such as ataxia telangiectasia, cerebellar dyssynergia, Friederich's Ataxia, Machado-Joseph Disease, olivopontocerebellar atrophy, cerebellar neoplasms such as infratentorial neoplasms, diffuse cerebral sclerosis such as encephalitis periaxialis, globoid cell leukodystrophy, metachromatic leukodystrophy and subacute sclerosing panencephalitis.

[00678] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include cerebrovascular disorders (such as carotid artery diseases which include carotid artery thrombosis, carotid stenosis and Moyamoya Disease), cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis such as carotid artery thrombosis, sinus thrombosis and Wallenberg's Syndrome, cerebral hemorrhage such as epidural hematoma, subdural hematoma and subarachnoid hemorrhage, cerebral infarction, cerebral ischemia such as transient cerebral ischemia, Subclavian Steal Syndrome and vertebrobasilar insufficiency, vascular dementia such as multi-infarct dementia, periventricular leukomalacia, vascular headache such as cluster headache and migraine.

[00679] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include dementia such as AIDS Dementia Complex, presenile dementia such as Alzheimer's Disease and Creutzfeldt-Jakob Syndrome, senile dementia such as Alzheimer's Disease and progressive supranuclear palsy, vascular dementia such as multi-infarct dementia, encephalitis which include encephalitis periaxialis, viral encephalitis such as epidemic encephalitis, Japanese Encephalitis, St. Louis Encephalitis, tick-borne encephalitis and West Nile Fever, acute disseminated encephalomyelitis, meningoencephalitis such as uveomeningoencephalitic syndrome, Postencephalitic Parkinson Disease and subacute sclerosing panencephalitis, encephalomalacia such as periventricular leukomalacia, epilepsy such as generalized epilepsy which includes infantile spasms, absence epilepsy,

myoclonic epilepsy which includes MERRF Syndrome, tonic-clonic epilepsy, partial epilepsy such as complex partial epilepsy, frontal lobe epilepsy and temporal lobe epilepsy, post-traumatic epilepsy, status epilepticus such as Epilepsia Partialis Continua, and Hallervorden-Spatz Syndrome.

[00680] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include hydrocephalus such as Dandy-Walker Syndrome and normal pressure hydrocephalus, hypothalamic diseases such as hypothalamic neoplasms, cerebral malaria, narcolepsy which includes cataplexy, bulbar poliomyelitis, cerebri pseudotumor, Rett Syndrome, Reye's Syndrome, thalamic diseases, cerebral toxoplasmosis, intracranial tuberculoma and Zellweger Syndrome, central nervous system infections such as AIDS Dementia Complex, Brain Abscess, subdural empyema, encephalomyelitis such as Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, Necrotizing Hemorrhagic Encephalomyelitis, Visna, and cerebral malaria.

[00681] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include meningitis such as arachnoiditis, aseptic meningitis such as viral meningitis which includes lymphocytic choriomeningitis, Bacterial meningitis which includes Haemophilus Meningitis, Listeria Meningitis, Meningococcal Meningitis such as Waterhouse-Friderichsen Syndrome, Pneumococcal Meningitis and meningeal tuberculosis, fungal meningitis such as Cryptococcal Meningitis, subdural effusion, meningoencephalitis such as uvemeningoencephalitic syndrome, myelitis such as transverse myelitis, neurosyphilis such as tabes dorsalis, poliomyelitis which includes bulbar poliomyelitis and postpoliomyelitis syndrome, prion diseases (such as Creutzfeldt-Jakob Syndrome, Bovine Spongiform Encephalopathy, Gerstmann-Straussler Syndrome, Kuru, Scrapie), and cerebral toxoplasmosis.

[00682] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include central nervous system neoplasms such as brain neoplasms that include cerebellar neoplasms such as infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms and supratentorial

neoplasms, meningeal neoplasms, spinal cord neoplasms which include epidural neoplasms, demyelinating diseases such as Canavan Diseases, diffuse cerebral sclerolosis which includes adrenoleukodystrophy, encephalitis periaxialis, globoid cell leukodystrophy, diffuse cerebral sclerosis such as metachromatic leukodystrophy, allergic encephalomyelitis, necrotizing hemorrhagic encephalomyelitis, progressive multifocal leukoencephalopathy, multiple sclerosis, central pontine myelinolysis, transverse myelitis, neuromyelitis optica, Scrapie, Swayback, Chronic Fatigue Syndrome, Visna, High Pressure Nervous Syndrome, Meningism, spinal cord diseases such as amyotonia congenita, amyotrophic lateral sclerosis, spinal muscular atrophy such as Werdnig-Hoffmann Disease, spinal cord compression, spinal cord neoplasms such as epidural neoplasms, syringomyelia, Tabes Dorsalis, Stiff-Man Syndrome, mental retardation such as Angelman Syndrome, Cri-du-Chat Syndrome, De Lange's Syndrome, Down Syndrome, Gangliosidoses such as gangliosidoses G(M1), Sandhoff Disease, Tay-Sachs Disease, Hartnup Disease, homocystinuria, Laurence-Moon- Biedl Syndrome, Lesch-Nyhan Syndrome, Maple Syrup Urine Disease, mucopolipidosis such as fucosidosis, neuronal ceroid-lipofuscinosis, oculocerebrorenal syndrome, phenylketonuria such as maternal phenylketonuria, Prader-Willi Syndrome, Rett Syndrome, Rubinstein-Taybi Syndrome, Tuberous Sclerosis, WAGR Syndrome, nervous system abnormalities such as holoprosencephaly, neural tube defects such as anencephaly which includes hydrangencephaly, Arnold-Chairi Deformity, encephalocele, meningocele, meningomyelocele, spinal dysraphism such as spina bifida cystica and spina bifida occulta.

**[00683]** Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include hereditary motor and sensory neuropathies which include Charcot-Marie Disease, Hereditary optic atrophy, Refsum's Disease, hereditary spastic paraplegia, Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies such as Congenital Analgesia and Familial Dysautonomia, Neurologic manifestations (such as agnosia that include Gerstmann's Syndrome, Amnesia such as retrograde amnesia, apraxia, neurogenic bladder, cataplexy, communicative disorders such as hearing disorders that includes deafness, partial hearing loss, loudness recruitment and tinnitus, language disorders such as aphasia which include agraphia, anomia,

broca aphasia, and Wernicke Aphasia, Dyslexia such as Acquired Dyslexia, language development disorders, speech disorders such as aphasia which includes anomia, broca aphasia and Wernicke Aphasia, articulation disorders, communicative disorders such as speech disorders which include dysarthria, echolalia, mutism and stuttering, voice disorders such as aphonia and hoarseness, decerebrate state, delirium, fasciculation, hallucinations, meningism, movement disorders such as angelman syndrome, ataxia, athetosis, chorea, dystonia, hypokinesia, muscle hypotonia, myoclonus, tic, torticollis and tremor, muscle hypertonia such as muscle rigidity such as stiff-man syndrome, muscle spasticity, paralysis such as facial paralysis which includes Herpes Zoster Oticus, Gastroparesis, Hemiplegia, ophthalmoplegia such as diplopia, Duane's Syndrome, Horner's Syndrome, Chronic progressive external ophthalmoplegia such as Kearns Syndrome, Bulbar Paralysis, Tropical Spastic Paraparesis, Paraplegia such as Brown-Sequard Syndrome, quadriplegia, respiratory paralysis and vocal cord paralysis, paresis, phantom limb, taste disorders such as ageusia and dysgeusia, vision disorders such as amblyopia, blindness, color vision defects, diplopia, hemianopsia, scotoma and subnormal vision, sleep disorders such as hypersomnia which includes Kleine-Levin Syndrome, insomnia, and somnambulism, spasm such as trismus, unconsciousness such as coma, persistent vegetative state and syncope and vertigo, neuromuscular diseases such as amyotonia congenita, amyotrophic lateral sclerosis, Lambert-Eaton Myasthenic Syndrome, motor neuron disease, muscular atrophy such as spinal muscular atrophy, Charcot-Marie Disease and Werdnig-Hoffmann Disease, Postpoliomyelitis Syndrome, Muscular Dystrophy, Myasthenia Gravis, Myotonia Atrophica, Myotonia Confenita, Nemaline Myopathy, Familial Periodic Paralysis, Multiplex Paramyoclonus, Tropical Spastic Paraparesis and Stiff-Man Syndrome, peripheral nervous system diseases such as acrodynia, amyloid neuropathies, autonomic nervous system diseases such as Adie's Syndrome, Barre-Lieou Syndrome, Familial Dysautonomia, Horner's Syndrome, Reflex Sympathetic Dystrophy and Shy-Drager Syndrome, Cranial Nerve Diseases such as Acoustic Nerve Diseases such as Acoustic Neuroma which includes Neurofibromatosis 2, Facial Nerve Diseases such as Facial Neuralgia, Melkersson-Rosenthal Syndrome, ocular motility disorders which includes amblyopia, nystagmus, oculomotor nerve paralysis, ophthalmoplegia such as Duane's Syndrome, Horner's Syndrome, Chronic

Progressive External Ophthalmoplegia which includes Kearns Syndrome, Strabismus such as Esotropia and Exotropia, Oculomotor Nerve Paralysis, Optic Nerve Diseases such as Optic Atrophy which includes Hereditary Optic Atrophy, Optic Disk Drusen, Optic Neuritis such as Neuromyelitis Optica, Papilledema, Trigeminal Neuralgia, Vocal Cord Paralysis, Demyelinating Diseases such as Neuromyelitis Optica and Swayback, and Diabetic neuropathies such as diabetic foot.

[00684] Additional neurologic diseases which can be treated or detected with polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include nerve compression syndromes such as carpal tunnel syndrome, tarsal tunnel syndrome, thoracic outlet syndrome such as cervical rib syndrome, ulnar nerve compression syndrome, neuralgia such as causalgia, cervico-brachial neuralgia, facial neuralgia and trigeminal neuralgia, neuritis such as experimental allergic neuritis, optic neuritis, polyneuritis, polyradiculoneuritis and radiculities such as polyradiculitis, hereditary motor and sensory neuropathies such as Charcot-Marie Disease, Hereditary Optic Atrophy, Refsum's Disease, Hereditary Spastic Paraplegia and Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies which include Congenital Analgesia and Familial Dysautonomia, POEMS Syndrome, Sciatica, Gustatory Sweating and Tetany).

### **Endocrine Disorders**

[00685] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose disorders and/or diseases related to hormone imbalance, and/or disorders or diseases of the endocrine system.

[00686] Hormones secreted by the glands of the endocrine system control physical growth, sexual function, metabolism, and other functions. Disorders may be classified in two ways: disturbances in the production of hormones, and the inability of tissues to respond to hormones. The etiology of these hormone imbalance or endocrine system diseases, disorders or conditions may be genetic, somatic, such as cancer and some autoimmune diseases, acquired (e.g., by chemotherapy, injury or toxins), or infectious. Moreover, polynucleotides, polypeptides, antibodies, and/or agonists or antagonists of the present invention can be used as a marker or detector

of a particular disease or disorder related to the endocrine system and/or hormone imbalance.

[00687] Endocrine system and/or hormone imbalance and/or diseases encompass disorders of uterine motility including, but not limited to: complications with pregnancy and labor (e.g., pre-term labor, post-term pregnancy, spontaneous abortion, and slow or stopped labor); and disorders and/or diseases of the menstrual cycle (e.g., dysmenorrhea and endometriosis).

[00688] Endocrine system and/or hormone imbalance disorders and/or diseases include disorders and/or diseases of the pancreas, such as, for example, diabetes mellitus, diabetes insipidus, congenital pancreatic agenesis, pheochromocytoma--islet cell tumor syndrome; disorders and/or diseases of the adrenal glands such as, for example, Addison's Disease, corticosteroid deficiency, virilizing disease, hirsutism, Cushing's Syndrome, hyperaldosteronism, pheochromocytoma; disorders and/or diseases of the pituitary gland, such as, for example, hyperpituitarism, hypopituitarism, pituitary dwarfism, pituitary adenoma, panhypopituitarism, acromegaly, gigantism; disorders and/or diseases of the thyroid, including but not limited to, hyperthyroidism, hypothyroidism, Plummer's disease, Graves' disease (toxic diffuse goiter), toxic nodular goiter, thyroiditis (Hashimoto's thyroiditis, subacute granulomatous thyroiditis, and silent lymphocytic thyroiditis), Pendred's syndrome, myxedema, cretinism, thyrotoxicosis, thyroid hormone coupling defect, thymic aplasia, Hurthle cell tumours of the thyroid, thyroid cancer, thyroid carcinoma, Medullary thyroid carcinoma; disorders and/or diseases of the parathyroid, such as, for example, hyperparathyroidism, hypoparathyroidism; disorders and/or diseases of the hypothalamus.

[00689] In addition, endocrine system and/or hormone imbalance disorders and/or diseases may also include disorders and/or diseases of the testes or ovaries, including cancer. Other disorders and/or diseases of the testes or ovaries further include, for example, ovarian cancer, polycystic ovary syndrome, Klinefelter's syndrome, vanishing testes syndrome (bilateral anorchia), congenital absence of Leydig's cells, cryptorchidism, Noonan's syndrome, myotonic dystrophy, capillary haemangioma of the testis (benign), neoplasias of the testis and neo-testis.

[00690] Moreover, endocrine system and/or hormone imbalance disorders and/or diseases may also include disorders and/or diseases such as, for example,

polyglandular deficiency syndromes, pheochromocytoma, neuroblastoma, multiple Endocrine neoplasia, and disorders and/or cancers of endocrine tissues.

[00691] In another embodiment, a polypeptide of the invention, or polynucleotides, antibodies, agonists, or antagonists corresponding to that polypeptide, may be used to diagnose, prognose, prevent, and/or treat endocrine diseases and/or disorders associated with the tissue(s) in which the polypeptide of the invention is expressed, including one, two, three, four, five, or more tissues disclosed in Table 1A, column 7 (Tissue Distribution Library Code).

### **Gastrointestinal Disorders**

[00692] Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat, prevent, diagnose, and/or prognose gastrointestinal disorders, including inflammatory diseases and/or conditions, infections, cancers (e.g., intestinal neoplasms (carcinoid tumor of the small intestine, non-Hodgkin's lymphoma of the small intestine, small bowel lymphoma)), and ulcers, such as peptic ulcers.

[00693] Gastrointestinal disorders include dysphagia, odynophagia, inflammation of the esophagus, peptic esophagitis, gastric reflux, submucosal fibrosis and stricturing, Mallory-Weiss lesions, leiomyomas, lipomas, epidermal cancers, adeoncarcinomas, gastric retention disorders, gastroenteritis, gastric atrophy, gastric/stomach cancers, polyps of the stomach, autoimmune disorders such as pernicious anemia, pyloric stenosis, gastritis (bacterial, viral, eosinophilic, stress-induced, chronic erosive, atrophic, plasma cell, and Ménétrier's), and peritoneal diseases (e.g., chyloperitoneum, hemoperitoneum, mesenteric cyst, mesenteric lymphadenitis, mesenteric vascular occlusion, panniculitis, neoplasms, peritonitis, pneumoperitoneum, bubphrenic abscess).

[00694] Gastrointestinal disorders also include disorders associated with the small intestine, such as malabsorption syndromes, distension, irritable bowel syndrome, sugar intolerance, celiac disease, duodenal ulcers, duodenitis, tropical sprue, Whipple's disease, intestinal lymphangiectasia, Crohn's disease, appendicitis, obstructions of the ileum, Meckel's diverticulum, multiple diverticula, failure of complete rotation of the small and large intestine, lymphoma, and bacterial and

parasitic diseases (such as Traveler's diarrhea, typhoid and paratyphoid, cholera, infection by Roundworms (*Ascariasis lumbricoides*), Hookworms (*Ancylostoma duodenale*), Threadworms (*Enterobius vermicularis*), Tapeworms (*Taenia saginata*, *Echinococcus granulosus*, *Diphyllobothrium spp.*, and *T. solium*).

- [00695] Liver diseases and/or disorders include intrahepatic cholestasis (alagille syndrome, biliary liver cirrhosis), fatty liver (alcoholic fatty liver, reye syndrome), hepatic vein thrombosis, hepatolentricular degeneration, hepatomegaly, hepatopulmonary syndrome, hepatorenal syndrome, portal hypertension (esophageal and gastric varices), liver abscess (amebic liver abscess), liver cirrhosis (alcoholic, biliary and experimental), alcoholic liver diseases (fatty liver, hepatitis, cirrhosis), parasitic (hepatic echinococcosis, fascioliasis, amebic liver abscess), jaundice (hemolytic, hepatocellular, and cholestatic), cholestasis, portal hypertension, liver enlargement, ascites, hepatitis (alcoholic hepatitis, animal hepatitis, chronic hepatitis (autoimmune, hepatitis B, hepatitis C, hepatitis D, drug induced), toxic hepatitis, viral human hepatitis (hepatitis A, hepatitis B, hepatitis C, hepatitis D, hepatitis E), Wilson's disease, granulomatous hepatitis, secondary biliary cirrhosis, hepatic encephalopathy, portal hypertension, varices, hepatic encephalopathy, primary biliary cirrhosis, primary sclerosing cholangitis, hepatocellular adenoma, hemangiomas, bile stones, liver failure (hepatic encephalopathy, acute liver failure), and liver neoplasms (angiomyolipoma, calcified liver metastases, cystic liver metastases, epithelial tumors, fibrolamellar hepatocarcinoma, focal nodular hyperplasia, hepatic adenoma, hepatobiliary cystadenoma, hepatoblastoma, hepatocellular carcinoma, hepatoma, liver cancer, liver hemangioendothelioma, mesenchymal hamartoma, mesenchymal tumors of liver, nodular regenerative hyperplasia, benign liver tumors (Hepatic cysts [Simple cysts, Polycystic liver disease, Hepatobiliary cystadenoma, Choledochal cyst], Mesenchymal tumors [Mesenchymal hamartoma, Infantile hemangioendothelioma, Hemangioma, Peliosis hepatis, Lipomas, Inflammatory pseudotumor, Miscellaneous], Epithelial tumors [Bile duct epithelium (Bile duct hamartoma, Bile duct adenoma), Hepatocyte (Adenoma, Focal nodular hyperplasia, Nodular regenerative hyperplasia)]), malignant liver tumors [hepatocellular, hepatoblastoma, hepatocellular carcinoma, cholangiocellular, cholangiocarcinoma, cystadenocarcinoma, tumors of blood vessels, angiosarcoma, Kaposi's sarcoma, hemangioendothelioma, other tumors,

10091504, 030703



embryonal sarcoma, fibrosarcoma, leiomyosarcoma, rhabdomyosarcoma, carcinosarcoma, teratoma, carcinoid, squamous carcinoma, primary lymphoma]], peliosis hepatis, erythrohepatic porphyria, hepatic porphyria (acute intermittent porphyria, porphyria cutanea tarda), Zellweger syndrome).

[00696] Pancreatic diseases and/or disorders include acute pancreatitis, chronic pancreatitis (acute necrotizing pancreatitis, alcoholic pancreatitis), neoplasms (adenocarcinoma of the pancreas, cystadenocarcinoma, insulinoma, gastrinoma, and glucagonoma, cystic neoplasms, islet-cell tumors, pancreoblastoma), and other pancreatic diseases (e.g., cystic fibrosis, cyst (pancreatic pseudocyst, pancreatic fistula, insufficiency)).

[00697] Gallbladder diseases include gallstones (cholelithiasis and choledocholithiasis), postcholecystectomy syndrome, diverticulosis of the gallbladder, acute cholecystitis, chronic cholecystitis, bile duct tumors, and mucocele.

[00698] Diseases and/or disorders of the large intestine include antibiotic-associated colitis, diverticulitis, ulcerative colitis, acquired megacolon, abscesses, fungal and bacterial infections, anorectal disorders (e.g., fissures, hemorrhoids), colonic diseases (colitis, colonic neoplasms [colon cancer, adenomatous colon polyps (e.g., villous adenoma), colon carcinoma, colorectal cancer], colonic diverticulitis, colonic diverticulosis, megacolon [Hirschsprung disease, toxic megacolon]; sigmoid diseases [proctocolitis, sigmoid neoplasms]), constipation, Crohn's disease, diarrhea (infantile diarrhea, dysentery), duodenal diseases (duodenal neoplasms, duodenal obstruction, duodenal ulcer, duodenitis), enteritis (enterocolitis), HIV enteropathy, ileal diseases (ileal neoplasms, ileitis), immunoproliferative small intestinal disease, inflammatory bowel disease (ulcerative colitis, Crohn's disease), intestinal atresia, parasitic diseases (anisakiasis, balantidiasis, blastocystis infections, cryptosporidiosis, dientamoebiasis, amebic dysentery, giardiasis), intestinal fistula (rectal fistula), intestinal neoplasms (cecal neoplasms, colonic neoplasms, duodenal neoplasms, ileal neoplasms, intestinal polyps, jejunal neoplasms, rectal neoplasms), intestinal obstruction (afferent loop syndrome, duodenal obstruction, impacted feces, intestinal pseudo-obstruction [cecal volvulus], intussusception), intestinal perforation, intestinal polyps (colonic polyps, gardner syndrome, peutz-jeghers syndrome), jejunal diseases (jejunal neoplasms), malabsorption syndromes (blind loop syndrome, celiac disease, lactose intolerance, short bowl syndrome, tropical

10091504 030702  
20X050 405T600T

sprue, whipple's disease), mesenteric vascular occlusion, pneumatosis cystoides intestinalis, protein-losing enteropathies (intestinal lymphangiectasis), rectal diseases (anus diseases, fecal incontinence, hemorrhoids, proctitis, rectal fistula, rectal prolapse, rectocele), peptic ulcer (duodenal ulcer, peptic esophagitis, hemorrhage, perforation, stomach ulcer, Zollinger-Ellison syndrome), postgastrectomy syndromes (dumping syndrome), stomach diseases (e.g., achlorhydria, duodenogastric reflux (bile reflux), gastric antral vascular ectasia, gastric fistula, gastric outlet obstruction, gastritis (atrophic or hypertrophic), gastroparesis, stomach dilatation, stomach diverticulum, stomach neoplasms (gastric cancer, gastric polyps, gastric adenocarcinoma, hyperplastic gastric polyp), stomach rupture, stomach ulcer, stomach volvulus), tuberculosis, visceroptosis, vomiting (e.g., hematemesis, hyperemesis gravidarum, postoperative nausea and vomiting) and hemorrhagic colitis.

- [00699] Further diseases and/or disorders of the gastrointestinal system include biliary tract diseases, such as, gastroschisis, fistula (e.g., biliary fistula, esophageal fistula, gastric fistula, intestinal fistula, pancreatic fistula), neoplasms (e.g., biliary tract neoplasms, esophageal neoplasms, such as adenocarcinoma of the esophagus, esophageal squamous cell carcinoma, gastrointestinal neoplasms, pancreatic neoplasms, such as adenocarcinoma of the pancreas, mucinous cystic neoplasm of the pancreas, pancreatic cystic neoplasms, pancreatoblastoma, and peritoneal neoplasms), esophageal disease (e.g., bullous diseases, candidiasis, glycogenic acanthosis, ulceration, barrett esophagus varices, atresia, cyst, diverticulum (e.g., Zenker's diverticulum), fistula (e.g., tracheoesophageal fistula), motility disorders (e.g., CREST syndrome, deglutition disorders, achalasia, spasm, gastroesophageal reflux), neoplasms, perforation (e.g., Boerhaave syndrome, Mallory-Weiss syndrome), stenosis, esophagitis, diaphragmatic hernia (e.g., hiatal hernia); gastrointestinal diseases, such as, gastroenteritis (e.g., cholera morbus, norwalk virus infection), hemorrhage (e.g., hematemesis, melena, peptic ulcer hemorrhage), stomach neoplasms (gastric cancer, gastric polyps, gastric adenocarcinoma, stomach cancer)), hernia (e.g., congenital diaphragmatic hernia, femoral hernia, inguinal hernia, obturator hernia, umbilical hernia, ventral hernia), and intestinal diseases (e.g., cecal diseases (appendicitis, cecal neoplasms)).

## Reproductive System Disorders

- [00700] The polynucleotides or polypeptides, or agonists or antagonists of the invention may be used for the diagnosis, treatment, or prevention of diseases and/or disorders of the reproductive system. Reproductive system disorders that can be treated by the compositions of the invention, include, but are not limited to, reproductive system injuries, infections, neoplastic disorders, congenital defects, and diseases or disorders which result in infertility, complications with pregnancy, labor, or parturition, and postpartum difficulties.
- [00701] Reproductive system disorders and/or diseases include diseases and/or disorders of the testes, including, but not limited to, testicular atrophy, testicular feminization, cryptorchism (unilateral and bilateral), anorchia, ectopic testis, epididymitis and orchitis (typically resulting from infections such as, for example, gonorrhea, mumps, tuberculosis, and syphilis), testicular torsion, vasitis nodosa, germ cell tumors (e.g., seminomas, embryonal cell carcinomas, teratocarcinomas, choriocarcinomas, yolk sac tumors, and teratomas), stromal tumors (e.g., Leydig cell tumors), hydrocele, hematocele, varicocele, spermatocele, inguinal hernia, and disorders of sperm production (e.g., immotile cilia syndrome, aspermia, asthenozoospermia, azoospermia, oligospermia, and teratozoospermia).
- [00702] Reproductive system disorders also include, but are not limited to, disorders of the prostate gland, such as acute non-bacterial prostatitis, chronic non-bacterial prostatitis, acute bacterial prostatitis, chronic bacterial prostatitis, prostatodystonia, prostatosis, granulomatous prostatitis, malacoplakia, benign prostatic hypertrophy or hyperplasia, and prostate neoplastic disorders, including adenocarcinomas, transitional cell carcinomas, ductal carcinomas, and squamous cell carcinomas.
- [00703] Additionally, the compositions of the invention may be useful in the diagnosis, treatment, and/or prevention of disorders or diseases of the penis and urethra, including, but not limited to, inflammatory disorders, such as balanoposthitis, balanitis xerotica obliterans, phimosis, paraphimosis, syphilis, herpes simplex virus, gonorrhea, non-gonococcal urethritis, chlamydia, mycoplasma, trichomonas, HIV, AIDS, Reiter's syndrome, condyloma acuminatum, condyloma latum, and pearly penile papules; urethral abnormalities, such as hypospadias, epispadias, and phimosis; premalignant lesions, including Erythroplasia of Queyrat, Bowen's disease, Bowenoid paplosis, giant condyloma of Buscke-Lowenstein, and verrucous

carcinoma; penile cancers, including squamous cell carcinomas, carcinoma in situ, verrucous carcinoma, and disseminated penile carcinoma; urethral neoplastic disorders, including penile urethral carcinoma, bulbomembranous urethral carcinoma, and prostatic urethral carcinoma; and erectile disorders, such as priapism, Peyronie's disease, erectile dysfunction, and impotence.

[00704] Moreover, diseases and/or disorders of the vas deferens include, but are not limited to, vasculitis and CBAVD (congenital bilateral absence of the vas deferens); additionally, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases and/or disorders of the seminal vesicles, including but not limited to, hydatid disease, congenital chloride diarrhea, and polycystic kidney disease.

[00705] Other disorders and/or diseases of the male reproductive system that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, Klinefelter's syndrome, Young's syndrome, premature ejaculation, diabetes mellitus, cystic fibrosis, Kartagener's syndrome, high fever, multiple sclerosis, and gynecomastia.

[00706] Further, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases and/or disorders of the vagina and vulva, including, but not limited to, bacterial vaginosis, candida vaginitis, herpes simplex virus, chancroid, granuloma inguinale, lymphogranuloma venereum, scabies, human papillomavirus, vaginal trauma, vulvar trauma, adenositis, chlamydia vaginitis, gonorrhea, trichomonas vaginitis, condyloma acuminatum, syphilis, molluscum contagiosum, atrophic vaginitis, Paget's disease, lichen sclerosus, lichen planus, vulvodynia, toxic shock syndrome, vaginismus, vulvovaginitis, vulvar vestibulitis, and neoplastic disorders, such as squamous cell hyperplasia, clear cell carcinoma, basal cell carcinoma, melanomas, cancer of Bartholin's gland, and vulvar intraepithelial neoplasia.

[00707] Disorders and/or diseases of the uterus that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, dysmenorrhea, retroverted uterus, endometriosis, fibroids, adenomyosis, anovulatory bleeding, amenorrhea, Cushing's syndrome, hydatidiform moles, Asherman's syndrome, premature menopause, precocious puberty, uterine polyps,

dysfunctional uterine bleeding (e.g., due to aberrant hormonal signals), and neoplastic disorders, such as adenocarcinomas, leiomyosarcomas, and sarcomas. Additionally, the polypeptides, polynucleotides, or agonists or antagonists of the invention may be useful as a marker or detector of, as well as in the diagnosis, treatment, and/or prevention of congenital uterine abnormalities, such as bicornuate uterus, septate uterus, simple unicornuate uterus, unicornuate uterus with a noncavitary rudimentary horn, unicornuate uterus with a non-communicating cavitary rudimentary horn, unicornuate uterus with a communicating cavitary horn, arcuate uterus, uterine didelphys, and T-shaped uterus.

[00708] Ovarian diseases and/or disorders that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, anovulation, polycystic ovary syndrome (Stein-Leventhal syndrome), ovarian cysts, ovarian hypofunction, ovarian insensitivity to gonadotropins, ovarian overproduction of androgens, right ovarian vein syndrome, amenorrhea, hirsutism, and ovarian cancer (including, but not limited to, primary and secondary cancerous growth, Sertoli-Leydig tumors, endometrioid carcinoma of the ovary, ovarian papillary serous adenocarcinoma, ovarian mucinous adenocarcinoma, and Ovarian Krukenberg tumors).

[00709] Cervical diseases and/or disorders that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, cervicitis, chronic cervicitis, mucopurulent cervicitis, cervical dysplasia, cervical polyps, Nabothian cysts, cervical erosion, cervical incompetence, and cervical neoplasms (including, for example, cervical carcinoma, squamous metaplasia, squamous cell carcinoma, adenosquamous cell neoplasia, and columnar cell neoplasia).

[00710] Additionally, diseases and/or disorders of the reproductive system that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, disorders and/or diseases of pregnancy, including miscarriage and stillbirth, such as early abortion, late abortion, spontaneous abortion, induced abortion, therapeutic abortion, threatened abortion, missed abortion, incomplete abortion, complete abortion, habitual abortion, missed abortion, and septic abortion; ectopic pregnancy, anemia, Rh incompatibility, vaginal bleeding during pregnancy, gestational diabetes, intrauterine growth retardation, polyhydramnios, HELLP

syndrome, abruptio placentae, placenta previa, hyperemesis, preeclampsia, eclampsia, herpes gestationis, and urticaria of pregnancy. Additionally, the polynucleotides, polypeptides, and agonists or antagonists of the present invention may be used in the diagnosis, treatment, and/or prevention of diseases that can complicate pregnancy, including heart disease, heart failure, rheumatic heart disease, congenital heart disease, mitral valve prolapse, high blood pressure, anemia, kidney disease, infectious disease (e.g., rubella, cytomegalovirus, toxoplasmosis, infectious hepatitis, chlamydia, HIV, AIDS, and genital herpes), diabetes mellitus, Graves' disease, thyroiditis, hypothyroidism, Hashimoto's thyroiditis, chronic active hepatitis, cirrhosis of the liver, primary biliary cirrhosis, asthma, systemic lupus erythematosus, rheumatoid arthritis, myasthenia gravis, idiopathic thrombocytopenic purpura, appendicitis, ovarian cysts, gallbladder disorders, and obstruction of the intestine.

- [00711] Complications associated with labor and parturition that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, premature rupture of the membranes, pre-term labor, post-term pregnancy, postmaturity, labor that progresses too slowly, fetal distress (e.g., abnormal heart rate (fetal or maternal), breathing problems, and abnormal fetal position), shoulder dystocia, prolapsed umbilical cord, amniotic fluid embolism, and aberrant uterine bleeding.
- [00712] Further, diseases and/or disorders of the postdelivery period, that may be diagnosed, treated, and/or prevented with the compositions of the invention, include, but are not limited to, endometritis, myometritis, parametritis, peritonitis, pelvic thrombophlebitis, pulmonary embolism, endotoxemia, pyelonephritis, saphenous thrombophlebitis, mastitis, cystitis, postpartum hemorrhage, and inverted uterus.
- [00713] Other disorders and/or diseases of the female reproductive system that may be diagnosed, treated, and/or prevented by the polynucleotides, polypeptides, and agonists or antagonists of the present invention include, but are not limited to, Turner's syndrome, pseudohermaphroditism, premenstrual syndrome, pelvic inflammatory disease, pelvic congestion (vascular engorgement), frigidity, anorgasmia, dyspareunia, ruptured fallopian tube, and Mittelschmerz.

#### Developmental and Inherited Disorders

20091504-030702

[00714] Polynuceotides or polypeptides, or agonists or antagonists of the present invention may be used to treat, prevent, diagnose, and/or prognose diseases associated with mixed fetal tissues, including, but not limited to, developmental and inherited disorders or defects of the nervous system, musculoskeletal system, excretory system, cardiovascular system, hematopoietic system, gastrointestinal system, reproductive system, and respiratory system. Compositions of the present invention may also be used to treat, prevent, diagnose, and/or prognose developmental and inherited disorders or defects associated with, but not limited to, skin, hair, visual, and auditory tissues, metabolism. Additionally, the compositions of the invention may be useful in the diagnosis, treatment, and/or prevention of disorders or diseases associated with, but not limited to, chromosomal or genetic abnormalities and hyperproliferation or neoplasia.

[00715] Disorders or defects of the nervous system associated with developmental or inherited abnormalities that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, adrenoleukodystrophy, agenesis of corpus callosum, Alexander disease, anencephaly, Angelman syndrome, Arnold-Chiari deformity, Batten disease, Canavan disease, cephalic disorders, Charcot-Marie-Tooth disease, encephalocele, Friedreich's ataxia, Gaucher's disease, Gorlin syndrome, Hallervorden-Spatz disease, hereditary spastic paraplegia, Huntington disease, hydranencephaly, hydrocephalus, Joubert syndrome, Lesch-Nyhan syndrome, leukodystrophy, Menkes disease, microcephaly, Niemann-Pick Type C1, neurofibromatosis, porencephaly, progeria, proteus syndrome, Refsum disease, spina bifida, Sturge-Weber syndrome, Tay-Sachs disease, tuberous sclerosis, and von Hippel-Lindau disease.

[00716] Developmental and inherited disorders resulting in disorders or defects of the musculoskeletal system that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, achondroplasia, atlanto-occipital fusion, arthrogryposis multiplex congenita, autosomal recessive muscular dystrophy, Becker's muscular dystrophy, cerebral palsy, choanal atresia, cleft lip, cleft palate, clubfoot, congenital amputation, congenital dislocation of the hip, congenital torticollis, congenital scoliosis, dopa-responsive dystonia, Duchenne muscular dystrophy, early-onset generalized dystonia, femoral torsion, Gorlin syndrome, hypophosphatasia, Klippel-Feil syndrome, knee dislocation, myoclonic

dystonia, myotonic dystrophy, nail-patella syndrome, osteogenesis imperfecta, paroxysmal dystonia, progeria, prune-belly syndrome, rapid-onset dystonia parkinsonism, scoliosis, syndactyly, Treacher Collins' syndrome, velocardiofacial syndrome, and X-linked dystonia-parkinsonism.

[00717] Developmental or hereditary disorders or defects of the excretory system that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, Alport's syndrome, Bartter's syndrome, bladder diverticula, bladder exstrophy, cystinuria, epispadias, Fanconi's syndrome, Hartnup disease, horseshoe kidney, hypospadias, kidney agenesis, kidney ectopia, kidney malrotation, Liddle's syndrome, medullary cystic disease, medullary sponge, multicystic kidney, kidney polycystic kidney disease, nail-patella syndrome, Potter's syndrome, urinary tract flow obstruction, vitamin D-resistant rickets, and Wilm's tumor.

[00718] Cardiovascular disorders or defects of developmental or hereditary origin that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, aortic valve stenosis, atrial septal defects, artioventricular (A-V) canal defect, bicuspid aortic valve, coarctation of the aorta, dextrocardia, Ebstein's anomaly, Eisenmenger's complex, hypoplastic left heart syndrome, Marfan syndrome, patent ductus arteriosus, progeria, pulmonary atresia, pulmonary valve stenosis, subaortic stenosis, tetralogy of fallot, total anomalous pulmonary venous (P-V) connection, transposition of the great arteries, tricuspid atresia, truncus arteriosus, ventricular septal defects. Developmental or inherited disorders resulting in disorders involving the hematopoietic system that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but not limited to, Bernard-Soulier syndrome, Chédiak-Higashi syndrome, hemophilia, Hermansky-Pudlak syndrome, sickle cell anemia, storage pool disease, thromboxane A2 dysfunction, thrombasthenia, and von Willebrand's disease.

[00719] The compositions of the invention may also be used to diagnose, treat, and/or prevent developmental and inherited disorders resulting in disorders or defects of the gastrointestinal system, including, but not limited to, anal atresia, biliary atresia, esophageal atresia, diaphragmatic hernia, Hirschsprung's disease, Meckel's diverticulum, oligohydramnios, omphalocele, polyhydramnios, porphyria, situs inversus viscera. Developmental or inherited disorders resulting in metabolic



disorders that may be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, alpha-1 antitrypsin deficiency, cystic fibrosis, hemochromatosis, lysosomal storage disease, phenylketonuria, Wilson's disease, and Zellweger syndrome.

[00720] Disorders of the reproductive system that are developmentally or hereditary related that may also be diagnosed, treated, and/or prevented with the compositions of the invention include, but are not limited to, androgen insensitivity syndrome, ambiguous genitalia, autosomal sex reversal, congenital adrenal hyperplasia, gonadoblastoma, ovarian germ cell cancer, pseudohermaphroditism, true hermaphroditism, undescended testis, XX male syndrome, and XY female type gonadal dysgenesis. The compositions of the invention may also be used to diagnose, treat, and/or prevent developmental or inherited respiratory defects including, but not limited to, askin tumor, azygos lobe, congenital diaphragmatic hernia, congenital lobar emphysema, cystic adenomatoid malformation, lobar emphysema, hyaline membrane disease, and pectus excavatum.

[00721] Developmental or inherited disorders may also result from chromosomal or genetic aberration that may be diagnosed, treated, and/or prevented with the compositions of the invention including, but not limited to, 4p- syndrome, cri du chat syndrome, Digeorge syndrome, Down's syndrome, Edward's syndrome, fragile X syndrome, Klinefelter's syndrome, Patau's syndrome, Prader-Willi syndrome, progeria, Turner's syndrome, triple X syndrome, and XYY syndrome. Other developmental disorders that can be diagnosed, treated, and/or prevented with the compositions of the invention, include, but are not limited to, fetal alcohol syndrome, and can be caused by environmental factors surrounding the developing fetus.

[00722] The compositions of the invention may further be able to be used to diagnose, treat, and/or prevent errors in development or a genetic disposition that may result in hyperproliferative disorders or neoplasms, including, but not limited to, acute childhood lymphoblastic leukemia, askin tumor, Beckwith-Wiedemann syndrome, childhood acute myeloid leukemia, childhood brain stem glioma, childhood cerebellar astrocytoma, childhood extracranial germ cell tumors childhood (primary), gonadoblastoma, hepatocellular cancer, childhood Hodgkin's disease, childhood Hodgkin's lymphoma, childhood hypothalamic and visual pathway

glioma, childhood (primary) liver cancer, childhood lymphoblastic leukemia, childhood medulloblastoma, childhood non-Hodgkin's lymphoma, childhood pineal and supratentorial primitive neuroectodermal tumors, childhood primary liver cancer, childhood rhabdomyosarcoma, childhood soft tissue sarcoma, Gorlin syndrome, familial multiple endocrine neoplasia type I, neuroblastoma, ovarian germ cell cancer, pheochromocytoma, retinoblastoma, and Wilm's tumor.

[00723] Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppositorial solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

#### Diseases at the Cellular Level

[00724] Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated, prevented, diagnosed and/or prognosed using polynucleotides or polypeptides, as well as antagonists or agonists of the present invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection.

[00725] In preferred embodiments, polynucleotides, polypeptides, and/or antagonists of the invention are used to inhibit growth, progression, and/or metastasis of cancers, in particular those [listed above] involving cardiovascular system tissues.

[00726] Additional diseases or conditions associated with increased cell survival that could be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

[00727] Diseases associated with increased apoptosis that could be treated, prevented, diagnosed, and/or prognosed using polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, include, but are not limited to, AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders

(such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestasis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

### **Wound Healing and Epithelial Cell Proliferation**

[00728] In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote dermal reestablishment subsequent to dermal loss.

[00729] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are types of grafts that polynucleotides or polypeptides, agonists or antagonists of the present invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic graft, autoepdermic grafts, avacular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft,

epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft, mesh graft, mucosal graft, Ollier-Thiersch graft, omentopal graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, can be used to promote skin strength and to improve the appearance of aged skin.

[00730] It is believed that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes, type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. Polynucleotides or polypeptides, agonists or antagonists of the present invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

[00731] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may have a cytoprotective effect on the small intestine mucosa. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may also stimulate healing of mucositis (mouth ulcers) that result from chemotherapy and viral infections.

[00732] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could also

be used to treat gastric and duodenal ulcers and help heal by scar formation of the mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases, which result in destruction of the mucosal surface of the small or large intestine, respectively. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with polynucleotides or polypeptides, agonists or antagonists of the present invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat diseases associated with the under expression.

[00733] Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to prevent and heal damage to the lungs due to various pathological states. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of alveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using polynucleotides or polypeptides, agonists or antagonists of the present invention. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to stimulate the proliferation and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

[00734] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis

and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

[00735] In addition, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent manifestation of the disease. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

#### **Infectious Disease**

[00736] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

[00737] Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papilloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g.,

Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiollitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever, yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat AIDS.

[00738] Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Cryptococcus neoformans, Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia (e.g., *Borrelia burgdorferi*, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, *E. coli* (e.g., Enterotoxigenic *E. coli* and Enterohemorrhagic *E. coli*), Enterobacteriaceae (Klebsiella, Salmonella (e.g., *Salmonella typhi*, and *Salmonella paratyphi*), Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, *Mycobacterium leprae*, *Vibrio cholerae*, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), *Meisseria meningitidis*, Pasteurellacea Infections (e.g., Actinobacillus, Heamophilus (e.g., Heamophilus influenza type B), Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Treponema spp., Leptospira spp.,



Shigella spp., Staphylococcal, Meningiococcal, Pneumococcal and Streptococcal (e.g., Streptococcus pneumoniae and Group B Streptococcus). These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid, pneumonia, Gonorrhea, meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections. Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, Polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat: tetanus, Diphtheria, botulism, and/or meningitis type B.

[00739] Moreover, parasitic agents causing disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax, Plasmodium falciparum, Plasmodium malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases.

[00740] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient,

supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

### **Regeneration**

[00741] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns, incisions, or ulcers), age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

[00742] Tissues that could be regenerated using the present invention include organs (e.g., pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac), vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone, cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased scarring. Regeneration also may include angiogenesis.

[00743] Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may increase regeneration of tissues difficult to heal. For example, increased tendon/ligament regeneration would quicken recovery time after damage. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention could also be used prophylactically in an effort to avoid damage. Specific diseases that could be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. A further example of tissue regeneration of non-healing wounds includes pressure ulcers, ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

[00744] Similarly, nerve and brain tissue could also be regenerated by using polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, to proliferate and differentiate nerve cells. Diseases that could be treated using this method include central and peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders (e.g., spinal cord disorders,

head trauma, cerebrovascular disease, and stroke). Specifically, diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from chemotherapy or other medical therapies), localized neuropathies, and central nervous system diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotides or polypeptides, as well as agonists or antagonists of the present invention.

### **Chemotaxis**

[00745] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells (e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells) to a particular site in the body, such as inflammation, infection, or site of hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

[00746] Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

[00747] It is also contemplated that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention could be used as an inhibitor of chemotaxis.

### **Binding Activity**

[00748] A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit

(antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

[00749] Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

[00750] Preferably, the screening for these molecules involves producing appropriate cells, which express the polypeptide. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

[00751] The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

[00752] Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

[00753] Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

[00754] Additionally, the receptor to which the polypeptide of the present invention binds can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in

Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labeled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

[00755] Following fixation and incubation, the slides are subjected to auto-radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

[00756] As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

[00757] Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of the polypeptide of the present invention thereby effectively generating agonists and antagonists of the polypeptide of the present invention. See generally, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., et al., Curr. Opinion Biotechnol. 8:724-33 (1997); Harayama, S. Trends Biotechnol. 16(2):76-82 (1998); Hansson L. O., et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. Biotechniques 24(2):308-13 (1998); each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into

10091504-030702

a desired molecule by homologous, or site-specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptide of the present invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF), TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

[00758] Other preferred fragments are biologically active fragments of the polypeptide of the present invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

[00759] Additionally, this invention provides a method of screening compounds to identify those, which modulate the action of the polypeptide of the present invention. An example of such an assay comprises combining a mammalian fibroblast cell, the polypeptide of the present invention, the compound to be screened and  $^3\text{[H]}$  thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound stimulates proliferation by determining the uptake of  $^3\text{[H]}$  thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography, which measures the incorporation of  $^3\text{[H]}$  thymidine. Both agonist and antagonist compounds may be identified by this procedure.

10091504, 030702  
2020ED, 405T600T

[00760] In another method, a mammalian cell or membrane preparation expressing a receptor for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

[00761] All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from suitably manipulated cells or tissues.

[00762] Therefore, the invention includes a method of identifying compounds, which bind to a polypeptide of the invention comprising the steps of: (a) incubating a candidate binding compound with a polypeptide of the present invention; and (b) determining if binding has occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with a polypeptide of the present invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

### Targeted Delivery

[00763] In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

[00764] As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of

compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

[00765] In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

[00766] By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxyacetamide derivatives of doxorubicin.

### Drug Screening

[00767] Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules, which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected



compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides following binding.

[00768] This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

[00769] Thus, the present invention provides methods of screening for drugs or any other agents, which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the present invention.

[00770] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

[00771] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide, which shares one or more antigenic epitopes with a polypeptide of the invention.

#### Antisense And Ribozyme (Antagonists)

[00772] In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to cDNA sequences contained in cDNA Clone ID NO:Z identified for example, in Table 1A. In one embodiment, antisense sequence is generated internally, by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, J., Neurochem. 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, J., Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

[00773] For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed *in vitro* by incubating cells with the oligoribonucleotide. A similar procedure for *in vivo* use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl<sub>2</sub>, 10mM

dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoR1/Hind III site of the retroviral vector PMV7 (WO 91/15580).

[00774] For example, the 5' coding portion of a polynucleotide that encodes the polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA *in vivo* and blocks translation of the mRNA molecule into receptor polypeptide.

[00775] In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding the polypeptide of the present invention or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, Nature 29:304-310 (1981), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., Cell 22:787-797 (1980), the herpes thymidine promoter (Wagner et al., Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster, et al., Nature 296:39-42 (1982)), etc.

[00776] The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of the present invention. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation

may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

[00777] Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335. Thus, oligonucleotides complementary to either the 5'- or 3'- non- translated, non-coding regions of polynucleotide sequences described herein could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5'-, 3'- or coding region of mRNA of the present invention, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

[00778] The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-

triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

[00779] The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylecytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

[00780] The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

[00781] In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

[00782] In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl.

Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

[00783] Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

[00784] While antisense nucleotides complementary to the coding region sequence could be used, those complementary to the transcribed untranslated region are most preferred.

[00785] Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within the nucleotide sequence of SEQ ID NO:X. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

[00786] As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express *in vivo*. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that

transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

[00787] Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

[00788] The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirable in cases such as restenosis after balloon angioplasty.

[00789] The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

[00790] The antagonist/agonist may also be employed to treat the diseases described herein.

[00791] Thus, the invention provides a method of treating disorders or diseases, including but not limited to the disorders or diseases listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a ribozyme directed to the polynucleotide of the present invention.

#### **Binding Peptides and Other Molecules**

[00792] The invention also encompasses screening methods for identifying polypeptides and nonpolypeptides that bind cardiovascular system antigen polypeptides, and the cardiovascular system antigen binding molecules identified thereby. These binding molecules are useful, for example, as agonists and antagonists of the cardiovascular system antigen polypeptides. Such agonists and antagonists can be used, in accordance with the invention, in the therapeutic embodiments described in detail, below.

[00793] This method comprises the steps of:

contacting cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides with a plurality of molecules; and

identifying a molecule that binds the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides.

[00794] The step of contacting the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides with the plurality of molecules may be effected in a number of ways. For example, one may contemplate immobilizing the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides on a solid support and bringing a solution of the plurality of molecules in contact with the immobilized cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides. Such a procedure would be akin to an affinity chromatographic process, with the affinity matrix being comprised of the immobilized cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides. The molecules having a selective affinity for the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides can then be purified by affinity selection. The nature of the solid support, process for attachment of the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides to the solid support, solvent, and conditions of the affinity isolation or selection are largely conventional and well known to those of ordinary skill in the art.

[00795] Alternatively, one may also separate a plurality of polypeptides into substantially separate fractions comprising a subset of or individual polypeptides. For instance, one can separate the plurality of polypeptides by gel electrophoresis, column chromatography, or like method known to those of ordinary skill for the separation of polypeptides. The individual polypeptides can also be produced by a transformed host cell in such a way as to be expressed on or about its outer surface (e.g., a recombinant phage). Individual isolates can then be "probed" by the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides, optionally in the presence of an inducer should one be required for expression, to determine if any selective affinity interaction takes place between the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides and the individual clone. Prior to contacting the cardiovascular system



antigen polypeptides or cardiovascular system antigen-like polypeptides with each fraction comprising individual polypeptides, the polypeptides could first be transferred to a solid support for additional convenience. Such a solid support may simply be a piece of filter membrane, such as one made of nitrocellulose or nylon. In this manner, positive clones could be identified from a collection of transformed host cells of an expression library, which harbor a DNA construct encoding a polypeptide having a selective affinity for cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides. Furthermore, the amino acid sequence of the polypeptide having a selective affinity for the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides can be determined directly by conventional means or the coding sequence of the DNA encoding the polypeptide can frequently be determined more conveniently. The primary sequence can then be deduced from the corresponding DNA sequence. If the amino acid sequence is to be determined from the polypeptide itself, one may use microsequencing techniques. The sequencing technique may include mass spectroscopy.

[00796] In certain situations, it may be desirable to wash away any unbound cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides, or alternatively, unbound polypeptides, from a mixture of the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides and the plurality of polypeptides prior to attempting to determine or to detect the presence of a selective affinity interaction. Such a wash step may be particularly desirable when the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides or the plurality of polypeptides is bound to a solid support.

[00797] The plurality of molecules provided according to this method may be provided by way of diversity libraries, such as random or combinatorial peptide or nonpeptide libraries which can be screened for molecules that specifically bind cardiovascular system antigen polypeptides. Many libraries are known in the art that can be used, e.g., chemically synthesized libraries, recombinant (e.g., phage display libraries), and *in vitro* translation-based libraries. Examples of chemically synthesized libraries are described in Fodor et al., 1991, Science 251:767-773; Houghten et al., 1991, Nature 354:84-86; Lam et al., 1991, Nature 354:82-84; Medynski, 1994,

Bio/Technology 12:709-710; Gallop et al., 1994, J. Medicinal Chemistry 37(9):1233-1251; Ohlmeyer et al., 1993, Proc. Natl. Acad. Sci. USA 90:10922-10926; Erb et al., 1994, Proc. Natl. Acad. Sci. USA 91:11422-11426; Houghten et al., 1992, Biotechniques 13:412; Jayawickreme et al., 1994, Proc. Natl. Acad. Sci. USA 91:1614-1618; Salmon et al., 1993, Proc. Natl. Acad. Sci. USA 90:11708-11712; PCT Publication No. WO 93/20242; and Brenner and Lerner, 1992, Proc. Natl. Acad. Sci. USA 89:5381-5383.

[00798] Examples of phage display libraries are described in Scott and Smith, 1990, Science 249:386-390; Devlin et al., 1990, Science, 249:404-406; Christian, R. B., et al., 1992, J. Mol. Biol. 227:711-718; Lenstra, 1992, J. Immunol. Meth. 152:149-157; Kay et al., 1993, Gene 128:59-65; and PCT Publication No. WO 94/18318 dated Aug. 18, 1994.

[00799] *In vitro* translation-based libraries include but are not limited to those described in PCT Publication No. WO 91/05058 dated Apr. 18, 1991; and Mattheakis et al., 1994, Proc. Natl. Acad. Sci. USA 91:9022-9026.

[00800] By way of examples of nonpeptide libraries, a benzodiazepine library (see e.g., Bunin et al., 1994, Proc. Natl. Acad. Sci. USA 91:4708-4712) can be adapted for use. Peptoid libraries (Simon et al., 1992, Proc. Natl. Acad. Sci. USA 89:9367-9371) can also be used. Another example of a library that can be used, in which the amide functionalities in peptides have been permethylated to generate a chemically transformed combinatorial library, is described by Ostresh et al. (1994, Proc. Natl. Acad. Sci. USA 91:11138-11142).

[00801] The variety of non-peptide libraries that are useful in the present invention is great. For example, Ecker and Crooke, 1995, Bio/Technology 13:351-360 list benzodiazepines, hydantoins, piperazinediones, biphenyls, sugar analogs, beta-mercaptoketones, arylacetic acids, acylpiperidines, benzopyrans, cubanes, xanthines, aminimides, and oxazolones as among the chemical species that form the basis of various libraries.

[00802] Non-peptide libraries can be classified broadly into two types: decorated monomers and oligomers. Decorated monomer libraries employ a relatively simple scaffold structure upon which a variety functional groups is added. Often the scaffold will be a molecule with a known useful pharmacological activity. For example, the scaffold might be the benzodiazepine structure.

[00803] Non-peptide oligomer libraries utilize a large number of monomers that are assembled together in ways that create new shapes that depend on the order of the monomers. Among the monomer units that have been used are carbamates, pyrrolinones, and morpholinos. Peptoids, peptide-like oligomers in which the side chain is attached to the alpha amino group rather than the alpha carbon, form the basis of another version of non-peptide oligomer libraries. The first non-peptide oligomer libraries utilized a single type of monomer and thus contained a repeating backbone. Recent libraries have utilized more than one monomer, giving the libraries added flexibility.

[00804] Screening the libraries can be accomplished by any of a variety of commonly known methods. See, e.g., the following references, which disclose screening of peptide libraries: Parmley and Smith, 1989, Adv. Exp. Med. Biol. 251:215-218; Scott and Smith, 1990, Science 249:386-390; Fowlkes et al., 1992, BioTechniques 13:422-427; Oldenburg et al., 1992, Proc. Natl. Acad. Sci. USA 89:5393-5397; Yu et al., 1994, Cell 76:933-945; Staudt et al., 1988, Science 241:577-580; Bock et al., 1992, Nature 355:564-566; Tuerk et al., 1992, Proc. Natl. Acad. Sci. USA 89:6988-6992; Ellington et al., 1992, Nature 355:850-852; U.S. Pat. No. 5,096,815, U.S. Pat. No. 5,223,409, and U.S. Pat. No. 5,198,346, all to Ladner et al.; Rebar and Pabo, 1993, Science 263:671-673; and CT Publication No. WO 94/18318.

[00805] In a specific embodiment, screening to identify a molecule that binds cardiovascular system antigen polypeptides can be carried out by contacting the library members with a cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides immobilized on a solid phase and harvesting those library members that bind to the cardiovascular system antigen polypeptides or cardiovascular system antigen-like polypeptides. Examples of such screening methods, termed "panning" techniques are described by way of example in Parmley and Smith, 1988, Gene 73:305-318; Fowlkes et al., 1992, BioTechniques 13:422-427; International Publication No. WO 94/18318; and in references cited herein.

[00806] In another embodiment, the two-hybrid system for selecting interacting proteins in yeast (Fields and Song, 1989, Nature 340:245-246; Chien et al., 1991, Proc. Natl. Acad. Sci. USA 88:9578-9582) can be used to identify molecules that specifically bind to cardiovascular system antigen polypeptides or cardiovascular system

antigen-like polypeptides.

[00807] Where the cardiovascular system antigen binding molecule is a polypeptide, the polypeptide can be conveniently selected from any peptide library, including random peptide libraries, combinatorial peptide libraries, or biased peptide libraries. The term "biased" is used herein to mean that the method of generating the library is manipulated so as to restrict one or more parameters that govern the diversity of the resulting collection of molecules, in this case peptides.

[00808] Thus, a truly random peptide library would generate a collection of peptides in which the probability of finding a particular amino acid at a given position of the peptide is the same for all 20 amino acids. A bias can be introduced into the library, however, by specifying, for example, that a lysine occur every fifth amino acid or that positions 4, 8, and 9 of a decapeptide library be fixed to include only arginine. Clearly, many types of biases can be contemplated, and the present invention is not restricted to any particular bias. Furthermore, the present invention contemplates specific types of peptide libraries, such as phage displayed peptide libraries and those that utilize a DNA construct comprising a lambda phage vector with a DNA insert.

[00809] As mentioned above, in the case of a cardiovascular system antigen binding molecule that is a polypeptide, the polypeptide may have about 6 to less than about 60 amino acid residues, preferably about 6 to about 10 amino acid residues, and most preferably, about 6 to about 22 amino acids. In another embodiment, a cardiovascular system antigen binding polypeptide has in the range of 15-100 amino acids, or 20-50 amino acids.

[00810] The selected cardiovascular system antigen binding polypeptide can be obtained by chemical synthesis or recombinant expression.

#### Other Activities

[00811] A polypeptide, polynucleotide, agonist, or antagonist of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. The polypeptide, polynucleotide, agonist, or antagonist of the present

invention may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

[00812] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

[00813] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. A polypeptide, polynucleotide, agonist, or antagonist of the present invention may have the ability to stimulate chondrocyte growth; therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

[00814] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

[00815] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, a polypeptide, polynucleotide, agonist, or antagonist of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

[00816] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues. A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

[00817] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, as discussed above, hematopoietic lineage.

10091504, 030702

[00818] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used to modulate mammalian characteristics, such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, a polypeptide, polynucleotide, agonist, or antagonist of the present invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

[00819] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, cardiac rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity), hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

[00820] A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

[00821] The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most preferred embodiments, the host is a human.

#### **Other Preferred Embodiments**

[00822] Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[00823] Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of the portion of SEQ ID NO:X as defined in column 4, "ORF (From-To)", in Table 1A.

[00824] Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of the portion of SEQ ID NO:X as defined in columns 8 and 9, "NT From" and "NT To" respectively, in Table 2.

[00825] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[00826] Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[00827] A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of the portion of SEQ ID NO:X defined in column 4, "ORF (From-To)", in Table 1A.

[00828] A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of the portion of SEQ ID NO:X defined in columns 8 and 9, "NT From" and "NT To", respectively, in Table 2.

[00829] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z.

[00830] Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, the

nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto, and/or cDNA contained in Clone ID NO:Z, wherein said nucleic acid molecule which hybridizes does not hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

[00831] Also preferred is a composition of matter comprising a DNA molecule which comprises the cDNA contained in Clone ID NO:Z.

[00832] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides of the cDNA sequence contained in Clone ID NO:Z.

[00833] Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of an open reading frame sequence encoded by cDNA contained in Clone ID NO:Z.

[00834] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[00835] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[00836] A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by cDNA contained in Clone ID NO:Z.

[00837] A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence encoded by cDNA contained in Clone ID NO:Z; which method comprises a step of comparing a nucleotide sequence of at least one nucleic acid molecule in said sample with a sequence selected from said group and determining

10094504, 030702



whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

[00838] Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

[00839] A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence of the cDNA contained in Clone ID NO:Z.

[00840] The method for identifying the species, tissue or cell type of a biological sample can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

[00841] Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto; or the cDNA contained in Clone ID NO:Z which encodes a protein, wherein the method comprises a step of detecting in a biological sample obtained from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 4 of

Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence of cDNA contained in Clone ID NO:Z.

[00842] The method for diagnosing a pathological condition can comprise a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

[00843] Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; the nucleotide sequence as defined in column 4 of Table 1A or columns 8 and 9 of Table 2 or the complementary strand thereto; and a nucleotide sequence encoded by cDNA contained in Clone ID NO:Z. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

[00844] Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a DNA microarray or "chip" of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 100, 150, 200, 250, 300, 500, 1000, 2000, 3000, or 4000 nucleotide sequences, wherein at least one sequence in said DNA microarray or "chip" is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X wherein X is any integer as defined in Table 1A; and a nucleotide sequence encoded by a human cDNA clone identified by a cDNA "Clone ID" in Table 1A.

[00845] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[00846] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the

amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[00847] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[00848] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[00849] Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a polypeptide encoded by contained in Clone ID NO:Z

[00850] Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a portion of said polypeptide encoded by cDNA contained in Clone ID NO:Z; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and/or the polypeptide sequence of SEQ ID NO:Y.

[00851] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00852] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of a polypeptide encoded by cDNA contained in Clone ID NO:Z.

[00853] Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00854] Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00855] Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

[00856] Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00857] Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in said sample with said sequence selected from said group.

[00858] Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00859] Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

[00860] Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleic acid sequence identified in Table 1A or Table 2 encoding a polypeptide, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00861] In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

[00862] Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at

least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00863] Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

[00864] Also preferred is a polypeptide molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z.

[00865] Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a recombinant host cell comprising introducing the vector into a host cell, as well as the recombinant host cell produced by this method.

[00866] Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a human protein comprising an amino acid sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto; the polypeptide encoded by the nucleotide sequence as defined in columns 8 and 9 of Table 2; and a polypeptide encoded by the cDNA contained in Clone ID NO:Z. The isolated polypeptide produced by this method is also preferred.

[00867] Also preferred is a method of treatment of an individual in need of an increased level of a protein activity, which method comprises administering to such an individual a Therapeutic comprising an amount of an isolated polypeptide,

polynucleotide, immunogenic fragment or analogue thereof, binding agent, antibody, or antigen binding fragment of the claimed invention effective to increase the level of said protein activity in said individual.

[00868] Also preferred is a method of treatment of an individual in need of a decreased level of a protein activity, which method comprised administering to such an individual a Therapeutic comprising an amount of an isolated polypeptide, polynucleotide, immunogenic fragment or analogue thereof, binding agent, antibody, or antigen binding fragment of the claimed invention effective to decrease the level of said protein activity in said individual.

[00869] Also preferred is a method of treatment of an individual in need of a specific delivery of toxic compositions to diseased cells (e.g., tumors, leukemias or lymphomas), which method comprises administering to such an individual a Therapeutic comprising an amount of an isolated polypeptide of the invention, including, but not limited to a binding agent, or antibody of the claimed invention that are associated with toxin or cytotoxic prodrugs.

[00870] Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

**TABLE 6**

ATCC Deposits	Deposit Date	ATCC Designation Number
LP01, LP02, LP03, LP04, LP05, LP06, LP07, LP08, LP09, LP10, LP11,	May-20-97	209059, 209060, 209061, 209062, 209063, 209064, 209065, 209066, 209067, 209068, 209069
LP12	Jan-12-98	209579
LP13	Jan-12-98	209578
LP14	Jul-16-98	203067
LP15	Jul-16-98	203068
LP16	Feb-1-99	203609
LP17	Feb-1-99	203610
LP20	Nov-17-98	203485

LP21	Jun-18-99	PTA-252
LP22	Jun-18-99	PTA-253
LP23	Dec-22-99	PTA-1081

10094504 030700



## Examples

### *Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample*

[00871] Each Clone ID NO:Z is contained in a plasmid. Table 7 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The following correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 7 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited Plasmid</u>
Lambda Zap	pBluescript (pBS)
Uni-Zap XR	pBluescript (pBS)
Zap Express	pBK
lafmid BA	plafmid BA
pSport1	pSport1
pCMVSport 2.0	pCMVSport 2.0
pCMVSport 3.0	pCMVSport 3.0
pCR®2.1	pCR®2.1

[00872] Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the orientation of the fl origin of replication ("ori"), such that in one

orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.

[00873] Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR<sup>®</sup>2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the vector sequences identified for the particular clone in Table 7, as well as the corresponding plasmid vector sequences designated above.

[00874] The deposited material in the sample assigned the ATCC Deposit Number cited by reference to Tables 1A, 2, 6 and 7 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each Clone ID NO:Z.

**TABLE 7**

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HUKA HUKB HUKC HUKD HUKF HUKG	Human Uterine Cancer	Lambda ZAP II	LP01
HCNA HCNB	Human Colon	Lambda Zap II	LP01
HFFA	Human Fetal Brain, random primed	Lambda Zap II	LP01
HTWA	Resting T-Cell	Lambda ZAP II	LP01
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP01
HLMB HLMF HLMG HLMH HLMJ HLMM HLMN	breast lymph node CDNA library	Lambda ZAP II	LP01
HCQA HCQB	human colon cancer	Lamda ZAP II	LP01
HMEA HMEC HMED HMEE	Human Microvascular Endothelial	Lambda ZAP II	LP01

10091504 030702

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HMEF HMEG HMEI HMEJ HMEK HMEL	Cells, fract. A		
HUSA HUSC	Human Umbilical Vein Endothelial Cells, fract. A	Lambda ZAP II	LP01
HLQA HLQB	Hepatocellular Tumor	Lambda ZAP II	LP01
HHGA HHGB HHGC HHGD	Hemangiopericytoma	Lambda ZAP II	LP01
HSDM	Human Striatum Depression, re-rescue	Lambda ZAP II	LP01
HUSH	H Umbilical Vein Endothelial Cells, fract A, re-excision	Lambda ZAP II	LP01
HSGS	Salivary gland, subtracted	Lambda ZAP II	LP01
HFXA HFXB HFXC HFXD HFXE HFXF HFXG HFXH	Brain frontal cortex	Lambda ZAP II	LP01
HPQA HPQB HPQC	PERM TF274	Lambda ZAP II	LP01
HFXJ HFXK	Brain Frontal Cortex, re-excision	Lambda ZAP II	LP01
HCWA HCWB HCWC HCWD HCWE HCWF HCWG HCWH HCWI HCWJ HCWK	CD34 positive cells (Cord Blood)	ZAP Express	LP02
HCUA HCUB HCUC	CD34 depleted Buffy Coat (Cord Blood)	ZAP Express	LP02
HRSM	A-14 cell line	ZAP Express	LP02
HRSA	A1-CELL LINE	ZAP Express	LP02
HCUD HCUE HCUF HCUG HCUH HCUI	CD34 depleted Buffy Coat (Cord Blood), re-excision	ZAP Express	LP02
HBXE HBXF HBXG	H. Whole Brain #2, re-excision	ZAP Express	LP02
HRLM	L8 cell line	ZAP Express	LP02
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP02
HUDA HUDB HUDC	Testes	ZAP Express	LP02
HHTM HHTN HHTO	H. hypothalamus, frac A; re-excision	ZAP Express	LP02
HHTL	H. hypothalamus, frac A	ZAP Express	LP02
HASA HASD	Human Adult Spleen	Uni-ZAP XR	LP03
HFKC HFKD HFKE HFKF HFKG	Human Fetal Kidney	Uni-ZAP XR	LP03
HE8A HE8B HE8C HE8D HE8E HE8F HE8M HE8N	Human 8 Week Whole Embryo	Uni-ZAP XR	LP03
HGBA HGBD HGBE HGBF HGBG HGBH HGBI	Human Gall Bladder	Uni-ZAP XR	LP03
HLHA HLHB HLHC HLHD HLHE HLHF HLHG HLHH HLHQ	Human Fetal Lung III	Uni-ZAP XR	LP03
HPMA HPMB HPMC HPMD	Human Placenta	Uni-ZAP XR	LP03

10091504.030702

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HPME HPMF HPMG HPMH			
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP03
HSIA HSIC HSID HSIE	Human Adult Small Intestine	Uni-ZAP XR	LP03
HTEA HTEB HTEC HTED	Human Testes	Uni-ZAP XR	LP03
HTEE HTEF HTEG HTEH HTEI HTEJ HTEK			
HTPA HTPB HTPC HTPD HTPE	Human Pancreas Tumor	Uni-ZAP XR	LP03
HTTA HTTB HTTC HTTD HTTE HTTF	Human Testes Tumor	Uni-ZAP XR	LP03
HAPA HAPB HAPC HAPM	Human Adult Pulmonary	Uni-ZAP XR	LP03
HETA HETB HETC HETD HETE HETF HETG HETH HETI	Human Endometrial Tumor	Uni-ZAP XR	LP03
HHFB HHFC HHFD HHFE HHFF HHFG HHFH HHFI	Human Fetal Heart	Uni-ZAP XR	LP03
HHPB HNPC HHPD HHPE HHPF HHPG HHPH	Human Hippocampus	Uni-ZAP XR	LP03
HCE1 HCE2 HCE3 HCE4 HCE5 HCEB HCEC HCED HCEE HCEF HCEG	Human Cerebellum	Uni-ZAP XR	LP03
HUVB HUV C HUVD HUVE	Human Umbilical Vein, Endo. remake	Uni-ZAP XR	LP03
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP03
HTAA HTAB HTAC HTAD HTAE	Human Activated T-Cells	Uni-ZAP XR	LP03
HFEA HFEB HFEC	Human Fetal Epithelium (Skin)	Uni-ZAP XR	LP03
HJPA HJPB HJPC HJPD	HUMAN JURKAT MEMBRANE BOUND POLYSOMES	Uni-ZAP XR	LP03
HESA	Human epithelioid sarcoma	Uni-Zap XR	LP03
HLTA HLTB HLTC HLTD HLTE HLTF	Human T-Cell Lymphoma	Uni-ZAP XR	LP03
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP03
HRDA HRDB HRDC HRDD HRDE HRDF	Human Rhabdomyosarcoma	Uni-ZAP XR	LP03
HCAA HCAB HCAC	Cem cells cyclohexamide treated	Uni-ZAP XR	LP03
HRGA HRGB HRGC HRGD	Raji Cells, cyclohexamide treated	Uni-ZAP XR	LP03
HSUA HSUB HSUC HSUM	Supt Cells, cyclohexamide treated	Uni-ZAP XR	LP03
HT4A HT4C HT4D	Activated T-Cells, 12 hrs.	Uni-ZAP XR	LP03
HE9A HE9B HE9C HE9D HE9E HE9F HE9G HE9H HE9M HE9N	Nine Week Old Early Stage Human	Uni-ZAP XR	LP03
HATA HATB HATC HATD	Human Adrenal Gland Tumor	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HATE			
HT5A	Activated T-Cells, 24 hrs.	Uni-ZAP XR	LP03
HFGA HFGM	Human Fetal Brain	Uni-ZAP XR	LP03
HNEA HNEB HNEC HNED HNEE	Human Neutrophil	Uni-ZAP XR	LP03
HBGB HBGD	Human Primary Breast Cancer	Uni-ZAP XR	LP03
HBNA HBNB	Human Normal Breast	Uni-ZAP XR	LP03
HCAS	Cem Cells, cyclohexamide treated, subtra	Uni-ZAP XR	LP03
HHPS	Human Hippocampus, subtracted	pBS	LP03
HKCS HKCU	Human Colon Cancer, subtracted	pBS	LP03
HRGS	Raji cells, cyclohexamide treated, subtracted	pBS	LP03
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBS	LP03
HT4S	Activated T-Cells, 12 hrs, subtracted	Uni-ZAP XR	LP03
HCDA HCDB HCDC HCDD HCDE	Human Chondrosarcoma	Uni-ZAP XR	LP03
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP03
HTLA HTLB HTLC HTLD HTLE HTLF	Human adult testis, large inserts	Uni-ZAP XR	LP03
HLMA HLMC HLMD	Breast Lymph node cDNA library	Uni-ZAP XR	LP03
H6EA H6EB H6EC	HL-60, PMA 4H	Uni-ZAP XR	LP03
HTXA HTXB HTXC HTXD HTXE HTXF HTXG HTXH	Activated T-Cell (12hs)/Thiouridine labelledEco	Uni-ZAP XR	LP03
HNFA HNFB HNFC HNFD HNFE HNFF HNFG HNFH HNFJ	Human Neutrophil, Activated	Uni-ZAP XR	LP03
HTOB HTOC	HUMAN TONSILS, FRACTION 2	Uni-ZAP XR	LP03
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP03
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP03
HORB	Human OB HOS treated (10 nM E2) fraction I	Uni-ZAP XR	LP03
HSVA HSVB HSVC	Human Chronic Synovitis	Uni-ZAP XR	LP03
HROA	HUMAN STOMACH	Uni-ZAP XR	LP03
HBJA HBJB HBJC HBJD HBJE HBJF HBJG HBJH HBJI HBJJ HBJK	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP03
HCRA HCRB HCRC	human corpus colosum	Uni-ZAP XR	LP03
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HDSA	Dermatofibrosarcoma Protuberance	Uni-ZAP XR	LP03
HMWA HMWB HMWC HMWD HMWE HMWF HMWG HMWH HMWI HMWJ	Bone Marrow Cell Line (RS4;11)	Uni-ZAP XR	LP03
HSOA	stomach cancer (human)	Uni-ZAP XR	LP03
HERA	SKIN	Uni-ZAP XR	LP03
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP03
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP03
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP03
HBCA HBCB	H. Lymph node breast Cancer	Uni-ZAP XR	LP03
HPWT	Human Prostate BPH, re-excision	Uni-ZAP XR	LP03
HFVG HFVH HFVI	Fetal Liver, subtraction II	pBS	LP03
HNFI	Human Neutrophils, Activated, re-excision	pBS	LP03
HBMB HBMC HBMD	Human Bone Marrow, re-excision	pBS	LP03
HKML HKMM HKMN	H. Kidney Medulla, re-excision	pBS	LP03
HKIX HKIY	H. Kidney Cortex, subtracted	pBS	LP03
HADT	H. Amygdala Depression, subtracted	pBS	LP03
H6AS	HL-60, untreated, subtracted	Uni-ZAP XR	LP03
H6ES	HL-60, PMA 4H, subtracted	Uni-ZAP XR	LP03
H6BS	HL-60, RA 4h, Subtracted	Uni-ZAP XR	LP03
H6CS	HL-60, PMA 1d, subtracted	Uni-ZAP XR	LP03
HTXJ HTXK	Activated T-cell(12h)/Thiouridine-re-excision	Uni-ZAP XR	LP03
HMSA HMSB HMSC HMSD HMSE HMSF HMSG HMSH HMSI HMSJ HMSK	Monocyte activated	Uni-ZAP XR	LP03
HAGA HAGB HAGC HAGD HAGE HAGF	Human Amygdala	Uni-ZAP XR	LP03
HSRA HSRB HSRE	STROMAL -OSTEOCLASTOMA	Uni-ZAP XR	LP03
HSRD HSRF HSRG HSRH	Human Osteoclastoma Stromal Cells - unamplified	Uni-ZAP XR	LP03
HSQA HSQB HSQC HSQD HSQE HSQF HSQG	Stromal cell TF274	Uni-ZAP XR	LP03
HSKA HSKB HSKC HSKD HSKE HSKF HSKZ	Smooth muscle, serum treated	Uni-ZAP XR	LP03
HSLA HSLB HSLC HSLD HSLE HSLF HSLG	Smooth muscle, control	Uni-ZAP XR	LP03
HSDA HSDD HSDE HSDF HSDG HSDH	Spinal cord	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HPWS	Prostate-BPH subtracted II	pBS	LP03
HSKW HSKX HSKY	Smooth Muscle- HASTE normalized	pBS	LP03
HFPB HFPC HFPD	H. Frontal cortex,epileptic;re-excision	Uni-ZAP XR	LP03
HSDI HSDJ HSDK	Spinal Cord, re-excision	Uni-ZAP XR	LP03
HSKN HSKO	Smooth Muscle Serum Treated, Norm	pBS	LP03
HSKG HSKH HSKI	Smooth muscle, serum induced,re-exc	pBS	LP03
HFCA HFCB HFCC HFCD HFCE HFCE	Human Fetal Brain	Uni-ZAP XR	LP04
HPTA HPTB HPTD	Human Pituitary	Uni-ZAP XR	LP04
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP04
HE6B HE6C HE6D HE6E HE6F HE6G HE6S	Human Whole Six Week Old Embryo	Uni-ZAP XR	LP04
HSSA HSSB HSSC HSSD HSSE HSSF HSSG HSSH HSSI HSSJ HSSK	Human Synovial Sarcoma	Uni-ZAP XR	LP04
HE7T	7 Week Old Early Stage Human, subtracted	Uni-ZAP XR	LP04
HEPA HEPB HEPC	Human Epididymus	Uni-ZAP XR	LP04
HSNA HSNB HSNB HSNM HSNN	Human Synovium	Uni-ZAP XR	LP04
HPFB HPFC HPFD HPFE	Human Prostate Cancer, Stage C fraction	Uni-ZAP XR	LP04
HE2A HE2D HE2E HE2H HE2I HE2M HE2N HE2O	12 Week Old Early Stage Human	Uni-ZAP XR	LP04
HE2B HE2C HE2F HE2G HE2P HE2Q	12 Week Old Early Stage Human, II	Uni-ZAP XR	LP04
HPTS HPTT HPTU	Human Pituitary, subtracted	Uni-ZAP XR	LP04
HAUA HAUB HAUC	Amniotic Cells - TNF induced	Uni-ZAP XR	LP04
HAQA HAQB HAQC HAQD	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP04
HUTA HWTB HWTG	wilm's tumor	Uni-ZAP XR	LP04
HBSD	Bone Cancer, re-excision	Uni-ZAP XR	LP04
HSGB	Salivary gland, re-excision	Uni-ZAP XR	LP04
HSJA HSJB HSJC	Smooth muscle-ILb induced	Uni-ZAP XR	LP04
HSXA HSXB HSXC HSXD	Human Substantia Nigra	Uni-ZAP XR	LP04
HSJA HSJB HSJC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP04
HOUA HOUB HOUH HOUD HOUE	Adipocytes	Uni-ZAP XR	LP04
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP04
HELA HELB HELC HELD	Endothelial cells-control	Uni-ZAP XR	LP04

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HELE HELF HELG HELH			
HEMA HEMB HEMC HEMD HEME HEMF HEMG HEMH	Endothelial-induced	Uni-ZAP XR	LP04
HBIA HBIB HBIC	Human Brain, Striatum	Uni-ZAP XR	LP04
HHSA HHSB HHSC HHSD HHSE	Human Hypothalamus, Schizophrenia	Uni-ZAP XR	LP04
HNGA HNGB HNGC HNGD HNGE HNGF HNGG HNGH HNGI HNGJ	neutrophils control	Uni-ZAP XR	LP04
HNHA HNHB HNHC HNHD HNHE HNHF HNHG HNHH HNHI HNHI	Neutrophils IL-1 and LPS induced	Uni-ZAP XR	LP04
HSDB HSDC	STRIATUM DEPRESSION	Uni-ZAP XR	LP04
HHPT	Hypothalamus	Uni-ZAP XR	LP04
HSAT HSAU HSAV HSAW HSAX HSAY HSAZ	Anergic T-cell	Uni-ZAP XR	LP04
HBMS HBMT HBMU HBMV HBMW HBMX	Bone marrow	Uni-ZAP XR	LP04
HOEA HOEB HOEC HOED HOEE HOEF HOEJ	Osteoblasts	Uni-ZAP XR	LP04
HAIA HAIB HAIC HAID HAIE HAIF	Epithelial-TNF $\alpha$ and INF induced	Uni-ZAP XR	LP04
HTGA HTGB HTGC HTGD	Apoptotic T-cell	Uni-ZAP XR	LP04
HMCA HMCB HMCC HMCD HMCE	Macrophage-oxLDL	Uni-ZAP XR	LP04
HMAA HMAB HMAB HMAD HMAE HMAF HMAG	Macrophage (GM-CSF treated)	Uni-ZAP XR	LP04
HPHA	Normal Prostate	Uni-ZAP XR	LP04
HPIA HPIB HPIC	LNCAP prostate cell line	Uni-ZAP XR	LP04
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP04
HOSE HOSF HOSG	Human Osteoclastoma, re-excision	Uni-ZAP XR	LP04
HTGE HTGF	Apoptotic T-cell, re-excision	Uni-ZAP XR	LP04
HMAJ HMAK	H Macrophage (GM-CSF treated), re-excision	Uni-ZAP XR	LP04
HACB HACC HACD	Human Adipose Tissue, re-excision	Uni-ZAP XR	LP04
HFPA	H. Frontal Cortex, Epileptic	Uni-ZAP XR	LP04
HFAA HFAB HFAC HFAD HFAE	Alzheimers, spongy change	Uni-ZAP XR	LP04
HFAM	Frontal Lobe, Dementia	Uni-ZAP XR	LP04
HMIA HMIB HMIC	Human Manic Depression Tissue	Uni-ZAP XR	LP04
HTSA HTSE HTSF HTSG	Human Thymus	pBS	LP05



10091504-030702

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HTSH			
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBS	LP05
HSAA HSAB HSAC	HSA 172 Cells	pBS	LP05
HSBA HSBB HSBC HSBM	HSC172 cells	pBS	LP05
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBS	LP05
HJBA HJBB HJBC HJBD	Jurkat T-Cell, S phase	pBS	LP05
HAFA HAFB	Aorta endothelial cells + TNF-a	pBS	LP05
HAWA HAWB HAWC	Human White Adipose	pBS	LP05
HTNA HTNB	Human Thyroid	pBS	LP05
HONA	Normal Ovary, Premenopausal	pBS	LP05
HARA HARB	Human Adult Retina	pBS	LP05
HLJA HLJB	Human Lung	pCMVSPORT 1	LP06
HOFM HOFN HOFO	H. Ovarian Tumor, II, OV5232	pCMVSPORT 2.0	LP07
HOGA HOGB HOGC	OV 10-3-95	pCMVSPORT 2.0	LP07
HCGL	CD34+cells, II	pCMVSPORT 2.0	LP07
HDLA	Hodgkin's Lymphoma I	pCMVSPORT 2.0	LP07
HDTA HDTB HDTC HDTD HDTE	Hodgkin's Lymphoma II	pCMVSPORT 2.0	LP07
HKAA HKAB HKAC HKAD HKAH HKAF HKAG HKAH	Keratinocyte	pCMVSPORT2.0	LP07
HCIM	CAPFINDER, Crohn's Disease, lib 2	pCMVSPORT 2.0	LP07
HKAL	Keratinocyte, lib 2	pCMVSPORT2.0	LP07
HKAT	Keratinocyte, lib 3	pCMVSPORT2.0	LP07
HNDA	Nasal polyps	pCMVSPORT2.0	LP07
HDRA	H. Primary Dendritic Cells, lib 3	pCMVSPORT2.0	LP07
HOHA HOHB HOHC	Human Osteoblasts II	pCMVSPORT2.0	LP07
HLDA HLDB HLDC	Liver, Hepatoma	pCMVSPORT3.0	LP08
HLDN HLDO HLDP	Human Liver, normal	pCMVSPORT3.0	LP08
HMTA	pBMC stimulated w/ poly I/C	pCMVSPORT3.0	LP08
HNTA	NTERA2, control	pCMVSPORT3.0	LP08
HDPH HDPB HDPD HDPD HDPF HDPG HDPH HDPH HDPJ HDPK	Primary Dendritic Cells, lib 1	pCMVSPORT3.0	LP08
HDPH HDPN HDPO HDPP	Primary Dendritic cells, frac 2	pCMVSPORT3.0	LP08
HMUA HMUB HMUC	Myeloid Progenitor Cell Line	pCMVSPORT3.0	LP08
HHEA HHEB HHEC HHED	T Cell helper I	pCMVSPORT3.0	LP08
HHEM HHEN HHEO HHEP	T cell helper II	pCMVSPORT3.0	LP08
HEQA HEQB HEQC	Human endometrial stromal cells	pCMVSPORT3.0	LP08
HJMA HJMB	Human endometrial stromal cells-	pCMVSPORT3.0	LP08

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
	treated with progesterone		
HSWA HSWB HSWC	Human endometrial stromal cells- treated with estradiol	pCMVSPORT3.0	LP08
HSYA HSYB HSYC	Human Thymus Stromal Cells	pCMVSPORT3.0	LP08
HLWA HLWB HLWC	Human Placenta	pCMVSPORT3.0	LP08
HRAA HRAB HRAC	Rejected Kidney, lib 4	pCMVSPORT3.0	LP08
HMTM	PCR, pBMC I/C treated	PCR11	LP09
HMJA	H. Meningioma, M6	pSport 1	LP10
HMKA HMKB HMKC HMKD HMKE	H. Meningioma, M1	pSport 1	LP10
HUSG HUSI	Human umbilical vein endothelial cells, IL-4 induced	pSport 1	LP10
HUSX HUSY	Human Umbilical Vein Endothelial Cells, uninduced	pSport 1	LP10
HOFA	Ovarian Tumor I, OV5232	pSport 1	LP10
HCFA HCFB HCFC HCFD	T-Cell PHA 16 hrs	pSport 1	LP10
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport 1	LP10
HADA HADC HADD HADE HADG HADG	Human Adipose	pSport 1	LP10
HOVA HOVB HOVC	Human Ovary	pSport 1	LP10
HTWB HTWC HTWD HTWE HTWF	Resting T-Cell Library, II	pSport 1	LP10
HMMMA	Spleen metastatic melanoma	pSport 1	LP10
HLYA HLYB HLYC HLYD HLYE	Spleen, Chronic lymphocytic leukemia	pSport 1	LP10
HCGA	CD34+ cell, I	pSport 1	LP10
HEOM HEON	Human Eosinophils	pSport 1	LP10
HTDA	Human Tonsil, Lib 3	pSport 1	LP10
HSPA	Salivary Gland, Lib 2	pSport 1	LP10
HCHA HCHB HCHC	Breast Cancer cell line, MDA 36	pSport 1	LP10
HCHM HCHN	Breast Cancer Cell line, angiogenic	pSport 1	LP10
HCIA	Crohn's Disease	pSport 1	LP10
HDAA HDAB HDAC	HEL cell line	pSport 1	LP10
HABA	Human Astrocyte	pSport 1	LP10
HUFA HUFB HUFC	Ulcerative Colitis	pSport 1	LP10
HNTM	NTERA2 + retinoic acid, 14 days	pSport 1	LP10
HDQA	Primary Dendritic cells, CapFinder2, frac 1	pSport 1	LP10
HDQM	Primary Dendritic Cells, CapFinder, frac 2	pSport 1	LP10

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HLDX	Human Liver, normal, CapFinder	pSport 1	LP10
HULA HULB HULC	Human Dermal Endothelial Cells, untreated	pSport1	LP10
HUMA	Human Dermal Endothelial cells, treated	pSport1	LP10
HCJA	Human Stromal Endometrial fibroblasts, untreated	pSport1	LP10
HCJM	Human Stromal endometrial fibroblasts, treated w/ estradiol	pSport1	LP10
HEDA	Human Stromal endometrial fibroblasts, treated with progesterone	pSport1	LP10
HFNA	Human ovary tumor cell OV350721	pSport1	LP10
HKGA HKGB HKGC HKGD	Merkel Cells	pSport1	LP10
HISA HISB HISC	Pancreas Islet Cell Tumor	pSport1	LP10
HLSA	Skin, burned	pSport1	LP10
HBZA	Prostate, BPH, Lib 2	pSport 1	LP10
HBZS	Prostate BPH, Lib 2, subtracted	pSport 1	LP10
HFIA HFIB HFIC	Synovial Fibroblasts (control)	pSport 1	LP10
HFIH HFII HFIJ	Synovial hypoxia	pSport 1	LP10
HFIT HFIU HFIV	Synovial IL-1/TNF stimulated	pSport 1	LP10
HGCA	Mesangial cell, frac 1	pSport1	LP10
HMVA HMVB HMVC	Bone Marrow Stromal Cell, untreated	pSport1	LP10
HFIY HFIZ	Synovial Fibroblasts (IL1/TNF), sub	pSport1	LP10
HFOX HFOY HFOZ	Synovial hypoxia-RSF subtracted	pSport1	LP10
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP11
HLIA HLIB HLIC	Human Liver	pCMVSPORT 1	LP012
HHBA HHBB HHBC HHBD HHBE	Human Heart	pCMVSPORT 1	LP012
HBBA HBBB	Human Brain	pCMVSPORT 1	LP012
HLJA HLJB HLJC HLJD HLJE	Human Lung	pCMVSPORT 1	LP012
HOGA HOGB HOGC	Ovarian Tumor	pCMVSPORT 2.0	LP012
HTJM	Human Tonsils, Lib 2	pCMVSPORT 2.0	LP012
HAMF HAMG	KMH2	pCMVSPORT 3.0	LP012
HAJA HAJB HAJC	L428	pCMVSPORT 3.0	LP012
HWBA HWBB HWBC HWBD HWBE	Dendritic cells, pooled	pCMVSPORT 3.0	LP012
HWAA HWAB HWAC HWAD HWAE	Human Bone Marrow, treated	pCMVSPORT 3.0	LP012
HYAA HYAB HYAC	B Cell lymphoma	pCMVSPORT 3.0	LP012
HWHG HWHH HWHI	Healing groin wound, 6.5 hours post	pCMVSPORT 3.0	LP012

10004504 40576001

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
	incision		
HWHP HWHQ HWHR	Healing groin wound; 7.5 hours post incision	pCMVSPORT 3.0	LP012
HARM	Healing groin wound - zero hr post-incision (control)	pCMVSPORT 3.0	LP012
HBIM	Olfactory epithelium; nasalcavity	pCMVSPORT 3.0	LP012
HWDA	Healing Abdomen wound; 70&90 min post incision	pCMVSPORT 3.0	LP012
HWEA	Healing Abdomen Wound;15 days post incision	pCMVSPORT 3.0	LP012
HWJA	Healing Abdomen Wound;21&29 days	pCMVSPORT 3.0	LP012
HNAL	Human Tongue, frac 2	pSport1	LP012
HMJA	H. Meningima, M6	pSport1	LP012
HMKA HMKB HMKC HMKD HMKF	H. Meningima, M1	pSport1	LP012
HOFA	Ovarian Tumor I, OV5232	pSport1	LP012
HCFA HCFB HCFC HCFD	T-Cell PHA 16 hrs	pSport1	LP012
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport1	LP012
HMMA HMMB HMMC	Spleen metastatic melanoma	pSport1	LP012
HTDA	Human Tonsil, Lib 3	pSport1	LP012
HDBA	Human Fetal Thymus	pSport1	LP012
HDUA	Pericardium	pSport1	LP012
HBZA	Prostate,BPH, Lib 2	pSport1	LP012
HWCA	Larynx tumor	pSport1	LP012
HWKA	Normal lung	pSport1	LP012
HSMB	Bone marrow stroma,treated	pSport1	LP012
HBHM	Normal trachea	pSport1	LP012
HLFC	Human Larynx	pSport1	LP012
HLRB	Siebben Polyposis	pSport1	LP012
HNIA	Mammary Gland	pSport1	LP012
HNJB	Palate carcinoma	pSport1	LP012
HNKA	Palate normal	pSport1	LP012
HMZA	Pharynx carcinoma	pSport1	LP012
HABG	Cheek Carcinoma	pSport1	LP012
HMZM	Pharynx Carcinoma	pSport1	LP012
HDRM	Larynx Carcinoma	pSport1	LP012
HVAA	Pancreas normal PCA4 No	pSport1	LP012
HICA	Tongue carcinoma	pSport1	LP012
HUKA HUKB HUKC HUKD HUKF	Human Uterine Cancer	Lambda ZAP II	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HFFA	Human Fetal Brain, random primed	Lambda ZAP II	LP013
HTUA	Activated T-cell labeled with 4-thioluri	Lambda ZAP II	LP013
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP013
HMEB	Human microvascular Endothelial cells, fract. B	Lambda ZAP II	LP013
HUSH	Human Umbilical Vein Endothelial cells, fract. A, re-excision	Lambda ZAP II	LP013
HLQC HLQD	Hepatocellular tumor, re-excision	Lambda ZAP II	LP013
HTWJ HTWK HTWL	Resting T-cell, re-excision	Lambda ZAP II	LP013
HF6S	Human Whole 6 week Old Embryo (II), subt	pBluescript	LP013
HHPS	Human Hippocampus, subtracted	pBluescript	LP013
HLIS	LNCAP, differential expression	pBluescript	LP013
HLHS HLHT	Early Stage Human Lung, Subtracted	pBluescript	LP013
HSUS	Supt cells, cyclohexamide treated, subtracted	pBluescript	LP013
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBluescript	LP013
HSDS	H. Striatum Depression, subtracted	pBluescript	LP013
HPTZ	Human Pituitary, Subtracted VII	pBluescript	LP013
HSDX	H. Striatum Depression, subt II	pBluescript	LP013
HSDZ	H. Striatum Depression, subt	pBluescript	LP013
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBluescript SK-	LP013
HRTA	Colorectal Tumor	pBluescript SK-	LP013
HSBA HSBB HSBC HSBM	HSC172 cells	pBluescript SK-	LP013
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBluescript SK-	LP013
HJBA HJBB HJBC HJBD	Jurkat T-cell, S1 phase	pBluescript SK-	LP013
HTNA HTNB	Human Thyroid	pBluescript SK-	LP013
HAHA HAHB	Human Adult Heart	Uni-ZAP XR	LP013
HE6A	Whole 6 week Old Embryo	Uni-ZAP XR	LP013
HFCA HFCB HFCC HFCD HFCE	Human Fetal Brain	Uni-ZAP XR	LP013
HFKE HFKE HFKE HFKE HFKE	Human Fetal Kidney	Uni-ZAP XR	LP013
HGBA HGBD HGBE HGBF HGBG	Human Gall Bladder	Uni-ZAP XR	LP013
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP013
HTEA HTEB HTEC HTED	Human Testes	Uni-ZAP XR	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HTEE			
HTTA HTTB HTTC HTTD HTTE	Human Testes Tumor	Uni-ZAP XR	LP013
HYBA HYBB	Human Fetal Bone	Uni-ZAP XR	LP013
HFLA	Human Fetal Liver	Uni-ZAP XR	LP013
HHFB HHFC HHFD HHFE HHFF	Human Fetal Heart	Uni-ZAP XR	LP013
HUVB HUV C HUVD HUVE	Human Umbilical Vein, End. remake	Uni-ZAP XR	LP013
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP013
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP013
HTAA HTAB HTAC HTAD HTAE	Human Activated T-cells	Uni-ZAP XR	LP013
HFEA HFEB HFEC	Human Fetal Epithelium (skin)	Uni-ZAP XR	LP013
HJPA HJPB HJPC HJPD	Human Jurkat Membrane Bound Polysomes	Uni-ZAP XR	LP013
HESA	Human Epithelioid Sarcoma	Uni-ZAP XR	LP013
HALS	Human Adult Liver, Subtracted	Uni-ZAP XR	LP013
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP013
HCAA HCAB HCAC	Cem cells, cyclohexamide treated	Uni-ZAP XR	LP013
HRGA HRGB HRGC HRGD	Raji Cells, cyclohexamide treated	Uni-ZAP XR	LP013
HE9A HE9B HE9C HE9D HE9E	Nine Week Old Early Stage Human	Uni-ZAP XR	LP013
HSFA	Human Fibrosarcoma	Uni-ZAP XR	LP013
HATA HATB HATC HATD HATE	Human Adrenal Gland Tumor	Uni-ZAP XR	LP013
HTRA	Human Trachea Tumor	Uni-ZAP XR	LP013
HE2A HE2D HE2E HE2H HE2I	12 Week Old Early Stage Human	Uni-ZAP XR	LP013
HE2B HE2C HE2F HE2G HE2P	12 Week Old Early Stage Human, II	Uni-ZAP XR	LP013
HNEA HNEB HNEC HNED HNEE	Human Neutrophil	Uni-ZAP XR	LP013
HBGA	Human Primary Breast Cancer	Uni-ZAP XR	LP013
HPTS HPTT HPTU	Human Pituitary, subtracted	Uni-ZAP XR	LP013
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP013
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP013
HTOA HTOD HTOE HTOF HTOG	human tonsils	Uni-ZAP XR	LP013
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP013
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP013
HOQB	Human OB HOS treated (1 nM E2) fraction I	Uni-ZAP XR	LP013
HAUA HAUB HAUC	Amniotic Cells - TNF induced	Uni-ZAP XR	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HAQA HAQB HAQC HAQD	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP013
HROA HROC	HUMAN STOMACH	Uni-ZAP XR	LP013
HBJA HBJB HBJC HBJD HBJE	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP013
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP013
HCPA	Corpus Callosum	Uni-ZAP XR	LP013
HSOA	stomach cancer (human)	Uni-ZAP XR	LP013
HERA	SKIN	Uni-ZAP XR	LP013
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP013
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP013
HWTA HWTB HWTC	wilm's tumor	Uni-ZAP XR	LP013
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP013
HAPN HAPO HAPR HAPQ	Human Adult Pulmonary;re-excision	Uni-ZAP XR	LP013
HLTG HLTH	Human T-cell lymphoma;re-excision	Uni-ZAP XR	LP013
HAHC HAHD HAHE	Human Adult Heart;re-excision	Uni-ZAP XR	LP013
HAGA HAGB HAGC HAGD HAGE	Human Amygdala	Uni-ZAP XR	LP013
HSJA HSJB HSJC	Smooth muscle-ILb induced	Uni-ZAP XR	LP013
HSHA HSHB HSHC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP013
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP013
HPJA HPJB HPJC	LNCAP prostate cell line	Uni-ZAP XR	LP013
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP013
HBTA	Bone Marrow Stroma, TNF&LPS ind	Uni-ZAP XR	LP013
HMCF HMCJ HMCH HMCJ	Macrophage-oxLDL; re-excision	Uni-ZAP XR	LP013
HAGG HAGH HAGI	Human Amygdala;re-excision	Uni-ZAP XR	LP013
HACA	H. Adipose Tissue	Uni-ZAP XR	LP013
HKFB	K562 + PMA (36 hrs),re-excision	ZAP Express	LP013
HCWT HCWU HCWV	CD34 positive cells (cord blood),re-ex	ZAP Express	LP013
HBWA	Whole brain	ZAP Express	LP013
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP013
HAVM	Temporal cortex-Alzheimer	pT-Adv	LP014
HAVT	Hippocampus, Alzheimer Subtracted	pT-Adv	LP014
HHAS	CHME Cell Line	Uni-ZAP XR	LP014
HAJR	Larynx normal	pSport 1	LP014
HWLE HWLF HWLG HWLH	Colon Normal	pSport 1	LP014
HCRM HCRN HCRO	Colon Carcinoma	pSport 1	LP014
HWLI HWLJ HWLK	Colon Normal	pSport 1	LP014

2020020405T500T

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HWLQ HWLR HWLS HWLT	Colon Tumor	pSport 1	LP014
HBFM	Gastrocnemius Muscle	pSport 1	LP014
HBOD HBOE	Quadriceps Muscle	pSport 1	LP014
HBKD HBKE	Soleus Muscle	pSport 1	LP014
HCCM	Pancreatic Langerhans	pSport 1	LP014
HWGA	Larynx carcinoma	pSport 1	LP014
HWGM HWGN	Larynx carcinoma	pSport 1	LP014
HWLA HWLB HWLC	Normal colon	pSport 1	LP014
HWLM HWLN	Colon Tumor	pSport 1	LP014
HVAM HVAN HVAO	Pancreas Tumor	pSport 1	LP014
HWGQ	Larynx carcinoma	pSport 1	LP014
HAQM HAQN	Salivary Gland	pSport 1	LP014
HASM	Stomach; normal	pSport 1	LP014
HBCM	Uterus; normal	pSport 1	LP014
HCDM	Testis; normal	pSport 1	LP014
HDJM	Brain; normal	pSport 1	LP014
HEFM	Adrenal Gland,normal	pSport 1	LP014
HBAA	Rectum normal	pSport 1	LP014
HFDM	Rectum tumour	pSport 1	LP014
HGAM	Colon, normal	pSport 1	LP014
HHMM	Colon, tumour	pSport 1	LP014
HCLB HCLC	Human Lung Cancer	Lambda Zap II	LP015
HRLA	L1 Cell line	ZAP Express	LP015
HHAM	Hypothalamus, Alzheimer's	pCMVSport 3.0	LP015
HKBA	Ku 812F Basophils Line	pSport 1	LP015
HS2S	Saos2, Dexamethosome Treated	pSport 1	LP016
HA5A	Lung Carcinoma A549 TNFalpha activated	pSport 1	LP016
HTFM	TF-1 Cell Line GM-CSF Treated	pSport 1	LP016
HYAS	Thyroid Tumour	pSport 1	LP016
HUTS	Larynx Normal	pSport 1	LP016
HXOA	Larynx Tumor	pSport 1	LP016
HEAH	Ea.hy.926 cell line	pSport 1	LP016
HINA	Adenocarcinoma Human	pSport 1	LP016
HRMA	Lung Mesothelium	pSport 1	LP016
HLCL	Human Pre-Differentiated Adipocytes	Uni-Zap XR	LP017
HS2A	Saos2 Cells	pSport 1	LP020
HS2I	Saos2 Cells; Vitamin D3 Treated	pSport 1	LP020
HUCM	CHME Cell Line, untreated	pSport 1	LP020
HEPN	Aryepiglottis Normal	pSport 1	LP020



10091504-030703

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HPSN	Sinus Piniformis Tumour	pSport 1	LP020
HNSA	Stomach Normal	pSport 1	LP020
HNSM	Stomach Tumour	pSport 1	LP020
HNLA	Liver Normal Met5No	pSport 1	LP020
HUTA	Liver Tumour Met 5 Tu	pSport 1	LP020
HOCN	Colon Normal	pSport 1	LP020
HOCT	Colon Tumor	pSport 1	LP020
HTNT	Tongue Tumour	pSport 1	LP020
HLXN	Larynx Normal	pSport 1	LP020
HLXT	Larynx Tumour	pSport 1	LP020
HTYN	Thymus	pSport 1	LP020
HPLN	Placenta	pSport 1	LP020
HTNG	Tongue Normal	pSport 1	LP020
HZAA	Thyroid Normal (SDCA2 No)	pSport 1	LP020
HWES	Thyroid Thyroiditis	pSport 1	LP020
HFHD	Ficoll Human Stromal Cells, 5Fu treated	pTriplEx2	LP021
HFHM,HFHN	Ficoll Human Stromal Cells, Untreated	pTriplEx2	LP021
HPCI	Hep G2 Cells, lambda library	lambda Zap-CMV XR	LP021
HBCA,HBCB,HBCC	H. Lymph node breast Cancer	Uni-ZAP XR	LP021
HCOK	Chondrocytes	pSPORT1	LP022
HDCA, HDCB, HDCC	Dendritic Cells From CD34 Cells	pSPORT1	LP022
HDMA, HDMB	CD40 activated monocyte dendritic cells	pSPORT1	LP022
HDDM, HDDN, HDDO	LPS activated derived dendritic cells	pSPORT1	LP022
HPCR	Hep G2 Cells, PCR library	lambda Zap-CMV XR	LP022
HAAA, HAAB, HAAC	Lung, Cancer (4005313A3): Invasive Poorly Differentiated Lung Adenocarcinoma	pSPORT1	LP022
HIPA, HIPB, HIPC	Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic	pSPORT1	LP022
HOOH, HOOI	Ovary, Cancer: (4004562 B6) Papillary Serous Cystic Neoplasm, Low Malignant Pot	pSPORT1	LP022
HIDA	Lung, Normal: (4005313 B1)	pSPORT1	LP022
HUJA,HUJB,HUJC,HUJD,HUJE	B-Cells	pCMVSPORT 3.0	LP022
HNOA,HNOB,HNOC,HNOD	Ovary, Normal: (9805C040R)	pSPORT1	LP022
HNLM	Lung, Normal: (4005313 B1)	pSPORT1	LP022

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HSCL	Stromal Cells	pSPORT1	LP022
HAAX	Lung, Cancer: (4005313 A3) Invasive Poorly-differentiated Metastatic lung adenocarcinoma	pSPORT1	LP022
HUUA,HUUB,HUUC,HUUD	B-cells (unstimulated)	pTriplEx2	LP022
HWWA,HWWB,HWWC,HWW D,HWWE,HWWF,HWWG	B-cells (stimulated)	pSPORT1	LP022
HCCC	Colon, Cancer: (9808C064R)	pCMVSPORT 3.0	LP023
HPDO HPDP HPDQ HPDR HPD	Ovary, Cancer (9809C332): Poorly differentiated adenocarcinoma	pSport 1	LP023
HPCO HPCP HPCQ HPCT	Ovary, Cancer (15395A1F): Grade II Papillary Carcinoma	pSport 1	LP023
HOCM HOCO HOC P HOCQ	Ovary, Cancer: (15799A1F) Poorly differentiated carcinoma	pSport 1	LP023
HCBM HCBN HCBO	Breast, Cancer: (4004943 A5)	pSport 1	LP023
HNBT HNBU HNBV	Breast, Normal: (4005522B2)	pSport 1	LP023
HBCP HBCQ	Breast, Cancer: (4005522 A2)	pSport 1	LP023
HBCJ	Breast, Cancer: (9806C012R)	pSport 1	LP023
HSAM HSAN	Stromal cells 3.88	pSport 1	LP023
HVCA HVCB HVCC HVCD	Ovary, Cancer: (4004332 A2)	pSport 1	LP023
HSCK HSEN HSEO	Stromal cells (HBM3.18)	pSport 1	LP023
HSCP HSCQ	stromal cell clone 2.5	pSport 1	LP023
HUXA	Breast Cancer: (4005385 A2)	pSport 1	LP023
HCOM HCON HCOO HCOP HCOQ	Ovary, Cancer (4004650 A3): Well-Differentiated Micropapillary Serous Carcinoma	pSport 1	LP023
HBNM	Breast, Cancer: (9802C020E)	pSport 1	LP023
HVVA HVVB HVVC HVVD HVVE	Human Bone Marrow, treated	pSport 1	LP023

[00875] Two nonlimiting examples are provided below for isolating a particular clone from the deposited sample of plasmid cDNAs cited for that clone in Table 7. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to the nucleotide sequence of SEQ ID NO:X.

[00876] Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with <sup>32</sup>P-γ-ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et

al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

[00877] Alternatively, two primers of 17-20 nucleotides derived from both ends of the nucleotide sequence of SEQ ID NO:X are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25  $\mu$ l of reaction mixture with 0.5  $\mu$ g of the above cDNA template. A convenient reaction mixture is 1.5-5 mM  $MgCl_2$ , 0.01% (w/v) gelatin, 20  $\mu$ M each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

[00878] Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not limited to, filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

[00879] Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set

10091504.030702

containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full length gene.

[00880] This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

[00881] This modified RNA preparation is used as a template for first strand cDNA synthesis using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

### ***Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide***

[00882] A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the sequence corresponding to SEQ ID NO:X according to the method described in Example 1. (See also, Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Edn., (1989), Cold Spring Harbor Laboratory Press).

### ***Example 3: Tissue specific expression analysis***

[00883] The Human Genome Sciences, Inc. (HGS) database is derived from sequencing tissue and/or disease specific cDNA libraries. Libraries generated from a particular tissue are selected and the specific tissue expression pattern of EST groups or assembled contigs within these libraries is determined by comparison of the

expression patterns of those groups or contigs within the entire database. ESTs and assembled contigs which show tissue specific expression are selected.

[00884] The original clone from which the specific EST sequence was generated, or in the case of an assembled contig, the clone from which the 5' most EST sequence was generated, is obtained from the catalogued library of clones and the insert amplified by PCR using methods known in the art. The PCR product is denatured and then transferred in 96 or 384 well format to a nylon membrane (Schleicher and Scheull) generating an array filter of tissue specific clones. Housekeeping genes, maize genes, and known tissue specific genes are included on the filters. These targets can be used in signal normalization and to validate assay sensitivity. Additional targets are included to monitor probe length and specificity of hybridization.

[00885] Radioactively labeled hybridization probes are generated by first strand cDNA synthesis per the manufacturer's instructions (Life Technologies) from mRNA/RNA samples prepared from the specific tissue being analyzed (e.g., heart, heart valve, vascular system, heart cancer, prostate, prostate cancer, ovarian, ovarian cancer, etc.). The hybridization probes are purified by gel exclusion chromatography, quantitated, and hybridized with the array filters in hybridization bottles at 65°C overnight. The filters are washed under stringent conditions and signals are captured using a Fuji phosphorimager.

[00886] Data is extracted using AIS software and following background subtraction, signal normalization is performed. This includes a normalization of filter-wide expression levels between different experimental runs. Genes that are differentially expressed in the tissue of interest are identified.

#### *Example 4: Chromosomal Mapping of the Polynucleotides*

[00887] An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions: 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated 32 times followed by one 5 minute cycle at 70°C. Human, mouse, and hamster DNA is used

as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions are analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR fragment in the particular somatic cell hybrid.

### ***Example 5: Bacterial Expression of a Polypeptide***

[00888] A polynucleotide encoding a polypeptide of the present invention is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp<sup>r</sup>), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

[00889] The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan<sup>r</sup>). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is isolated and confirmed by restriction analysis.

[00890] Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.<sup>600</sup>) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto pyranoside) is then added to a final concentration of 1

mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

[00891] Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

[00892] Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8. The column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8, then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

[00893] The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl. The purified protein is stored at 4°C or frozen at -80°C.

[00894] In addition to the above expression vector, the present invention further includes an expression vector, called pHE4a (ATCC Accession Number 209645, deposited on February 25, 1998) which contains phage operator and promoter elements operatively linked to a polynucleotide of the present invention. This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (lacIq). The origin of replication (oriC) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter and operator sequences are made synthetically.

[00895] DNA can be inserted into the pHE4a by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

[00896] The engineered vector could easily be substituted in the above protocol to express protein in a bacterial system.

### ***Example 6: Purification of a Polypeptide from an Inclusion Body***

[00897] The following alternative method can be used to purify a polypeptide expressed in *E. coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

[00898] Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

[00899] The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

[00900] The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.



- 100901-030700
- [00901] Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.
- [00902] To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16  $\mu$ m membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.
- [00903] Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant  $A_{280}$  monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.
- [00904] The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5  $\mu$ g of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

**Example 7: Cloning and Expression of a Polypeptide in a Baculovirus  
Expression System**

- [00905] In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.
- [00906] Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).
- [00907] Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon, is amplified using the PCR protocol described in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present invention, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).
- [00908] The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

[00909] The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

[00910] The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

[00911] Five  $\mu$ g of a plasmid-containing the polynucleotide is co-transfected with 1.0  $\mu$ g of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7417 (1987). One  $\mu$ g of BaculoGold™ virus DNA and 5  $\mu$ g of the plasmid are mixed in a sterile well of a microtiter plate containing 50  $\mu$ l of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10  $\mu$ l Lipofectin plus 90  $\mu$ l Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

[00912] After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant

viruses is then resuspended in a microcentrifuge tube containing 200  $\mu$ l of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

[00913] To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5  $\mu$ Ci of  $^{35}$ S-methionine and 5  $\mu$ Ci  $^{35}$ S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

[00914] Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

### *Example 8: Expression of a Polypeptide in Mammalian Cells*

[00915] The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV1, HIV1 and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

[00916] Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109),

pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

[00917] Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as DHFR, gpt, neomycin, or hygromycin allows the identification and isolation of the transfected cells.

[00918] The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

[00919] Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No.209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

[00920] Specifically, the plasmid pC6, for example, is digested with appropriate restriction enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

[00921] A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present invention, the vector does not need a second signal peptide. Alternatively, if a naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., International Publication No. WO 96/34891.)

[00922] The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

[00923] The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

[00924] Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five  $\mu$ g of the expression plasmid pC6 or pC4 is cotransfected with 0.5  $\mu$ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1  $\mu$ M, 2  $\mu$ M, 5  $\mu$ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200  $\mu$ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

### Example 9: Protein Fusions

[00925] The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time *in vivo*. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

[00926] Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

[00927] For example, if pC4 (ATCC Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without a stop codon, otherwise a fusion protein will not be produced.

[00928] If the naturally occurring signal sequence is used to produce the polypeptide of the present invention, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., International Publication No. WO 96/34891.)

Human IgG Fc region:

GGGATCCGGAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCCCA  
GCACCTGAATTCGAGGGTGCACCGTCAGTCTTCCTCTTCCCCC AAAACCCAAG  
GACACCCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGTGGACGTA  
AGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGAGGT  
GCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTG  
TGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTAC  
AAGTGCAAGGTCTCCAACAAAGCCCTCCCAACCCCCATCGAGAAAACCATCTC  
CAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCC  
GGGATGAGCTGACCAAGAACCAGGTGACCTGACCTGCCTGGTCAAAGGCTTC  
TATCCAAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAA  
CTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAG  
CAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCT  
CCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGT  
CTCCGGGTAAATGAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO: 1)

### *Example 10: Production of an Antibody from a Polypeptide*

#### Hybridoma Technology

[00929] The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing polypeptide of the present invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide of the present invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

[00930] Monoclonal antibodies specific for polypeptide of the present invention are prepared using hybridoma technology (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In general, an animal (preferably a mouse) is immunized with polypeptide of the present invention or, more preferably, with a



secreted polypeptide of the present invention-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

[00931] The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones, which secrete antibodies capable of binding the polypeptide of the present invention.

[00932] Alternatively, additional antibodies capable of binding to polypeptide of the present invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody, which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the polypeptide of the present invention-specific antibody can be blocked by polypeptide of the present invention. Such antibodies comprise anti-idiotypic antibodies to the polypeptide of the present invention-specific antibody and are used to immunize an animal to induce formation of further polypeptide of the present invention-specific antibodies.

[00933] For *in vivo* use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric and humanized antibodies are known in the art and are discussed herein. (See, for review, Morrison, Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et

al., International Publication No. WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

Isolation Of Antibody Fragments Directed Against Polypeptide of the Present Invention  
From A Library Of scFvs

[00934] Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against polypeptide of the present invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

[00935] *Rescue of the Library.* A library of scFvs is constructed from the RNA of human PBLs as described in International Publication No. WO 92/01047. To rescue phage displaying antibody fragments, approximately  $10^9$  *E. coli* harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU,  $2 \times 10^8$  TU of delta gene 3 helper (M13 delta gene III, see International Publication No. WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 µg/ml ampicillin and 50 µg/ml kanamycin and grown overnight. Phage are prepared as described in International Application No. WO 92/01047.

[00936] M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations (Sambrook et al., 1990), resuspended in 2 ml PBS and passed through

a 0.45  $\mu\text{m}$  filter (Minisart NML; Sartorius) to give a final concentration of approximately  $10^{13}$  transducing units/ml (ampicillin-resistant clones).

[00937] *Panning of the Library.* Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100  $\mu\text{g/ml}$  or 10  $\mu\text{g/ml}$  of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately  $10^{13}$  TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100  $\mu\text{g/ml}$  ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

[00938] *Characterization of Binders.* Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with 10  $\mu\text{g/ml}$  of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., International Application No. WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

*Example 11: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide*

- 10091504-030702
- [00939] RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X; and/or the nucleotide sequence of the cDNA contained in Clone ID NO:Z. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).
- [00940] PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase (Epicentre Technologies). The intron-exon boundaries of selected exons is also determined and genomic PCR products analyzed to confirm the results. PCR products harboring suspected mutations are then cloned and sequenced to validate the results of the direct sequencing.
- [00941] PCR products are cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected individuals.
- [00942] Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenindeoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991). Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.
- [00943] Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et

al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions, deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

***Example 12: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample***

[00944] A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

[00945] For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

[00946] The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbound polypeptide.

[00947] Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbound conjugate.

[00948] Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis

(log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

### *Example 13: Formulation*

[00949] The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a Therapeutic. By therapeutic is meant polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

[00950] The Therapeutic will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

[00951] As a general proposition, the total pharmaceutically effective amount of the Therapeutic administered parenterally per dose will be in the range of about 1 ug/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 ug/kg/hour to about 50 ug/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

[00952] Therapeutics can be administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[00953] Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

[00954] Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

[00955] Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., *Biopolymers* 22:547-556 (1983)), poly (2-hydroxyethyl methacrylate) (Langer et al., *J. Biomed. Mater. Res.* 15:167-277 (1981), and Langer, *Chem. Tech.* 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., *Id.*) or poly-D-(-)-3-hydroxybutyric acid (EP 133,988).

[00956] Sustained-release Therapeutics also include liposomally entrapped Therapeutics of the invention (see generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317-327 and 353-365 (1989)). Liposomes

containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci.(USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal Therapeutic.

[00957] In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

[00958] Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

[00959] For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

[00960] Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

[00961] The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten



10091504 030702

residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

[00962] The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

[00963] Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[00964] Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is lyophilized. The infusion solution is prepared by reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

[00965] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention. Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

[00966] The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG (e.g.,

10091504-030702

THERACYS®), MPL and nonviable preparations of *Corynebacterium parvum*. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to, vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B, whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

[00967] The Therapeutics of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not limited to, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, and/or therapeutic treatments described below. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

10094504-030702  
202009105T60T

[00968] In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), and/or protease inhibitors (PIs). NRTIs that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). NNRTIs that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIXIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

[00969] Additional NRTIs include LODENOSINE™ (F-ddA; an acid-stable adenosine NRTI; Triangle/Abbott; COVIRACIL™ (emtricitabine/FTC; structurally related to lamivudine (3TC) but with 3- to 10-fold greater activity *in vitro*; Triangle/Abbott); dOTC (BCH-10652, also structurally related to lamivudine but retains activity against a substantial proportion of lamivudine-resistant isolates; Biochem Pharma); Adefovir (refused approval for anti-HIV therapy by FDA; Gilead Sciences); PREVEON® (Adefovir Dipivoxil, the active prodrug of adefovir; its active form is PMEA-pp); TENOFOVIR™ (bis-POC PMPA, a PMPA prodrug; Gilead); DAPD/DXG (active metabolite of DAPD; Triangle/Abbott); D-D4FC (related to 3TC, with activity against AZT/3TC-resistant virus); GW420867X (Glaxo Wellcome); ZIAGEN™ (abacavir/159U89; Glaxo Wellcome Inc.); CS-87 (3'-azido-2',3'-dideoxyuridine; WO 99/66936); and S-acyl-2-thioethyl (SATE)-bearing prodrug forms of  $\beta$ -L-FD4C and  $\beta$ -L-FddC (WO 98/17281).

[00970] Additional NNRTIs include COACTINON™ (Emivirine/MKC-442, potent NNRTI of the HEPT class; Triangle/Abbott); CAPRAVIRINE™ (AG-1549/S-1153,

a next generation NNRTI with activity against viruses containing the K103N mutation; Agouron); PNU-142721 (has 20- to 50-fold greater activity than its predecessor delavirdine and is active against K103N mutants; Pharmacia & Upjohn); DPC-961 and DPC-963 (second-generation derivatives of efavirenz, designed to be active against viruses with the K103N mutation; DuPont); GW-420867X (has 25-fold greater activity than HBY097 and is active against K103N mutants; Glaxo Wellcome); CALANOLIDE A (naturally occurring agent from the latex tree; active against viruses containing either or both the Y181C and K103N mutations); and Propolis (WO 99/49830).

[00971] Additional protease inhibitors include LOPINAVIR™ (ABT378/r; Abbott Laboratories); BMS-232632 (an azapeptide; Bristol-Myers Squibb); TIPRANAVIR™ (PNU-140690, a non-peptic dihydropyrone; Pharmacia & Upjohn); PD-178390 (a nonpeptidic dihydropyrone; Parke-Davis); BMS 232632 (an azapeptide; Bristol-Myers Squibb); L-756,423 (an indinavir analog; Merck); DMP-450 (a cyclic urea compound; Avid & DuPont); AG-1776 (a peptidomimetic with *in vitro* activity against protease inhibitor-resistant viruses; Agouron); VX-175/GW-433908 (phosphate prodrug of amprenavir; Vertex & Glaxo Wellcome); CGP61755 (Ciba); and AGENERASE™ (amprenavir; Glaxo Wellcome Inc.).

[00972] Additional antiretroviral agents include fusion inhibitors/gp41 binders. Fusion inhibitors/gp41 binders include T-20 (a peptide from residues 643-678 of the HIV gp41 transmembrane protein ectodomain which binds to gp41 in its resting state and prevents transformation to the fusogenic state; Trimeris) and T-1249 (a second-generation fusion inhibitor; Trimeris).

[00973] Additional antiretroviral agents include fusion inhibitors/chemokine receptor antagonists. Fusion inhibitors/chemokine receptor antagonists include CXCR4 antagonists such as AMD 3100 (a bicyclam), SDF-1 and its analogs, and ALX40-4C (a cationic peptide), T22 (an 18 amino acid peptide; Trimeris) and the T22 analogs T134 and T140; CCR5 antagonists such as RANTES (9-68), AOP-RANTES, NNY-RANTES, and TAK-779; and CCR5/CXCR4 antagonists such as NSC 651016 (a distamycin analog). Also included are CCR2B, CCR3, and CCR6 antagonists. Chemokine receptor agonists such as RANTES, SDF-1, MIP-1 $\alpha$ , MIP-1 $\beta$ , etc., may also inhibit fusion.

[00974] Additional antiretroviral agents include integrase inhibitors. Integrase inhibitors include dicaffeoylquinic (DFQA) acids; L-chicoric acid (a dicaffeoyltartaric (DCTA) acid); quinalizarin (QLC) and related anthraquinones; ZINTEVIR™ (AR 177, an oligonucleotide that probably acts at cell surface rather than being a true integrase inhibitor; Arondex); and naphthols such as those disclosed in WO 98/50347.

[00975] Additional antiretroviral agents include hydroxyurea-like compounds such as BCX-34 (a purine nucleoside phosphorylase inhibitor; Biocryst); ribonucleotide reductase inhibitors such as DIDOX™ (Molecules for Health); inosine monophosphate dehydrogenase (IMPDH) inhibitors such as VX-497 (Vertex); and mycophenolic acids such as CellCept (mycophenolate mofetil; Roche).

[00976] Additional antiretroviral agents include inhibitors of viral integrase, inhibitors of viral genome nuclear translocation such as arylene bis(methylketone) compounds; inhibitors of HIV entry such as AOP-RANTES, NNY-RANTES, RANTES-IgG fusion protein, soluble complexes of RANTES and glycosaminoglycans (GAG), and AMD-3100; nucleocapsid zinc finger inhibitors such as dithiane compounds; targets of HIV Tat and Rev; and pharmacoenhancers such as ABT-378.

[00977] Other antiretroviral therapies and adjunct therapies include cytokines and lymphokines such as MIP-1 $\alpha$ , MIP-1 $\beta$ , SDF-1 $\alpha$ , IL-2, PROLEUKIN™ (aldesleukin/L2-7001; Chiron), IL-4, IL-10, IL-12, and IL-13; interferons such as IFN- $\alpha$ 2a; antagonists of TNFs, NF $\kappa$ B, GM-CSF, M-CSF, and IL-10; agents that modulate immune activation such as cyclosporin and prednisone; vaccines such as Remune™ (HIV Immunogen), APL 400-003 (Apollon), recombinant gp120 and fragments, bivalent (B/E) recombinant envelope glycoprotein, rgp120CM235, MN rgp120, SF-2 rgp120, gp120/soluble CD4 complex, Delta JR-FL protein, branched synthetic peptide derived from discontinuous gp120 C3/C4 domain, fusion-competent immunogens, and Gag, Pol, Nef, and Tat vaccines; gene-based therapies such as genetic suppressor elements (GSEs; WO 98/54366), and intrakines (genetically modified CC chemokines targetted to the ER to block surface expression of newly synthesized CCR5 (Yang *et al.*, *PNAS* 94:11567-72 (1997); Chen *et al.*, *Nat. Med.* 3:1110-16 (1997)); antibodies such as the anti-CXCR4 antibody 12G5, the anti-CCR5 antibodies 2D7, 5C7, PA8, PA9, PA10, PA11, PA12, and PA14, the anti-CD4 antibodies Q4120 and RPA-T4, the anti-CCR3 antibody

7B11, the anti-gp120 antibodies 17b, 48d, 447-52D, 257-D, 268-D and 50.1, anti-Tat antibodies, anti-TNF- $\alpha$  antibodies, and monoclonal antibody 33A; aryl hydrocarbon (AH) receptor agonists and antagonists such as TCDD, 3,3',4,4',5-pentachlorobiphenyl, 3,3',4,4'-tetrachlorobiphenyl, and  $\alpha$ -naphthoflavone (WO 98/30213); and antioxidants such as  $\gamma$ -L-glutamyl-L-cysteine ethyl ester ( $\gamma$ -GCE; WO 99/56764).

[00978] In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

[00979] In other embodiments, Therapeutics of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICLOVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment, Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment,

10093504-030702

Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat or prevent an opportunistic cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial infection.

[00980] In a further embodiment, the Therapeutics of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin, beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin, erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rapamycin, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

[00981] In other embodiments, Therapeutics of the invention are administered in combination with immunosuppressive agents. Immunosuppressive agents that may be administered in combination with the Therapeutics of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells. Other immunosuppressive agents that may be administered in combination with the Therapeutics of the invention include, but are not limited to, prednisolone, methotrexate, thalidomide, methoxsalen, rapamycin, leflunomide, mizoribine (BREDININ™), brequinar, deoxyspergualin, and azaspirane (SKF 105685),

ORTHOCLONE OKT® 3 (muromonab-CD3), SANDIMMUNE™, NEORAL™, SANGDYA™ (cyclosporine), PROGRAF® (FK506, tacrolimus), CELLCEPT® (mycophenolate mofetil, of which the active metabolite is mycophenolic acid), IMURAN™ (azathioprine), glucocorticosteroids, adrenocortical steroids such as DELTASONE™ (prednisone) and HYDELTRASOL™ (prednisolone), FOLEX™ and MEXATE™ (methotrexate), OXSORALEN-ULTRA™ (methoxsalen) and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

[00982] In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, ATGAM™ (antithymocyte globulin), and GAMIMUNE™. In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

[00983] In certain embodiments, the Therapeutics of the invention are administered alone or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, corticosteroids (e.g. betamethasone, budesonide, cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, and triamcinolone), nonsteroidal anti-inflammatory drugs (e.g., diclofenac, diflunisal, etodolac, fenoprofen, floctafenine, flurbiprofen, ibuprofen, indomethacin, ketoprofen, meclofenamate, mefenamic acid, meloxicam, nabumetone, naproxen, oxaprozin, phenylbutazone, piroxicam, sulindac, tenoxicam, tiaprofenic acid, and tolmetin.), as well as antihistamines, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.



[00984] In an additional embodiment, the compositions of the invention are administered alone or in combination with an anti-angiogenic agent. Anti-angiogenic agents that may be administered with the compositions of the invention include, but are not limited to, Angiostatin (Entremed, Rockville, MD), Troponin-1 (Boston Life Sciences, Boston, MA), anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel (Taxol), Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, VEGF, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

[00985] Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

[00986] Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

[00987] Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

**[00988]** A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include, but are not limited to, platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., *Cancer Res.* 51:22-26, (1991)); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., *J. Bio. Chem.* 267:17321-17326, (1992)); Chymostatin (Tomkinson et al., *Biochem J.* 286:475-480, (1992)); Cyclodextrin Tetradecasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., *Nature* 348:555-557, (1990)); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, *J. Clin. Invest.* 79:1440-1446, (1987)); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., *J. Biol. Chem.* 262(4):1659-1664, (1987)); Bisantrone (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4- chloroanthronilic acid disodium or "CCA"; (Takeuchi et al., *Agents Actions* 36:312-316, (1992)); and metalloproteinase inhibitors such as BB94.

**[00989]** Additional anti-angiogenic factors that may also be utilized within the context of the present invention include Thalidomide, (Celgene, Warren, NJ); Angiostatic steroid; AGM-1470 (H. Brem and J. Folkman *J Pediatr. Surg.* 28:445-51 (1993)); an integrin alpha v beta 3 antagonist (C. Storgard et al., *J Clin. Invest.* 103:47-54 (1999)); carboxynaminolmidazole; Carboxyamidotriazole (CAI) (National Cancer Institute, Bethesda, MD); Conbretastatin A-4 (CA4P) (OXiGENE, Boston, MA); Squalamine (Magainin Pharmaceuticals, Plymouth Meeting, PA); TNP-470, (Tap Pharmaceuticals, Deerfield, IL); ZD-0101 AstraZeneca (London, UK); APRA (CT2584); Benefin, Byrostatin-1 (SC339555); CGP-41251 (PKC 412); CM101; Dexrazoxane (ICRF187); DMXAA; Endostatin; Flavopridiol; Genestein; GTE; ImmTher; Iressa (ZD1839); Octreotide (Somatostatin); Panretin; Penacillamine; Photopoint; PI-88; Prinomastat (AG-3340) Purlytin; Suradista (FCE26644); Tamoxifen (Nolvadex); Tazarotene; Tetrathiomolybdate; Xeloda (Capecitabine); and 5-Fluorouracil.

10091504-030702

[00990] Anti-angiogenic agents that may be administered in combination with the compounds of the invention may work through a variety of mechanisms including, but not limited to, inhibiting proteolysis of the extracellular matrix, blocking the function of endothelial cell-extracellular matrix adhesion molecules, by antagonizing the function of angiogenesis inducers such as growth factors, and inhibiting integrin receptors expressed on proliferating endothelial cells. Examples of anti-angiogenic inhibitors that interfere with extracellular matrix proteolysis and which may be administered in combination with the compositions of the invention include, but are not limited to, AG-3340 (Agouron, La Jolla, CA), BAY-12-9566 (Bayer, West Haven, CT), BMS-275291 (Bristol Myers Squibb, Princeton, NJ), CGS-27032A (Novartis, East Hanover, NJ), Marimastat (British Biotech, Oxford, UK), and Metastat (Aeterna, St-Foy, Quebec). Examples of anti-angiogenic inhibitors that act by blocking the function of endothelial cell-extracellular matrix adhesion molecules and which may be administered in combination with the compositions of the invention include, but are not limited to, EMD-121974 (Merck KgaA Darmstadt, Germany) and Vitaxin (Ixsys, La Jolla, CA/Medimmune, Gaithersburg, MD). Examples of anti-angiogenic agents that act by directly antagonizing or inhibiting angiogenesis inducers and which may be administered in combination with the compositions of the invention include, but are not limited to, Angiozyme (Ribozyme, Boulder, CO), Anti-VEGF antibody (Genentech, S. San Francisco, CA), PTK-787/ZK-225846 (Novartis, Basel, Switzerland), SU-101 (Sugen, S. San Francisco, CA), SU-5416 (Sugen/ Pharmacia Upjohn, Bridgewater, NJ), and SU-6668 (Sugen). Other anti-angiogenic agents act to indirectly inhibit angiogenesis. Examples of indirect inhibitors of angiogenesis which may be administered in combination with the compositions of the invention include, but are not limited to, IM-862 (Cytran, Kirkland, WA), Interferon-alpha, IL-12 (Roche, Nutley, NJ), and Pentosan polysulfate (Georgetown University, Washington, DC).

[00991] In particular embodiments, the use of compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of an autoimmune disease, such as for example, an autoimmune disease described herein.

[00992] In a particular embodiment, the use of compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment,

prevention, and/or amelioration of arthritis. In a more particular embodiment, the use of compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of rheumatoid arthritis.

[00993] In another embodiment, the polynucleotides encoding a polypeptide of the present invention are administered in combination with an angiogenic protein, or polynucleotides encoding an angiogenic protein. Examples of angiogenic proteins that may be administered with the compositions of the invention include, but are not limited to, acidic and basic fibroblast growth factors, VEGF-1, VEGF-2, VEGF-3, epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin-like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

[00994] In additional embodiments, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to alkylating agents such as nitrogen mustards (for example, Mechlorethamine, cyclophosphamide, Cyclophosphamide Ifosfamide, Melphalan (L-sarcosine), and Chlorambucil), ethylenimines and methylmelamines (for example, Hexamethylmelamine and Thiotepa), alkyl sulfonates (for example, Busulfan), nitrosoureas (for example, Carmustine (BCNU), Lomustine (CCNU), Semustine (methyl-CCNU), and Streptozocin (streptozotocin)), triazines (for example, Dacarbazine (DTIC; dimethyltriazenoimidazolecarboxamide)), folic acid analogs (for example, Methotrexate (amethopterin)), pyrimidine analogs (for example, Fluorouracil (5-fluorouracil; 5-FU), Floxuridine (fluorodeoxyuridine; FudR), and Cytarabine (cytosine arabinoside)), purine analogs and related inhibitors (for example, Mercaptopurine (6-mercaptopurine; 6-MP), Thioguanine (6-thioguanine; TG), and Pentostatin (2'-deoxycoformycin)), vinca alkaloids (for example, Vinblastine (VLB, vinblastine sulfate) and Vincristine (vincristine sulfate)), epipodophyllotoxins (for example, Etoposide and Teniposide), antibiotics (for example, Dactinomycin (actinomycin D), Daunorubicin (daunomycin; rubidomycin), Doxorubicin, Bleomycin, Plicamycin (mithramycin), and Mitomycin

40091504-030702

(mitomycin C), enzymes (for example, L-Asparaginase), biological response modifiers (for example, Interferon-alpha and interferon-alpha-2b), platinum coordination compounds (for example, Cisplatin (cis-DDP) and Carboplatin), anthracenedione (Mitoxantrone), substituted ureas (for example, Hydroxyurea), methylhydrazine derivatives (for example, Procarbazine (N-methylhydrazine; MIH), adrenocorticosteroids (for example, Prednisone), progestins (for example, Hydroxyprogesterone caproate, Medroxyprogesterone, Medroxyprogesterone acetate, and Megestrol acetate), estrogens (for example, Diethylstilbestrol (DES), Diethylstilbestrol diphosphate, Estradiol, and Ethinyl estradiol), antiestrogens (for example, Tamoxifen), androgens (Testosterone propionate, and Fluoxymesterone), antiandrogens (for example, Flutamide), gonadotropin-releasing hormone analogs (for example, Leuprolide), other hormones and hormone analogs (for example, methyltestosterone, estramustine, estramustine phosphate sodium, chlorotrianisene, and testolactone), and others (for example, dicarbazine, glutamic acid, and mitotane).

[00995] In one embodiment, the compositions of the invention are administered in combination with one or more of the following drugs: infliximab (also known as Remicade™ Centocor, Inc.), Trocade (Roche, RO-32-3555), Leflunomide (also known as Arava™ from Hoechst Marion Roussel), Kineret™ (an IL-1 Receptor antagonist also known as Anakinra from Amgen, Inc.).

[00996] In a specific embodiment, compositions of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or combination of one or more of the components of CHOP. In one embodiment, the compositions of the invention are administered in combination with anti-CD20 antibodies, human monoclonal anti-CD20 antibodies. In another embodiment, the compositions of the invention are administered in combination with anti-CD20 antibodies and CHOP, or anti-CD20 antibodies and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. In a specific embodiment, compositions of the invention are administered in combination with Rituximab. In a further embodiment, compositions of the invention are administered with Rituximab and CHOP, or Rituximab and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. In a specific embodiment,

compositions of the invention are administered in combination with tositumomab. In a further embodiment, compositions of the invention are administered with tositumomab and CHOP, or tositumomab and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. The anti-CD20 antibodies may optionally be associated with radioisotopes, toxins or cytotoxic prodrugs.

[00997] In another specific embodiment, the compositions of the invention are administered in combination Zevalin™. In a further embodiment, compositions of the invention are administered with Zevalin™ and CHOP, or Zevalin™ and any combination of one or more of the components of CHOP, particularly cyclophosphamide and/or prednisone. Zevalin™ may be associated with one or more radisotopes. Particularly preferred isotopes are <sup>90</sup>Y and <sup>111</sup>In.

[00998] In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

[00999] In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-IBB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TRANK, TR9 (International

Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

[001000] In an additional embodiment, the Therapeutics of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PlGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PlGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are herein incorporated by reference in their entireties.

[001001] In an additional embodiment, the Therapeutics of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

[001002] In an additional embodiment, the Therapeutics of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, granulocyte macrophage colony stimulating factor

(GM-CSF) (sargramostim, LEUKINE™, PROKINE™), granulocyte colony stimulating factor (G-CSF) (filgrastim, NEUPOGEN™), macrophage colony stimulating factor (M-CSF, CSF-1) erythropoietin (epoetin alfa, EPOGEN™, PROCRIT™), stem cell factor (SCF, c-kit ligand, steel factor), megakaryocyte colony stimulating factor, PIXY321 (a GMCSF/IL-3 fusion protein), interleukins, especially any one or more of IL-1 through IL-12, interferon-gamma, or thrombopoietin.

[001003] In certain embodiments, Therapeutics of the present invention are administered in combination with adrenergic blockers, such as, for example, acebutolol, atenolol, betaxolol, bisoprolol, carteolol, labetalol, metoprolol, nadolol, oxprenolol, penbutolol, pindolol, propranolol, sotalol, and timolol.

[001004] In another embodiment, the Therapeutics of the invention are administered in combination with an antiarrhythmic drug (e.g., adenosine, amiodarone, bretylium, digitalis, digoxin, digitoxin, diltiazem, disopyramide, esmolol, flecainide, lidocaine, mexiletine, moricizine, phenytoin, procainamide, N-acetyl procainamide, propafenone, propranolol, quinidine, sotalol, tocainide, and verapamil).

[001005] In another embodiment, the Therapeutics of the invention are administered in combination with diuretic agents, such as carbonic anhydrase-inhibiting agents (e.g., acetazolamide, dichlorphenamide, and methazolamide), osmotic diuretics (e.g., glycerin, isosorbide, mannitol, and urea), diuretics that inhibit  $\text{Na}^+\text{-K}^+\text{-2Cl}^-$  symport (e.g., furosemide, bumetanide, azosemide, piretanide, triparamide, ethacrynic acid, muzolimine, and torsemide), thiazide and thiazide-like diuretics (e.g., bendroflumethiazide, benzthiazide, chlorothiazide, hydrochlorothiazide, hydroflumethiazide, methyclothiazide, polythiazide, trichormethiazide, chlorthalidone, indapamide, metolazone, and quinethazone), potassium sparing diuretics (e.g., amiloride and triamterene), and mineralcorticoid receptor antagonists (e.g., spironolactone, canrenone, and potassium canrenoate).

[001006] In one embodiment, the Therapeutics of the invention are administered in combination with treatments for endocrine and/or hormone imbalance disorders. Treatments for endocrine and/or hormone imbalance disorders include, but are not limited to,  $^{127}\text{I}$ , radioactive isotopes of iodine such as  $^{131}\text{I}$  and  $^{123}\text{I}$ ; recombinant growth hormone, such as HUMATROPE™ (recombinant somatropin); growth hormone analogs such as PROTROPIN™ (somatrem); dopamine agonists such as



PARLODEL™ (bromocriptine); somatostatin analogs such as SANDOSTATIN™ (octreotide); gonadotropin preparations such as PREGNYL™, A.P.L.™ and PROFASI™ (chorionic gonadotropin (CG)), PERGONAL™ (menotropins), and METRODIN™ (urofollitropin (uFSH)); synthetic human gonadotropin releasing hormone preparations such as FACTREL™ and LUTREPULSE™ (gonadorelin hydrochloride); synthetic gonadotropin agonists such as LUPRON™ (leuprolide acetate), SUPPRELIN™ (histrelin acetate), SYNAREL™ (nafarelin acetate), and ZOLADEX™ (goserelin acetate); synthetic preparations of thyrotropin-releasing hormone such as RELEFACT TRH™ and THYPINONE™ (protirelin); recombinant human TSH such as THYROGEN™; synthetic preparations of the sodium salts of the natural isomers of thyroid hormones such as L-T<sub>4</sub>™, SYNTHROID™ and LEVOTHROID™ (levothyroxine sodium), L-T<sub>3</sub>™, CYTOMEL™ and TRIOSTAT™ (liothyroine sodium), and THYROLAR™ (liotrix); antithyroid compounds such as 6-*n*-propylthiouracil (propylthiouracil), 1-methyl-2-mercaptoimidazole and TAPAZOLE™ (methimazole), NEO-MERCAZOLE™ (carbimazole); beta-adrenergic receptor antagonists such as propranolol and esmolol; Ca<sup>2+</sup> channel blockers; dexamethasone and iodinated radiological contrast agents such as TELEPAQUE™ (iopanoic acid) and ORAGRAFIN™ (sodium ipodate); estrogens or conjugated estrogens such as ESTRACE™ (estradiol), ESTINYL™ (ethinyl estradiol), PREMARIN™, ESTRATAB™, ORTHO-EST™, OGEN™ and estropipate (estrone), ESTROVIS™ (quinestrol), ESTRADERM™ (estradiol), DELESTROGEN™ and VALERGEN™ (estradiol valerate), DEPO-ESTRADIOL CYPIONATE™ and ESTROJECT LA™ (estradiol cypionate); antiestrogens such as NOLVADEX™ (tamoxifen), SEROPHENE™ and CLOMID™ (clomiphene); progestins such as DURALUTIN™ (hydroxyprogesterone caproate), MPA™ and DEPO-PROVERA™ (medroxyprogesterone acetate), PROVERA™ and CYCRIN™ (MPA), MEGACE™ (megestrol acetate), NORLUTIN™ (norethindrone), and NORLUTATE™ and AYGESTIN™ (norethindrone acetate); progesterone implants such as NORPLANT SYSTEM™ (subdermal implants of norgestrel); antiprogestins such as RU 486™ (mifepristone); hormonal contraceptives such as ENOVID™ (norethynodrel plus mestranol), PROGESTASERT™ (intrauterine device that releases progesterone), LOESTRIN™,

BREVICON™, MODICON™, GENORA™, NELONA™, NORINYL™, OVACON-35™ and OVACON-50™ (ethinyl estradiol/norethindrone), LEVLEN™, NORDETTE™, TRI-LEVLEN™ and TRIPHASIL-21™ (ethinyl estradiol/levonorgestrel) LO/OVRAL™ and OVRAL™ (ethinyl estradiol/norgestrel), DEMULEN™ (ethinyl estradiol/ethynodiol diacetate), NORINYL™, ORTHO-NOVUM™, NORETHIN™, GENORA™, and NELOVA™ (norethindrone/mestranol), DESOGEN™ and ORTHO-CEPT™ (ethinyl estradiol/desogestrel), ORTHO-CYCLEN™ and ORTHO-TRICYCLEN™ (ethinyl estradiol/norgestimate), MICRONOR™ and NOR-QD™ (norethindrone), and OVRETTE™ (norgestrel); testosterone esters such as methenolone acetate and testosterone undecanoate; parenteral and oral androgens such as TESTOJECT-50™ (testosterone), TESTEX™ (testosterone propionate), DELATESTRYL™ (testosterone enanthate), DEPO-TESTOSTERONE™ (testosterone cypionate), DANOCRINE™ (danazol), HALOTESTIN™ (fluoxymesterone), ORETON METHYL™, TESTRED™ and VIRILON™ (methyltestosterone), and OXANDRIN™ (oxandrolone); testosterone transdermal systems such as TESTODERM™; androgen receptor antagonist and 5-alpha-reductase inhibitors such as ANDROCUR™ (cyproterone acetate), EULEXIN™ (flutamide), and PROSCAR™ (finasteride); adrenocorticotrophic hormone preparations such as CORTROSYN™ (cosyntropin); adrenocortical steroids and their synthetic analogs such as ACLOVATE™ (alclometasone dipropionate), CYCLOCORT™ (amcinonide), BECLOVENT™ and VANCERIL™ (beclomethasone dipropionate), CELESTONE™ (betamethasone), BENISONE™ and UTICORT™ (betamethasone benzoate), DIPROSONE™ (betamethasone dipropionate), CELESTONE PHOSPHATE™ (betamethasone sodium phosphate), CELESTONE SOLUSPAN™ (betamethasone sodium phosphate and acetate), BETA-VAL™ and VALISONE™ (betamethasone valerate), TEMOVATE™ (clobetasol propionate), CLODERM™ (clocortolone pivalate), CORTEF™ and HYDROCORTONE™ (cortisol (hydrocortisone)), HYDROCORTONE ACETATE™ (cortisol (hydrocortisone) acetate), LOCOID™ (cortisol (hydrocortisone) butyrate), HYDROCORTONE PHOSPHATE™ (cortisol (hydrocortisone) sodium phosphate), A-HYDROCORT™ and SOLU CORTEF™ (cortisol (hydrocortisone) sodium succinate), WESTCORT™

(cortisol (hydrocortisone) valerate), CORTISONE ACETATE™ (cortisone acetate), DESOWEN™ and TRIDESILON™ (desonide), TOPICORT™ (desoximetasone), DECADRON™ (dexamethasone), DECADRON LA™ (dexamethasone acetate), DECADRON PHOSPHATE™ and HEXADROL PHOSPHATE™ (dexamethasone sodium phosphate), FLORONE™ and MAXIFLOR™ (diflorasone diacetate), FLORINEF ACETATE™ (fludrocortisone acetate), AEROBID™ and NASALIDE™ (flunisolide), FLUONID™ and SYNALAR™ (fluocinolone acetonide), LIDEX™ (fluocinonide), FLUOR-OP™ and FML™ (fluorometholone), CORDRAN™ (flurandrenolide), HALOG™ (halcinonide), HMS LIZUIFILM™ (medrysone), MEDROL™ (methylprednisolone), DEPO-MEDROL™ and MEDROL ACETATE™ (methylprednisone acetate), A-METHAPRED™ and SOLUMEDROL™ (methylprednisolone sodium succinate), ELOCON™ (mometasone furoate), HALDRONE™ (paramethasone acetate), DELTA-CORTEF™ (prednisolone), ECONOPRED™ (prednisolone acetate), HYDELTRASOL™ (prednisolone sodium phosphate), HYDELTRA-T.B.A™ (prednisolone tebutate), DELTASONE™ (prednisone), ARISTOCORT™ and KENACORT™ (triamcinolone), KENALOG™ (triamcinolone acetonide), ARISTOCORT™ and KENACORT DIACETATE™ (triamcinolone diacetate), and ARISTOSPAN™ (triamcinolone hexacetonide); inhibitors of biosynthesis and action of adrenocortical steroids such as CYTADREN™ (aminoglutethimide), NIZORAL™ (ketoconazole), MODRASTANE™ (trilostane), and METOPIRONE™ (metyrapone); bovine, porcine or human insulin or mixtures thereof; insulin analogs; recombinant human insulin such as HUMULIN™ and NOVOLIN™; oral hypoglycemic agents such as ORAMIDE™ and ORINASE™ (tolbutamide), DIABINESE™ (chlorpropamide), TOLAMIDE™ and TOLINASE™ (tolazamide), DYMELOR™ (acetohexamide), glibenclamide, MICRONASE™, DIBETA™ and GLYNASE™ (glyburide), GLUCOTROL™ (glipizide), and DIAMICRON™ (gliclazide), GLUCOPHAGE™ (metformin), ciglitazone, pioglitazone, and alpha-glucosidase inhibitors; bovine or porcine glucagon; somatostatins such as SANDOSTATIN™ (octreotide); and diazoxides such as PROGLYCEM™ (diazoxide).

[001007] In one embodiment, the Therapeutics of the invention are administered in combination with treatments for uterine motility disorders. Treatments for uterine motility disorders include, but are not limited to, estrogen drugs such as conjugated estrogens (e.g., PREMARIN® and ESTRATAB®), estradiols (e.g., CLIMARA® and ALORA®), estropipate, and chlorotrianisene; progestin drugs (e.g., AMEN® (medroxyprogesterone), MICRONOR® (norethidrone acetate), PROMETRIUM® progesterone, and megestrol acetate); and estrogen/progesterone combination therapies such as, for example, conjugated estrogens/medroxyprogesterone (e.g., PREMPRO™ and PREMPHASE®) and norethindrone acetate/ethinyl estradiol (e.g., FEMHRT™).

[001008] In an additional embodiment, the Therapeutics of the invention are administered in combination with drugs effective in treating iron deficiency and hypochromic anemias, including but not limited to, ferrous sulfate (iron sulfate, FEOSOL™), ferrous fumarate (e.g., FEOSTAT™), ferrous gluconate (e.g., FERGON™), polysaccharide-iron complex (e.g., NIFEREX™), iron dextran injection (e.g., INFED™), cupric sulfate, pyroxidine, riboflavin, Vitamin B<sub>12</sub>, cyanocobalamin injection (e.g., REDISOL™, RUBRAMIN PCT™), hydroxocobalamin, folic acid (e.g., FOLVITE™), leucovorin (folinic acid, 5-CHOH4PteGlu, citrovorum factor) or WELLCOVORIN (Calcium salt of leucovorin), transferrin or ferritin.

[001009] In certain embodiments, the Therapeutics of the invention are administered in combination with agents used to treat psychiatric disorders. Psychiatric drugs that may be administered with the Therapeutics of the invention include, but are not limited to, antipsychotic agents (e.g., chlorpromazine, chlorprothixene, clozapine, fluphenazine, haloperidol, loxapine, mesoridazine, molindone, olanzapine, perphenazine, pimozide, quetiapine, risperidone, thioridazine, thiothixene, trifluoperazine, and triflupromazine), antimanic agents (e.g., carbamazepine, divalproex sodium, lithium carbonate, and lithium citrate), antidepressants (e.g., amitriptyline, amoxapine, bupropion, citalopram, clomipramine, desipramine, doxepin, fluvoxamine, fluoxetine, imipramine, isocarboxazid, maprotiline, mirtazapine, nefazodone, nortriptyline, paroxetine, phenelzine, protriptyline, sertraline, tranylcypromine, trazodone, trimipramine, and venlafaxine), antianxiety agents (e.g., alprazolam, buspirone, chlordiazepoxide, clorazepate, diazepam,

halazepam, lorazepam, oxazepam, and prazepam), and stimulants (e.g., d-amphetamine, methylphenidate, and pemoline).

[001010] In other embodiments, the Therapeutics of the invention are administered in combination with agents used to treat neurological disorders. Neurological agents that may be administered with the Therapeutics of the invention include, but are not limited to, antiepileptic agents (e.g., carbamazepine, clonazepam, ethosuximide, phenobarbital, phenytoin, primidone, valproic acid, divalproex sodium, felbamate, gabapentin, lamotrigine, levetiracetam, oxcarbazepine, tiagabine, topiramate, zonisamide, diazepam, lorazepam, and clonazepam), antiparkinsonian agents (e.g., levodopa/carbidopa, selegiline, amantidine, bromocriptine, pergolide, ropinirole, pramipexole, benzotropine; biperiden; ethopropazine; procyclidine; trihexyphenidyl, tolcapone), and ALS therapeutics (e.g. riluzole).

[001011] In another embodiment, Therapeutics of the invention are administered in combination with vasodilating agents and/or calcium channel blocking agents. Vasodilating agents that may be administered with the Therapeutics of the invention include, but are not limited to, Angiotensin Converting Enzyme (ACE) inhibitors (e.g., papaverine, isoxsuprine, benazepril, captopril, cilazapril, enalapril, enalaprilat, fosinopril, lisinopril, moexipril, perindopril, quinapril, ramipril, spirapril,trandolapril, and nylidrin), and nitrates (e.g., isosorbide dinitrate, isosorbide mononitrate, and nitroglycerin). Examples of calcium channel blocking agents that may be administered in combination with the Therapeutics of the invention include, but are not limited to amlodipine, bepridil, diltiazem, felodipine, flunarizine, isradipine, nifedipine, nimodipine, and verapamil.

[001012] In additional embodiments, the Therapeutics of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

#### ***Example 14: Method of Treating Decreased Levels of the Polypeptide***

[001013] The present invention relates to a method for treating an individual in need of an increased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically

effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a polypeptide of the present invention in an individual can be treated by administering the agonist or antagonist of the present invention. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the agonist or antagonist to increase the activity level of the polypeptide in such an individual.

[001014] For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the agonist or antagonist for six consecutive days. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 13.

#### ***Example 15: Method of Treating Increased Levels of the Polypeptide***

[001015] The present invention also relates to a method of treating an individual in need of a decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

[001016] In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, due to a variety of etiologies, such as cancer.

[001017] For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 13.

#### ***Example 16: Method of Treatment Using Gene Therapy-Ex Vivo***

[001018] One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a

subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

[001019] At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

[001020] pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

[001021] The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

[001022] The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

10094504.030702

[001023] Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

[001024] The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

***Example 17: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention***

[001025] Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

[001026] Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends.



Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

[001027] The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel, then purified by phenol extraction and ethanol precipitation.

[001028] In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

[001029] Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

[001030] Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na<sub>2</sub> HPO<sub>4</sub>, 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3X10<sup>6</sup> cells/ml. Electroporation should be performed immediately following resuspension.

20200404030702

[001031] Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

[001032] Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120 µg/ml. 0.5 ml of the cell suspension (containing approximately  $1.5 \times 10^6$  cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 µF and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

[001033] Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

[001034] The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a patient as described above.

**Example 18: Method of Treatment Using Gene Therapy - In Vivo**

[001035] Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to (i.e., associated with) a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata et al., Cardiovasc. Res. 35(3):470-479 (1997); Chao et al., Pharmacol. Res. 35(6):517-522 (1997); Wolff, Neuromuscul. Disord. 7(5):314-318 (1997); Schwartz et al., Gene Ther. 3(5):405-411 (1996); Tsurumi et al., Circulation 94(12):3281-3290 (1996) (incorporated herein by reference).

[001036] The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

[001037] The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) Ann. NY Acad. Sci. 772:126-139 and Abdallah B. et al. (1995) Biol. Cell 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

[001038] The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene

therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

[001039] The polynucleotide construct can be delivered to the interstitial space of tissues within an animal, including muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

[001040] For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial

tissues, throat or mucous membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

[001041] The dose response effects of injected polynucleotide in muscle *in vivo* is determined as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various amounts of the template DNA.

[001042] Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

[001043] After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in mice can be used to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

### ***Example 19: Transgenic Animals***

[001044] The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human

primates, *e.g.*, baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides of the invention in humans, as part of a gene therapy protocol.

[001045] Any technique known in the art may be used to introduce the transgene (*i.e.*, polynucleotides of the invention) into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, *e.g.*, Ulmer et al., Science 259:1745 (1993); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

[001046] Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

[001047] The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The

regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example, the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

**[001048]** Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

**[001049]** Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the

transgene on a distinct background that is appropriate for an experimental model of interest.

- [001050] Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

### *Example 20: Knock-Out Animals*

- [001051] Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (See e.g., Smithies et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety.) For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (e.g., see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

- [001052] In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (e.g., knockouts) are



administered to a patient *in vivo*. Such cells may be obtained from the patient (i.e., animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (e.g., lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, e.g., by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the patient systemically, e.g., in the circulation, or intraperitoneally.

[001053] Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

[001054] When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

[001055] Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions

and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

***Example 21: Assays Detecting Stimulation or Inhibition of B cell  
Proliferation and Differentiation***

[001056] Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can, in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, tolerance and death among B cell populations.

[001057] One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective ligands CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

[001058] *In vitro* Assay- Agonists or antagonists of the invention can be assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the agonists or antagonists of the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed *Staphylococcus aureus* Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as

IL-2 and IL-15 synergize with SAC and IgM crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation. Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

[001059] Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added  $10^5$  B-cells suspended in culture medium (RPMI 1640 containing 10% FBS,  $5 \times 10^{-5}$ M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and  $10^{-5}$  dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The positive and negative controls are IL2 and medium respectively.

[001060] In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or 2 mg/Kg of agonists or antagonists of the invention, or truncated forms thereof. Mice receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with agonists or antagonists of the invention identify the results of the activity of the agonists or antagonists on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths, and/or significant increases in the nucleated cellularity of the red pulp regions, which may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell zones that infiltrate established T-cell regions.

[001061] Flow cytometric analyses of the spleens from mice treated with agonist or antagonist is used to indicate whether the agonists or antagonists specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice.

[001062] Likewise, a predicted consequence of increased mature B-cell representation in vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and agonists or antagonists-treated mice.

[001063] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

### *Example 22: T Cell Proliferation Assay*

[001064] A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of  $^3\text{H}$ -thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100  $\mu\text{l}$ /well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1  $\mu\text{g}/\text{ml}$  in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells ( $5 \times 10^4$ /well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of agonists or antagonists of the invention (total volume 200  $\mu\text{l}$ ). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100  $\mu\text{l}$  of supernatant is removed and stored -20 degrees C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100  $\mu\text{l}$  of medium containing 0.5  $\mu\text{Ci}$  of  $^3\text{H}$ -thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of  $^3\text{H}$ -thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation of T cells is used as the negative control for the effects of agonists or antagonists of the invention.

[001065] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

*Example 23: Effect of Agonists or Antagonists of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells*

[001066] Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and MHC class II antigens). Treatment with activating factors, such as TNF- $\alpha$ , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC $\gamma$ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

[001067] FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of agonist or antagonist of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

[001068] Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells ( $10^6$ /ml) are treated with increasing concentrations of agonists or antagonists of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for IL-12 content using commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

[001069] Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increased expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

[001070] FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of agonists or antagonists of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

[001071] Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. Agonists or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

[001072] Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated processes (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in polypropylene

tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of  $2 \times 10^6$ /ml in PBS containing PI at a final concentration of 5  $\mu$ g/ml, and then incubated at room temperature for 5 minutes before FACScan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

[001073] Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of  $5 \times 10^5$  cells/ml with increasing concentrations of agonists or antagonists of the invention and under the same conditions, but in the absence of agonists or antagonists. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in the presence of agonist or antagonist of the invention. LPS (10 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e.g., R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

[001074] Oxidative burst. Purified monocytes are plated in 96-w plate at  $2 \times 10^5$  cell/well. Increasing concentrations of agonists or antagonists of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of HRP) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20  $\mu$ l 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of  $H_2O_2$  produced by the macrophages, a standard curve of a  $H_2O_2$  solution of known molarity is performed for each experiment.

[001075] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

#### ***Example 24: Biological Effects of Agonists or Antagonists of the Invention***

##### Astrocyte and Neuronal Assays

[001076] Agonists or antagonists of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate an agonist or antagonist of the invention's activity on these cells.

[001077] Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of an agonist or antagonist of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

##### Fibroblast and endothelial cell assays



10091504.030702

[001078] Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE<sub>2</sub> assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or agonists or antagonists of the invention with or without IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for PGE<sub>2</sub> by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without agonists or antagonists of the invention IL-1 $\alpha$  for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

[001079] Human lung fibroblasts are cultured with FGF-2 or agonists or antagonists of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with agonists or antagonists of the invention.

#### Parkinson Models.

[001080] The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP<sup>+</sup>) and released. Subsequently, MPP<sup>+</sup> is actively accumulated in dopaminergic neurons by the

high-affinity reuptake transporter for dopamine. MPP<sup>+</sup> is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

[001081] It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989). Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

[001082] Based on the data with FGF-2, agonists or antagonists of the invention can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of an agonist or antagonist of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm<sup>2</sup> on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

[001083] Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if an agonist or antagonist of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the agonist or antagonist may be involved in Parkinson's Disease.

[001084] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified

studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

***Example 25: The Effect of Agonists or Antagonists of the Invention on the Growth of Vascular Endothelial Cells***

[001085] On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at  $2 \times 10^4$  cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. An agonist or antagonist of the invention, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

[001086] An increase in the number of HUVEC cells indicates that the compound of the invention may proliferate vascular endothelial cells, while a decrease in the number of HUVEC cells indicates that the compound of the invention inhibits vascular endothelial cells.

[001087] The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

***Example 26: Rat Corneal Wound Healing Model***

[001088] This animal model shows the effect of an agonist or antagonist of the invention on neovascularization. The experimental protocol includes:

Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.

Inserting a spatula below the lip of the incision facing the outer corner of the eye.

Making a pocket (its base is 1-1.5 mm from the edge of the eye).

Positioning a pellet, containing 50ng- 5ug of an agonist or antagonist of the invention, within the pocket.

Treatment with an agonist or antagonist of the invention can also be applied

topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five days).

[001089] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

### ***Example 27: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models***

#### *Diabetic db+/db+ Mouse Model.*

[001090] To demonstrate that an agonist or antagonist of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

[001091] The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman, D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is

resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

[001092] The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.* 136:1235-1246 (1990)).

[001093] Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

[001094] Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

[001095] Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

[001096] An agonist or antagonist of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

[001097] Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

[001098] Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

[001099] Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:

$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

[001100] Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with an agonist or antagonist of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated lens micrometer is used by a blinded observer.

[001101] Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated lens micrometer.

[001102] Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer served as a positive tissue control and human brain tissue is used as a negative tissue control. Each specimen included a section with omission of the primary antibody and substitution with non-immune mouse IgG. Ranking of these

sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

- [001103] Experimental data are analyzed using an unpaired t test. A p value of  $< 0.05$  is considered significant.

#### *Steroid Impaired Rat Model*

- [001104] The inhibition of wound healing by steroids has been well documented in various *in vitro* and *in vivo* systems (Wahl, Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *Am. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

- [001105] To demonstrate that an agonist or antagonist of the invention can accelerate the healing process, the effects of multiple topical applications of the agonist or antagonist on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

- [001106] Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc.

Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

[001107] The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

[001108] Wounds are visually examined and photographed at a fixed distance at the day of wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

[001109] The agonist or antagonist of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

[001110] Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

[001111] Three groups of 10 animals each (5 with methylprednisolone and 5 without glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated groups.

[001112] Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm<sup>2</sup>, the corresponding size of the dermal punch. Calculations are made using the following formula:



[Open area on day 8] - [Open area on day 1] / [Open area on day 1]

[001113] Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds. Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with an agonist or antagonist of the invention. A calibrated lens micrometer is used by a blinded observer to determine the distance of the wound gap.

[001114] Experimental data are analyzed using an unpaired t test. A p value of  $< 0.05$  is considered significant.

[001115] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

### *Example 28: Lymphadema Animal Model*

[001116] The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of an agonist or antagonist of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

[001117] Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's

Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

[001118] Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated or suture ligated.

[001119] Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

[001120] Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck). The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

[001121] To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect of plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places. Analysis is performed in a blind manner.

[001122] Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people and those 2 readings are averaged. Readings are taken from both control and edematous limbs.

[001123] Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), and both legs are shaved and equally marked using waterproof marker on legs. Legs are first

dipped in water, then dipped into instrument to each marked level, then measured by Buxco edema software(Chen/Victor). Data is recorded by one person, while the other is dipping the limb to marked area.

[001124] Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and  $\text{Ca}^{2+}$  comparison.

[001125] Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.

[001126] Histological Preparations: The transverse muscle located behind the knee (popliteal) area is dissected and arranged in a metal mold, filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

[001127] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

***Example 29: Suppression of TNF alpha-induced adhesion molecule expression by a Agonist or Antagonist of the Invention***

[001128] The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

[001129] Tumor necrosis factor alpha (TNF-a), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

[001130] The potential of an agonist or antagonist of the invention to mediate a suppression of TNF-a induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF-a treated ECs when co-stimulated with a member of the FGF family of proteins.

[001131] To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics, San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a 37 degree C humidified incubator containing 5% CO<sub>2</sub>. HUVECs are seeded in 96-well plates at concentrations of  $1 \times 10^4$  cells/well in EGM medium at 37 degree C for 18-24 hrs or until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

[001132] Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 ul of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 ul volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100  $\mu$ l of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min.

[001133] Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10  $\mu$ l of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10  $\mu$ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

[001134] Then add 20  $\mu$ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with

PBS(+Ca,Mg)+0.5% BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100  $\mu$ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 ( $10^0$ ) >  $10^{-0.5}$  >  $10^{-1}$  >  $10^{-1.5}$ . 5  $\mu$ l of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100  $\mu$ l of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50  $\mu$ l of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [ 5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

[001135] The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

### ***Example 30: Production Of Polypeptide of the Invention For High-Throughput Screening Assays***

[001136] The following protocol produces a supernatant containing polypeptide of the present invention to be tested. This supernatant can then be used in the Screening Assays described in Examples 32-41.

[001137] First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200  $\mu$ l of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

[001138] Plate 293T cells (do not carry cells past P+20) at  $2 \times 10^5$  cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

[001139] The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8-10, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

[001140] Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a 12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degree C for 6 hours.

[001141] While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or HGS CHO-5 media (116.6 mg/L of  $\text{CaCl}_2$  (anhyd); 0.00130 mg/L  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ ; 0.050 mg/L of  $\text{Fe}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ ; 0.417 mg/L of  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ; 311.80 mg/L of Kcl; 28.64 mg/L of  $\text{MgCl}_2$ ; 48.84 mg/L of  $\text{MgSO}_4$ ; 6995.50 mg/L of NaCl; 2400.0 mg/L of  $\text{NaHCO}_3$ ; 62.50 mg/L of  $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ ; 71.02 mg/L of  $\text{Na}_2\text{HPO}_4$ ; .4320 mg/L of  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ ; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine;

20091504-030702

147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H<sub>2</sub>O; 6.65 mg/ml of L-Aspartic Acid; 29.56 mg/ml of L-Cystine-2HCL-H<sub>2</sub>O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H<sub>2</sub>O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H<sub>2</sub>O; and 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; 0.680 mg/L of Vitamin B<sub>12</sub>; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic Acid; 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal Acetate. Adjust osmolarity to 327 mOsm) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

[001142] The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degree C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

[001143] On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 32-39.

[001144] It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the

polypeptide of the present invention directly (e.g., as a secreted protein) or by polypeptide of the present invention inducing expression of other proteins, which are then secreted into the supernatant. Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

### *Example 31: Construction of GAS Reporter Construct*

[001145] One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

[001146] GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class I, cells after treatment with IL-12. Stat5 was originally called mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

[001147] The STATs are activated to translocate from the cytoplasm to the nucleus upon tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

[001148] The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g, and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and



one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xaa-Trp-Ser (SEQ ID NO:2)).

[001149] Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is encompassed in the Jaks-STATs signal transduction pathway.

[001150] Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter molecules, activators of the Jaks-STATs pathway can be identified.

20090409T0001

<u>JAKs</u>	<u>STATs</u>	<u>GAS(elements) or ISRE</u>				
<u>Ligand</u>	<u>tyk2</u>	<u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>		
<u>IFN family</u>						
IFN-a/B	+	+	-	-	1,2,3	ISRE
IFN-g		+	+	-	1	GAS (IRF1>Lys6>IFP)
Il-10	+	?	?	-	1,3	
<u>gp130 family</u>						
IL-6 (Pleiotropic) +	+	+	?		1,3	GAS (IRF1>Lys6>IFP)
Il-11(Pleiotropic) ?	+	?	?		1,3	
OnM(Pleiotropic) ?	+	+	?		1,3	
LIF(Pleiotropic) ?	+	+	?		1,3	
CNTF(Pleiotropic)	-/+	+	+	?	1,3	
G-CSF(Pleiotropic)	?	+	?	?	1,3	
IL-12(Pleiotropic)	+	-	+	+	1,3	
<u>g-C family</u>						
IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 =IFP >>Ly6)(IgH)
IL-7 (lymphocytes)	-	+	-	+	5	GAS
IL-9 (lymphocytes)	-	+	-	+	5	GAS
IL-13 (lymphocyte)	-	+	?	?	6	GAS
IL-15	?	+	?	+	5	GAS
<u>gp140 family</u>						
IL-3 (myeloid)	-	-	+	-	5	GAS (IRF1>IFP>>Ly6)
IL-5 (myeloid)	-	-	+	-	5	GAS
GM-CSF (myeloid)	-	-	+	-	5	GAS
<u>Growth hormone family</u>						
GH	?	-	+	-	5	
PRL	?	+/-	+	-	1,3,5	
EPO	?	-	+	-	5	GAS(B-CAS>IRF1=IFP>>Ly6)
<u>Receptor Tyrosine Kinases</u>						
EGF	?	+	+	-	1,3	GAS (IRF1)
PDGF	?	+	+	-	1,3	
CSF-1	?	+	+	-	1,3	GAS (not IRF1)

[001151] To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 32-33, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5':GCGCCTCGAGATTTCCCGAAATCTAGATTTCCCGAAATGATTTCCCG  
GAAATGATTTCCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO: 3)

[001152] The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO: 4)

[001153] PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5':CTCGAGATTTCCCGAAATCTAGATTTCCCGAAATGATTTCCCGAAA  
TGATTTCCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCG  
CCCCTAACTCCGCCCATCCCGCCCCTAACTCCGCCCAGTTCGCCCATTTCT  
CCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCC  
TCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTA  
GGCTTTTGGCAAAAAAGCTT:3' (SEQ ID NO: 5)

[001154] With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be used instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

10091504-030302

[001155] The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

[001156] Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 32-33.

[001157] Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 34 and 35. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

***Example 32: High-Throughput Screening Assay for T-cell Activity.***

[001158] The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 31. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152),

although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC Accession No. CRL-1582) cells can also be used.

[001159] Jurkat T-cells are lymphoblastic CD4+ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

[001160] Specifically, the following protocol will yield sufficient cells for 75 wells containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

[001161] During the incubation period, count cell concentration, spin down the required number of cells ( $10^7$  per transfection), and resuspend in OPTI-MEM to a final concentration of  $10^7$  cells/ml. Then add 1ml of  $1 \times 10^7$  cells in OPTI-MEM to T25 flask and incubate at 37 degree C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

[001162] The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing polypeptide of the present invention or polypeptide of the present invention induced polypeptides as produced by the protocol described in Example 30.

[001163] On the day of treatment with the supernatant, the cells should be washed and resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

[001164] Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100, 000 cells per well).

[001165] After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

[001166] The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and stored at -20 degree C until SEAP assays are performed according to Example 36. The plates containing the remaining treated cells are placed at 4 degree C and serve as a source of material for repeating the assay on a specific well if desired.

[001167] As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

[001168] The above protocol may be used in the generation of both transient, as well as stable, transfected cells, which would be apparent to those of skill in the art.

### ***Example 33: High-Throughput Screening Assay Identifying Myeloid Activity***

[001169] The following protocol is used to assess myeloid activity of polypeptide of the present invention by determining whether polypeptide of the present invention proliferates and/or differentiates myeloid cells. Myeloid cell activity is assessed using the GAS/SEAP/Neo construct produced in Example 31. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

2009-03-04 10:57:50

[001170] To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 31, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest  $2 \times 10^7$  U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 mg/ml streptomycin.

[001171] Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM  $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ , 1 mM  $\text{MgCl}_2$ , and 675 uM  $\text{CaCl}_2$ . Incubate at 37 degrees C for 45 min.

[001172] Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degree C for 36 hr.

[001173] The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

[001174] These cells are tested by harvesting  $1 \times 10^8$  cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of  $5 \times 10^5$  cells/ml. Plate 200 ul cells per well in the 96-well plate (or  $1 \times 10^5$  cells/well).

[001175] Add 50 ul of the supernatant prepared by the protocol described in Example 30. Incubate at 37 degree C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 36.

### ***Example 34: High-Throughput Screening Assay Identifying Neuronal Activity.***

[001176] When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types

upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed by polypeptide of the present invention.

[001177] Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat phenochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells by polypeptide of the present invention can be assessed.

[001178] The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO: 6)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO: 7)

[001179] Using the GAS:SEAP/Neo vector produced in Example 31, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

[001180] To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

[001181] PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.



20091504-030702

[001182] Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 30. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

[001183] To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

[001184] The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as  $5 \times 10^5$  cells/ml.

[001185] Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to  $1 \times 10^5$  cells/well). Add 50 ul supernatant produced by Example 30, 37 degree C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 36.

### ***Example 35: High-Throughput Screening Assay for T-cell Activity***

[001186] NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF, CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

[001187] In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded,

causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.

[001188] Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 30. Activators or inhibitors of NF-KB would be useful in treating, preventing, and/or diagnosing diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

[001189] To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO: 8), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC  
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO: 9)

[001190] The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTTGCAAAGCCTAGGC:3' (SEQ ID NO: 4)

[001191] PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene) Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGGACTTTCC  
ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCC  
ATCCCGCCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGA  
CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTA  
TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA  
GCTT:3' (SEQ ID NO: 10)

[001192] Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and

HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

[001193] In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP cassette is removed from the above NF-KB/SEAP vector using restriction enzymes Sall and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with Sall and NotI.

[001194] Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are created and maintained according to the protocol described in Example 32. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 32. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

### *Example 36: Assay for SEAP Activity*

[001195] As a reporter molecule for the assays described in Examples 32-35, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution, Assay, and Reaction Buffers used below.

[001196] Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

[001197] Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the Table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it takes about 10 minutes to read 5 plates on a luminometer, thus one should treat 5 plates at each time and start the second set 10 minutes later.

[001198] Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

[001199] Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5
25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25

44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

---

***Example 37: High-Throughput Screening Assay Identifying Changes in  
Small Molecule Concentration and Membrane Permeability***

**[001200]** Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

**[001201]** The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule, fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

**[001202]** For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO<sub>2</sub> incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

**[001203]** A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO<sub>2</sub> incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

**[001204]** For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to  $2-5 \times 10^6$  cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml

fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to  $1 \times 10^6$  cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley Cell Wash with 200 ul, followed by an aspiration step to 100 ul final volume.

[001205] For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

[001206] To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm; and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event caused by the a molecule, either polypeptide of the present invention or a molecule induced by polypeptide of the present invention, which has resulted in an increase in the intracellular  $\text{Ca}^{++}$  concentration.

### *Example 38: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity*

[001207] The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine Kinase (RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

[001208] Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and

non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family, members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

[001209] Because of the wide range of known factors capable of stimulating tyrosine kinase activity, identifying whether polypeptide of the present invention or a molecule induced by polypeptide of the present invention is capable of activating tyrosine kinase signal transduction pathways is of interest. Therefore, the following protocol is designed to identify such molecules capable of activating the tyrosine kinase signal transduction pathways.

[001210] Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with 100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

[001211] To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 30, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na<sub>3</sub>VO<sub>4</sub>, 2 mM Na<sub>4</sub>P<sub>2</sub>O<sub>7</sub> and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN)) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4°C. The plate is then placed in a

vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degree C at 16,000 x g.

[001212] Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

[001213] Generally, the tyrosine kinase activity of a supernatant is evaluated by determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

[001214] The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg<sub>2</sub><sup>+</sup> (5mM ATP/50mM MgCl<sub>2</sub>), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl<sub>2</sub>, 5 mM MnCl<sub>2</sub>, 0.5 mg/ml BSA), then 5ul of Sodium Vanadate (1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degree C for 2 min. Initiate the reaction by adding 10ul of the control enzyme or the filtered supernatant.

[001215] The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

[001216] Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degree C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase (anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degree C for one hour. Wash the well as above.



[001217] Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

***Example 39: High-Throughput Screening Assay Identifying Phosphorylation Activity***

[001218] As a potential alternative and/or complement to the assay of protein tyrosine kinase activity described in Example 38, an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP, Map kinase kinase (MEK), MEK kinase, Src, Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

[001219] Specifically, assay plates are made by coating the wells of a 96-well ELISA plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degree C until use.

[001220] A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants obtained in Example 30 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

10091504-030702  
20/05/04-405T600T

[001221] After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased fluorescent signal over background indicates a phosphorylation by polypeptide of the present invention or a molecule induced by polypeptide of the present invention.

***Example 40: Assay for the Stimulation of Bone Marrow CD34+ Cell Proliferation***

[001222] This assay is based on the ability of human CD34+ to proliferate in the presence of hematopoietic growth factors and evaluates the ability of isolated polypeptides expressed in mammalian cells to stimulate proliferation of CD34+ cells.

[001223] It has been previously shown that most mature precursors will respond to only a single signal. More immature precursors require at least two signals to respond. Therefore, to test the effect of polypeptides on hematopoietic activity of a wide range of progenitor cells, the assay contains a given polypeptide in the presence or absence of other hematopoietic growth factors. Isolated cells are cultured for 5 days in the presence of Stem Cell Factor (SCF) in combination with tested sample. SCF alone has a very limited effect on the proliferation of bone marrow (BM) cells, acting in such conditions only as a "survival" factor. However, combined with any factor exhibiting stimulatory effect on these cells (e.g., IL-3), SCF will cause a synergistic effect. Therefore, if the tested polypeptide has a stimulatory effect on hematopoietic progenitors, such activity can be easily detected. Since normal BM cells have a low level of cycling cells, it is likely that any inhibitory effect of a given polypeptide, or agonists or antagonists thereof, might not be detected. Accordingly, assays for an inhibitory effect on progenitors is preferably tested in cells that are first subjected to

*in vitro* stimulation with SCF+IL+3, and then contacted with the compound that is being evaluated for inhibition of such induced proliferation.

[001224] Briefly, CD34+ cells are isolated using methods known in the art. The cells are thawed and resuspended in medium (QBSF 60 serum-free medium with 1% L-glutamine (500ml) Quality Biological, Inc., Gaithersburg, MD Cat# 160-204-101). After several gentle centrifugation steps at 200 x g, cells are allowed to rest for one hour. The cell count is adjusted to  $2.5 \times 10^5$  cells/ml. During this time, 100  $\mu$ l of sterile water is added to the peripheral wells of a 96-well plate. The cytokines that can be tested with a given polypeptide in this assay is rhSCF (R&D Systems, Minneapolis, MN, Cat# 255-SC) at 50 ng/ml alone and in combination with rhSCF and rhIL-3 (R&D Systems, Minneapolis, MN, Cat# 203-ML) at 30 ng/ml. After one hour, 10  $\mu$ l of prepared cytokines, 50  $\mu$ l of the supernatants prepared in Example 30 (supernatants at 1:2 dilution = 50  $\mu$ l) and 20  $\mu$ l of diluted cells are added to the media which is already present in the wells to allow for a final total volume of 100  $\mu$ l. The plates are then placed in a 37°C/5% CO<sub>2</sub> incubator for five days.

[001225] Eighteen hours before the assay is harvested, 0.5  $\mu$ Ci/well of [3H] Thymidine is added in a 10  $\mu$ l volume to each well to determine the proliferation rate. The experiment is terminated by harvesting the cells from each 96-well plate to a filtermat using the Tomtec Harvester 96. After harvesting, the filtermats are dried, trimmed and placed into OmniFilter assemblies consisting of one OmniFilter plate and one OmniFilter Tray. 60  $\mu$ l Microscint is added to each well and the plate sealed with TopSeal-A press-on sealing film. A bar code 15 sticker is affixed to the first plate for counting. The sealed plates are then loaded and the level of radioactivity determined via the Packard Top Count and the printed data collected for analysis. The level of radioactivity reflects the amount of cell proliferation.

[001226] The studies described in this example test the activity of a given polypeptide to stimulate bone marrow CD34+ cell proliferation. One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof. As a nonlimiting example, potential antagonists tested in this assay would be expected to inhibit cell proliferation in the presence of cytokines and/or to increase the inhibition of cell proliferation in the presence of cytokines and a given polypeptide.

In contrast, potential agonists tested in this assay would be expected to enhance cell proliferation and/or to decrease the inhibition of cell proliferation in the presence of cytokines and a given polypeptide.

- [001227] The ability of a gene to stimulate the proliferation of bone marrow CD34+ cells indicates that polynucleotides and polypeptides corresponding to the gene are useful for the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections above, and elsewhere herein.

***Example 41: Assay for Extracellular Matrix Enhanced Cell Response  
(EMECCR)***

- [001228] The objective of the Extracellular Matrix Enhanced Cell Response (EMECCR) assay is to identify gene products (e.g., isolated polypeptides) that act on the hematopoietic stem cells in the context of the extracellular matrix (ECM) induced signal.

- [001229] Cells respond to the regulatory factors in the context of signal(s) received from the surrounding microenvironment. For example, fibroblasts, and endothelial and epithelial stem cells fail to replicate in the absence of signals from the ECM. Hematopoietic stem cells can undergo self-renewal in the bone marrow, but not in *in vitro* suspension culture. The ability of stem cells to undergo self-renewal *in vitro* is dependent upon their interaction with the stromal cells and the ECM protein fibronectin (fn). Adhesion of cells to fn is mediated by the  $\alpha_5\beta_1$  and  $\alpha_4\beta_1$  integrin receptors, which are expressed by human and mouse hematopoietic stem cells. The factor(s) which integrate with the ECM environment and are responsible for stimulating stem cell self-renewal have not yet been identified. Discovery of such factors should be of great interest in gene therapy and bone marrow transplant applications

40091504-030700

[001230] Briefly, polystyrene, non tissue culture treated, 96-well plates are coated with fn fragment at a coating concentration of  $0.2 \mu\text{g}/\text{cm}^2$ . Mouse bone marrow cells are plated (1,000 cells/well ) in 0.2 ml of serum-free medium. Cells cultured in the presence of IL-3 ( 5 ng/ml ) + SCF ( 50 ng/ml ) would serve as the positive control, conditions under which little self-renewal but pronounced differentiation of the stem cells is to be expected. Gene products of the invention (e.g., including, but not limited to, polynucleotides and polypeptides of the present invention, and supernatants produced in Example 30), are tested with appropriate negative controls in the presence and absence of SCF(5.0 ng/ml), where test factor supernatants represent 10% of the total assay volume. The plated cells are then allowed to grow by incubating in a low oxygen environment ( 5%  $\text{CO}_2$ , 7%  $\text{O}_2$ , and 88%  $\text{N}_2$  ) tissue culture incubator for 7 days. The number of proliferating cells within the wells is then quantitated by measuring thymidine incorporation into cellular DNA. Verification of the positive hits in the assay will require phenotypic characterization of the cells, which can be accomplished by scaling up of the culture system and using appropriate antibody reagents against cell surface antigens and FACScan.

[001231] One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

[001232] If a particular polypeptide of the present invention is found to be a stimulator of hematopoietic progenitors, polynucleotides and polypeptides corresponding to the gene encoding said polypeptide may be useful for the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections above, and elsewhere herein. The gene product may also be useful in the expansion of stem cells and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

[001233] Additionally, the polynucleotides and/or polypeptides of the gene of interest and/or agonists and/or antagonists thereof, may also be employed to inhibit the proliferation and differentiation of hematopoietic cells and therefore may be employed to protect bone marrow stem cells from chemotherapeutic agents during chemotherapy. This antiproliferative effect may allow administration of higher doses

of chemotherapeutic agents and, therefore, more effective chemotherapeutic treatment.

- [001234] Moreover, polynucleotides and polypeptides corresponding to the gene of interest may also be useful for the treatment and diagnosis of hematopoietic related disorders such as, for example, anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of neoplasia.

***Example 42: Human Dermal Fibroblast and Aortic Smooth Muscle Cell Proliferation***

- [001235] The polypeptide of interest is added to cultures of normal human dermal fibroblasts (NHDF) and human aortic smooth muscle cells (AoSMC) and two co-assays are performed with each sample. The first assay examines the effect of the polypeptide of interest on the proliferation of normal human dermal fibroblasts (NHDF) or aortic smooth muscle cells (AoSMC). Aberrant growth of fibroblasts or smooth muscle cells is a part of several pathological processes, including fibrosis, and restenosis. The second assay examines IL6 production by both NHDF and SMC. IL6 production is an indication of functional activation. Activated cells will have increased production of a number of cytokines and other factors, which can result in a proinflammatory or immunomodulatory outcome. Assays are run with and without co-TNF $\alpha$  stimulation, in order to check for costimulatory or inhibitory activity.

- [001236] Briefly, on day 1, 96-well black plates are set up with 1000 cells/well (NHDF) or 2000 cells/well (AoSMC) in 100  $\mu$ l culture media. NHDF culture media contains: Clonetics FB basal media, 1mg/ml hFGF, 5mg/ml insulin, 50mg/ml gentamycin, 2%FBS, while AoSMC culture media contains Clonetics SM basal media, 0.5  $\mu$ g/ml hEGF, 5mg/ml insulin, 1 $\mu$ g/ml hFGF, 50mg/ml gentamycin, 50  $\mu$ g/ml Amphotericin B, 5%FBS. After incubation at 37°C for at least 4-5 hours culture media is aspirated and replaced with growth arrest media. Growth arrest media for NHDF contains

fibroblast basal media, 50mg/ml gentamycin, 2% FBS, while growth arrest media for AoSMC contains SM basal media, 50mg/ml gentamycin, 50µg/ml Amphotericin B, 0.4% FBS. Incubate at 37 °C until day 2.

[001237] On day 2, serial dilutions and templates of the polypeptide of interest are designed such that they always include media controls and known-protein controls. For both stimulation and inhibition experiments, proteins are diluted in growth arrest media. For inhibition experiments, TNFα is added to a final concentration of 2ng/ml (NHDF) or 5ng/ml (AoSMC). Add 1/3 vol media containing controls or polypeptides of the present invention and incubate at 37 degrees C/5% CO<sub>2</sub> until day 5.

[001238] Transfer 60µl from each well to another labeled 96-well plate, cover with a plate-sealer, and store at 4 degrees C until Day 6 (for IL6 ELISA). To the remaining 100 µl in the cell culture plate, aseptically add Alamar Blue in an amount equal to 10% of the culture volume (10µl). Return plates to incubator for 3 to 4 hours. Then measure fluorescence with excitation at 530nm and emission at 590nm using the CytoFluor. This yields the growth stimulation/inhibition data.

[001239] On day 5, the IL6 ELISA is performed by coating a 96 well plate with 50-100 ul/well of Anti-Human IL6 Monoclonal antibody diluted in PBS, pH 7.4, incubate ON at room temperature.

[001240] On day 6, empty the plates into the sink and blot on paper towels. Prepare Assay Buffer containing PBS with 4% BSA. Block the plates with 200 µl/well of Pierce Super Block blocking buffer in PBS for 1-2 hr and then wash plates with wash buffer (PBS, 0.05% Tween-20). Blot plates on paper towels. Then add 50 µl/well of diluted Anti-Human IL-6 Monoclonal, Biotin-labeled antibody at 0.50 mg/ml. Make dilutions of IL-6 stock in media (30, 10, 3, 1, 0.3, 0 ng/ml). Add duplicate samples to top row of plate. Cover the plates and incubate for 2 hours at RT on shaker.

[001241] Plates are washed with wash buffer and blotted on paper towels. Dilute EU-labeled Streptavidin 1:1000 in Assay buffer, and add 100 µl/well. Cover the plate and incubate 1 h at RT. Plates are again washed with wash buffer and blotted on paper towels.

[001242] Add 100 µl/well of Enhancement Solution. Shake for 5 minutes. Read the plate on the Wallac DELFIA Fluorometer. Readings from triplicate samples in each assay were tabulated and averaged.

70094504, 030702

[001243] A positive result in this assay suggests AoSMC cell proliferation and that the polypeptide of the present invention may be involved in dermal fibroblast proliferation and/or smooth muscle cell proliferation. A positive result also suggests many potential uses of polypeptides, polynucleotides, agonists and/or antagonists of the polynucleotide/polypeptide of the present invention which gives a positive result. For example, inflammation and immune responses, wound healing, and angiogenesis, as detailed throughout this specification. Particularly, polypeptides of the present invention and polynucleotides of the present invention may be used in wound healing and dermal regeneration, as well as the promotion of vasculogenesis, both of the blood vessels and lymphatics. The growth of vessels can be used in the treatment of, for example, cardiovascular diseases. Additionally, antagonists of polypeptides and polynucleotides of the invention may be useful in treating diseases, disorders, and/or conditions which involve angiogenesis by acting as an anti-vascular agent (e.g., anti-angiogenesis). These diseases, disorders, and/or conditions are known in the art and/or are described herein, such as, for example, malignancies, solid tumors, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; atherosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis. Moreover, antagonists of polypeptides and polynucleotides of the invention may be useful in treating anti-hyperproliferative diseases and/or anti-inflammatory known in the art and/or described herein.

[001244] One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.



**Example 43: Cellular Adhesion Molecule (CAM) Expression  
on Endothelial Cells**

**[001245]** The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

**[001246]** Briefly, endothelial cells (e.g., Human Umbilical Vein Endothelial cells (HUVECs)) are grown in a standard 96 well plate to confluence, growth medium is removed from the cells and replaced with 100  $\mu$ l of 199 Medium (10% fetal bovine serum (FBS)). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10  $\mu$ l volumes). Plates are then incubated at 37°C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100  $\mu$ l of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min. Fixative is removed from the wells and wells are washed 1X with PBS(+Ca,Mg) + 0.5% BSA and drained. 10  $\mu$ l of diluted primary antibody is added to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10  $\mu$ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed three times with PBS(+Ca,Mg) + 0.5% BSA. 20  $\mu$ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution, referred to herein as the working dilution) are added to each well and incubated at 37°C for 30 min. Wells are washed three times with

PBS(+Ca,Mg)+0.5% BSA. Dissolve 1 tablet of p-Nitrophenol Phosphate pNPP per 5 ml of glycine buffer (pH 10.4). 100  $\mu$ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer:  $1:5,000$  ( $10^0$ )  $> 10^{-0.5} > 10^{-1} > 10^{-1.5}$ . 5  $\mu$ l of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100  $\mu$ l of pNPP reagent is then added to each of the standard wells. The plate is incubated at 37°C for 4h. A volume of 50  $\mu$ l of 3M NaOH is added to all wells. The plate is read on a plate reader at 405 nm using the background subtraction option on blank wells filled with glycine buffer only. Additionally, the template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

#### ***Example 44: Alamar Blue Endothelial Cells Proliferation Assay***

[001247] This assay may be used to quantitatively determine protein mediated inhibition of bFGF-induced proliferation of Bovine Lymphatic Endothelial Cells (LECs), Bovine Aortic Endothelial Cells (BAECs) or Human Microvascular Uterine Myometrial Cells (UTMECs). This assay incorporates a fluorometric growth indicator based on detection of metabolic activity. A standard Alamar Blue Proliferation Assay is prepared in EGM-2MV with 10 ng /ml of bFGF added as a source of endothelial cell stimulation. This assay may be used with a variety of endothelial cells with slight changes in growth medium and cell concentration. Dilutions of the protein batches to be tested are diluted as appropriate. Serum-free medium (GIBCO SFM) without bFGF is used as a non-stimulated control and Angiostatin or TSP-1 are included as a known inhibitory controls.

[001248] Briefly, LEC, BAECs or UTMECs are seeded in growth media at a density of 5000 to 2000 cells/well in a 96 well plate and placed at 37degrees C overnight. After the overnight incubation of the cells, the growth media is removed and replaced with GIBCO EC-SFM. The cells are treated with the appropriate dilutions of the protein of interest or control protein sample(s) (prepared in SFM ) in triplicate wells with additional bFGF to a concentration of 10 ng/ ml. Once the cells have been treated

with the samples, the plate(s) is/are placed back in the 37° C incubator for three days. After three days 10 ml of stock alamar blue (Biosource Cat# DAL1100) is added to each well and the plate(s) is/are placed back in the 37°C incubator for four hours. The plate(s) are then read at 530nm excitation and 590nm emission using the CytoFluor fluorescence reader. Direct output is recorded in relative fluorescence units.

[001249] Alamar blue is an oxidation-reduction indicator that both fluoresces and changes color in response to chemical reduction of growth medium resulting from cell growth. As cells grow in culture, innate metabolic activity results in a chemical reduction of the immediate surrounding environment. Reduction related to growth causes the indicator to change from oxidized (non-fluorescent blue) form to reduced (fluorescent red) form (i.e., stimulated proliferation will produce a stronger signal and inhibited proliferation will produce a weaker signal and the total signal is proportional to the total number of cells as well as their metabolic activity). The background level of activity is observed with the starvation medium alone. This is compared to the output observed from the positive control samples (bFGF in growth medium) and protein dilutions.

#### ***Example 45: Detection of Inhibition of a Mixed Lymphocyte Reaction***

[001250] This assay can be used to detect and evaluate inhibition of a Mixed Lymphocyte Reaction (MLR) by gene products (e.g., isolated polypeptides). Inhibition of a MLR may be due to a direct effect on cell proliferation and viability, modulation of costimulatory molecules on interacting cells, modulation of adhesiveness between lymphocytes and accessory cells, or modulation of cytokine production by accessory cells. Multiple cells may be targeted by these polypeptides since the peripheral blood mononuclear fraction used in this assay includes T, B and natural killer lymphocytes, as well as monocytes and dendritic cells.

[001251] Polypeptides of interest found to inhibit the MLR may find application in diseases associated with lymphocyte and monocyte activation or proliferation. These include, but are not limited to, diseases such as asthma, arthritis, diabetes, inflammatory skin conditions, psoriasis, eczema, systemic lupus erythematosus, multiple sclerosis, glomerulonephritis, inflammatory bowel disease, crohn's disease,

ulcerative colitis, arteriosclerosis, cirrhosis, graft vs. host disease, host vs. graft disease, hepatitis, leukemia and lymphoma.

[001252] Briefly, PBMCs from human donors are purified by density gradient centrifugation using Lymphocyte Separation Medium (LSM<sup>®</sup>, density 1.0770 g/ml, Organon Teknika Corporation, West Chester, PA). PBMCs from two donors are adjusted to  $2 \times 10^6$  cells/ml in RPMI-1640 (Life Technologies, Grand Island, NY) supplemented with 10% FCS and 2 mM glutamine. PBMCs from a third donor is adjusted to  $2 \times 10^5$  cells/ml. Fifty microliters of PBMCs from each donor is added to wells of a 96-well round bottom microtiter plate. Dilutions of test materials (50  $\mu$ l) is added in triplicate to microtiter wells. Test samples (of the protein of interest) are added for final dilution of 1:4; rhuIL-2 (R&D Systems, Minneapolis, MN, catalog number 202-IL) is added to a final concentration of 1  $\mu$ g/ml; anti-CD4 mAb (R&D Systems, clone 34930.11, catalog number MAB379) is added to a final concentration of 10  $\mu$ g/ml. Cells are cultured for 7-8 days at 37°C in 5% CO<sub>2</sub>, and 1  $\mu$ C of [<sup>3</sup>H] thymidine is added to wells for the last 16 hrs of culture. Cells are harvested and thymidine incorporation determined using a Packard TopCount. Data is expressed as the mean and standard deviation of triplicate determinations.

[001253] Samples of the protein of interest are screened in separate experiments and compared to the negative control treatment, anti-CD4 mAb, which inhibits proliferation of lymphocytes and the positive control treatment, IL-2 (either as recombinant material or supernatant), which enhances proliferation of lymphocytes.

[001254] One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

#### *Example 46: Assays for Protease Activity*

[001255] The following assay may be used to assess protease activity of the polypeptides of the invention.

[001256] Gelatin and casein zymography are performed essentially as described (Heusen et al., Anal. Biochem., 102:196-202 (1980); Wilson et al., Journal of

Urology, 149:653-658 (1993)). Samples are run on 10% polyacrylamide/0.1% SDS gels containing 1% gelatin or casein, soaked in 2.5% triton at room temperature for 1 hour, and in 0.1M glycine, pH 8.3 at 37°C 5 to 16 hours. After staining in amido black areas of proteolysis appear as clear areas against the blue-black background. Trypsin (Sigma T8642) is used as a positive control.

[001257] Protease activity is also determined by monitoring the cleavage of n-benzoyl-L-arginine ethyl ester (BAEE) (Sigma B-4500). Reactions are set up in (25mM NaPO<sub>4</sub>, 1mM EDTA, and 1mM BAEE), pH 7.5. Samples are added and the change in absorbance at 260nm is monitored on the Beckman DU-6 spectrophotometer in the time-drive mode. Trypsin is used as a positive control.

[001258] Additional assays based upon the release of acid-soluble peptides from casein or hemoglobin measured as absorbance at 280 nm or colorimetrically using the Folin method are performed as described in Bergmeyer, et al., Methods of Enzymatic Analysis, 5 (1984). Other assays involve the solubilization of chromogenic substrates (Ward, Applied Science, 251-317 (1983)).

#### ***Example 47: Identifying Serine Protease Substrate Specificity***

[001259] Methods known in the art or described herein may be used to determine the substrate specificity of the polypeptides of the present invention having serine protease activity. A preferred method of determining substrate specificity is by the use of positional scanning synthetic combinatorial libraries as described in GB 2 324 529 (incorporated herein in its entirety).

#### ***Example 48: Ligand Binding Assays***

[001260] The following assay may be used to assess ligand binding activity of the polypeptides of the invention.

[001261] Ligand binding assays provide a direct method for ascertaining receptor pharmacology and are adaptable to a high throughput format. The purified ligand for a polypeptide is radiolabeled to high specific activity (50-2000 Ci/mmol) for binding studies. A determination is then made that the process of radiolabeling does not

diminish the activity of the ligand towards its polypeptide. Assay conditions for buffers, ions, pH and other modulators such as nucleotides are optimized to establish a workable signal to noise ratio for both membrane and whole cell polypeptide sources. For these assays, specific polypeptide binding is defined as total associated radioactivity minus the radioactivity measured in the presence of an excess of unlabeled competing ligand. Where possible, more than one competing ligand is used to define residual nonspecific binding.

#### ***Example 49: Functional Assay in *Xenopus* Oocytes***

[001262] Capped RNA transcripts from linearized plasmid templates encoding the polypeptides of the invention are synthesized *in vitro* with RNA polymerases in accordance with standard procedures. *In vitro* transcripts are suspended in water at a final concentration of 0.2 mg/ml. Ovarian lobes are removed from adult female toads, Stage V defolliculated oocytes are obtained, and RNA transcripts (10 ng/oocyte) are injected in a 50 nl bolus using a microinjection apparatus. Two electrode voltage clamps are used to measure the currents from individual *Xenopus oocytes* in response to polypeptides and polypeptide agonist exposure. Recordings are made in Ca<sup>2+</sup> free Barth's medium at room temperature. The *Xenopus* system can be used to screen known ligands and tissue/cell extracts for activating ligands.

#### ***Example 50: Microphysiometric Assays***

[001263] Activation of a wide variety of secondary messenger systems results in extrusion of small amounts of acid from a cell. The acid formed is largely as a result of the increased metabolic activity required to fuel the intracellular signaling process. The pH changes in the media surrounding the cell are very small but are detectable by the CYTOSENSOR microphysiometer (Molecular Devices Ltd., Menlo Park, Calif.). The CYTOSENSOR is thus capable of detecting the activation of polypeptide which is coupled to an energy utilizing intracellular signaling pathway.

### *Example 51: Extract/Cell Supernatant Screening*

[001264] A large number of mammalian receptors exist for which there remains, as yet, no cognate activating ligand (agonist). Thus, active ligands for these receptors may not be included within the ligands banks as identified to date. Accordingly, the polypeptides of the invention can also be functionally screened (using calcium, cAMP, microphysiometer, oocyte electrophysiology, etc., functional screens) against tissue extracts to identify its natural ligands. Extracts that produce positive functional responses can be sequentially subfractionated until an activating ligand is isolated and identified.

### *Example 52: Calcium and cAMP Functional Assays*

[001265] Seven transmembrane receptors which are expressed in HEK 293 cells have been shown to be coupled functionally to activation of PLC and calcium mobilization and/or cAMP stimulation or inhibition. Basal calcium levels in the HEK 293 cells in receptor-transfected or vector control cells were observed to be in the normal, 100 nM to 200 nM, range. HEK 293 cells expressing recombinant receptors are loaded with fura 2 and in a single day >150 selected ligands or tissue/cell extracts are evaluated for agonist induced calcium mobilization. Similarly, HEK 293 cells expressing recombinant receptors are evaluated for the stimulation or inhibition of cAMP production using standard cAMP quantitation assays. Agonists presenting a calcium transient or cAMP fluctuation are tested in vector control cells to determine if the response is unique to the transfected cells expressing receptor.

### *Example 53: ATP-binding assay*

[001266] The following assay may be used to assess ATP-binding activity of polypeptides of the invention.

10091504-030702

[001267] ATP-binding activity of the polypeptides of the invention may be detected using the ATP-binding assay described in U.S. Patent 5,858,719, which is herein incorporated by reference in its entirety. Briefly, ATP-binding to polypeptides of the invention is measured via photoaffinity labeling with 8-azido-ATP in a competition assay. Reaction mixtures containing 1 mg/ml of the ABC transport protein of the present invention are incubated with varying concentrations of ATP, or the non-hydrolyzable ATP analog adenylyl-5'-imidodiphosphate for 10 minutes at 4°C. A mixture of 8-azido-ATP (Sigma Chem. Corp., St. Louis, MO.) plus 8-azido-ATP (<sup>32</sup>P-ATP) (5 mCi/μmol, ICN, Irvine CA.) is added to a final concentration of 100 μM and 0.5 ml aliquots are placed in the wells of a porcelain spot plate on ice. The plate is irradiated using a short wave 254 nm UV lamp at a distance of 2.5 cm from the plate for two one-minute intervals with a one-minute cooling interval in between. The reaction is stopped by addition of dithiothreitol to a final concentration of 2mM. The incubations are subjected to SDS-PAGE electrophoresis, dried, and autoradiographed. Protein bands corresponding to the particular polypeptides of the invention are excised, and the radioactivity quantified. A decrease in radioactivity with increasing ATP or adenylyl-5'-imidodiphosphate provides a measure of ATP affinity to the polypeptides.

#### *Example 54: Small Molecule Screening*

[001268] This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and polypeptide of the invention.



10094504-030792

[001269] Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the invention. These methods comprise contacting such an agent with a polypeptide of the invention or fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or fragment thereof, by methods well known in the art. In such a competitive binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of a particular agent to bind to the polypeptides of the invention.

[001270] Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is herein incorporated by reference in its entirety. Briefly stated, large numbers of different small molecule test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with polypeptides of the invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

[001271] This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

#### *Example 55: Phosphorylation Assay*

[001272] In order to assay for phosphorylation activity of the polypeptides of the invention, a phosphorylation assay as described in U.S. Patent 5,958,405 (which is herein incorporated by reference) is utilized. Briefly, phosphorylation activity may be

measured by phosphorylation of a protein substrate using gamma-labeled  $^{32}\text{P}$ -ATP and quantitation of the incorporated radioactivity using a gamma radioisotope counter. The polypeptides of the invention are incubated with the protein substrate,  $^{32}\text{P}$ -ATP, and a kinase buffer. The  $^{32}\text{P}$  incorporated into the substrate is then separated from free  $^{32}\text{P}$ -ATP by electrophoresis, and the incorporated  $^{32}\text{P}$  is counted and compared to a negative control. Radioactivity counts above the negative control are indicative of phosphorylation activity of the polypeptides of the invention.

***Example 56: Detection of Phosphorylation Activity (Activation) of the Polypeptides of the Invention in the Presence of Polypeptide Ligands***

[001273] Methods known in the art or described herein may be used to determine the phosphorylation activity of the polypeptides of the invention. A preferred method of determining phosphorylation activity is by the use of the tyrosine phosphorylation assay as described in US 5,817,471 (incorporated herein by reference).

***Example 57: Identification Of Signal Transduction Proteins That Interact With Polypeptides Of The Present Invention***

[001274] The purified polypeptides of the invention are research tools for the identification, characterization and purification of additional signal transduction pathway proteins or receptor proteins. Briefly, labeled receptor PTK polypeptide is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, receptor PTK polypeptide is covalently coupled to a chromatography column. Cell-free extract derived from putative target cells, such as carcinoma tissues, is passed over the column, and molecules with appropriate affinity bind to the receptor PTK polypeptides, or specific phosphotyrosine-recognition domains thereof. The receptor PTK polypeptide interacting protein-complex is recovered from the column, dissociated, and the recovered molecule subjected to N-terminal protein sequencing. This amino acid sequence is then used to

identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

### ***Example 58: IL-6 Bioassay***

[001275] To test the proliferative effects of the polypeptides of the invention, the IL-6 Bioassay as described by Marz *et al.* is utilized (*Proc. Natl. Acad. Sci., U.S.A.*, 95:3251-56 (1998), which is herein incorporated by reference). Briefly, IL-6 dependent B9 murine cells are washed three times in IL-6 free medium and plated at a concentration of 5,000 cells per well in 50  $\mu$ l, and 50  $\mu$ l of the IL-6-like polypeptide is added. After 68 hrs. at 37°C, the number of viable cells is measured by adding the tetrazolium salt thiazolyl blue (MTT) and incubating for a further 4 hrs. at 37°C. B9 cells are lysed by SDS and optical density is measured at 570 nm. Controls containing IL-6 (positive) and no cytokine (negative) are utilized. Enhanced proliferation in the test sample(s) relative to the negative control is indicative of proliferative effects mediated by polypeptides of the invention.

### ***Example 59: Support of Chicken Embryo Neuron Survival***

[001276] To test whether sympathetic neuronal cell viability is supported by polypeptides of the invention, the chicken embryo neuronal survival assay of Senaldi *et al* is utilized (*Proc. Natl. Acad. Sci., U.S.A.*, 96:11458-63 (1998), which is herein incorporated by reference). Briefly, motor and sympathetic neurons are isolated from chicken embryos, resuspended in L15 medium (with 10% FCS, glucose, sodium selenite, progesterone, conalbumin, putrescine, and insulin; Life Technologies, Rockville, MD.) and Dulbecco's modified Eagles medium [with 10% FCS, glutamine, penicillin, and 25 mM Hepes buffer (pH 7.2); Life Technologies, Rockville, MD.], respectively, and incubated at 37°C in 5% CO<sub>2</sub> in the presence of different concentrations of the purified IL-6-like polypeptide, as well as a negative control lacking any cytokine. After 3 days, neuron survival is determined by evaluation of cellular morphology, and through the use of the colorimetric assay of Mosmann (Mossmann, T., *J. Immunol. Methods*, 65:55-63 (1983)). Enhanced neuronal cell

viability as compared to the controls lacking cytokine is indicative of the ability of the inventive purified IL-6-like polypeptide(s) to enhance the survival of neuronal cells.

### ***Example 60: Assay for Phosphatase Activity***

[001277] The following assay may be used to assess serine/threonine phosphatase (PTPase) activity of the polypeptides of the invention.

[001278] In order to assay for serine/threonine phosphatase (PTPase) activity, assays can be utilized which are widely known to those skilled in the art. For example, the serine/threonine phosphatase (PSPase) activity is measured using a PSPase assay kit from New England Biolabs, Inc. Myelin basic protein (MyBP), a substrate for PSPase, is phosphorylated on serine and threonine residues with cAMP-dependent Protein Kinase in the presence of [<sup>32</sup>P]ATP. Protein serine/threonine phosphatase activity is then determined by measuring the release of inorganic phosphate from 32P-labeled MyBP.

### ***Example 61: Interaction of Serine/Threonine Phosphatases with other Proteins***

[001279] The polypeptides of the invention with serine/threonine phosphatase activity as determined in Example 60 are research tools for the identification, characterization and purification of additional interacting proteins or receptor proteins, or other signal transduction pathway proteins. Briefly, labeled polypeptide(s) of the invention is useful as a reagent for the purification of molecules with which it interacts. In one embodiment of affinity purification, polypeptide of the invention is covalently coupled to a chromatography column. Cell-free extract derived from putative target cells, such as neural or liver cells, is passed over the column, and molecules with appropriate affinity bind to the polypeptides of the invention. The polypeptides of the invention -complex is recovered from the column, dissociated, and the recovered molecule subjected to N-terminal protein sequencing. This amino acid sequence is

then used to identify the captured molecule or to design degenerate oligonucleotide probes for cloning the relevant gene from an appropriate cDNA library.

### ***Example 62: Assaying for Heparanase Activity***

[001280] In order to assay for heparanase activity of the polypeptides of the invention, the heparanase assay described by Vlodavsky et al is utilized (Vlodavsky, I., et al., Nat. Med., 5:793-802 (1999)). Briefly, cell lysates, conditioned media or intact cells ( $1 \times 10^6$  cells per 35-mm dish) are incubated for 18 hrs at 37°C, pH 6.2-6.6, with  $^{35}\text{S}$ -labeled ECM or soluble ECM derived peak I proteoglycans. The incubation medium is centrifuged and the supernatant is analyzed by gel filtration on a Sepharose CL-6B column (0.9 x 30 cm). Fractions are eluted with PBS and their radioactivity is measured. Degradation fragments of heparan sulfate side chains are eluted from Sepharose 6B at  $0.5 < K_{av} < 0.8$  (peak II). Each experiment is done at least three times. Degradation fragments corresponding to "peak II," as described by Vlodavsky et al., is indicative of the activity of the polypeptides of the invention in cleaving heparan sulfate.

### ***Example 63: Immobilization of biomolecules***

[001281] This example provides a method for the stabilization of polypeptides of the invention in non-host cell lipid bilayer constructs (see, e.g., Bieri et al., Nature Biotech 17:1105-1108 (1999), hereby incorporated by reference in its entirety herein) which can be adapted for the study of polypeptides of the invention in the various functional assays described above. Briefly, carbohydrate-specific chemistry for biotinylation is used to confine a biotin tag to the extracellular domain of the polypeptides of the invention, thus allowing uniform orientation upon immobilization. A 50uM solution of polypeptides of the invention in washed membranes is incubated with 20 mM NaIO<sub>4</sub> and 1.5 mg/ml (4mM) BACH or 2 mg/ml (7.5mM) biotin-hydrazide for 1 hr at room temperature (reaction volume, 150ul). Then the sample is dialyzed (Pierce Slidealizer Cassett, 10 kDa cutoff; Pierce Chemical Co., Rockford IL) at 4C first for 5

h, exchanging the buffer after each hour, and finally for 12 h against 500 ml buffer R (0.15 M NaCl, 1 mM MgCl<sub>2</sub>, 10 mM sodium phosphate, pH7). Just before addition into a cuvette, the sample is diluted 1:5 in buffer ROG50 (Buffer R supplemented with 50 mM octylglucoside).

#### *Example 64: TAQMAN*

[001282] Quantitative PCR (QPCR). Total RNA from cells in culture are extracted by Trizol separation as recommended by the supplier (LifeTechnologies). (Total RNA is treated with DNase I (Life Technologies) to remove any contaminating genomic DNA before reverse transcription.) Total RNA (50 ng) is used in a one-step, 50ul, RT-QPCR, consisting of Taqman Buffer A (Perkin-Elmer; 50 mM KCl/10 mM Tris, pH 8.3), 5.5 mM MgCl<sub>2</sub>, 240 μM each dNTP, 0.4 units RNase inhibitor(Promega), 8%glycerol, 0.012% Tween-20, 0.05% gelatin, 0.3uM primers, 0.1uM probe, 0.025units Amplitaq Gold (Perkin-Elmer) and 2.5 units Superscript II reverse transcriptase (Life Technologies). As a control for genomic contamination, parallel reactions are setup without reverse transcriptase. The relative abundance of (unknown) and 18S RNAs are assessed by using the Applied Biosystems Prism 7700 Sequence Detection System (Livak, K. J., Flood, S. J., Marmaro, J., Giusti, W. & Deetz, K. (1995) PCR Methods Appl. 4, 357-362). Reactions are carried out at 48°C for 30 min, 95°C for 10 min, followed by 40 cycles of 95°C for 15s, 60°C for 1 min. Reactions are performed in triplicate.

[001283] Primers (f & r) and FRET probes sets are designed using Primer Express Software (Perkin-Elmer). Probes are labeled at the 5'-end with the reporter dye 6-FAM and on the 3'-end with the quencher dye TAMRA (Biosource International, Camarillo, CA or Perkin-Elmer).

#### *Example 65: Assays for Metalloproteinase Activity*

[001284] Metalloproteinases (EC 3.4.24.-) are peptide hydrolases which use metal ions, such as Zn<sup>2+</sup>, as the catalytic mechanism. Metalloproteinase activity of polypeptides

of the present invention can be assayed according to the following methods.

#### *Proteolysis of alpha-2-macroglobulin*

[001285] To confirm protease activity, purified polypeptides of the invention are mixed with the substrate alpha-2-macroglobulin (0.2 unit/ml; Boehringer Mannheim, Germany) in 1x assay buffer (50 mM HEPES, pH 7.5, 0.2 M NaCl, 10 mM CaCl<sub>2</sub>, 25  $\mu$ M ZnCl<sub>2</sub> and 0.05% Brij-35) and incubated at 37°C for 1-5 days. Trypsin is used as positive control. Negative controls contain only alpha-2-macroglobulin in assay buffer. The samples are collected and boiled in SDS-PAGE sample buffer containing 5% 2-mercaptoethanol for 5-min, then loaded onto 8% SDS-polyacrylamide gel. After electrophoresis the proteins are visualized by silver staining. Proteolysis is evident by the appearance of lower molecular weight bands as compared to the negative control.

#### *Inhibition of alpha-2-macroglobulin proteolysis by inhibitors of metalloproteinases*

[001286] Known metalloproteinase inhibitors (metal chelators (EDTA, EGTA, AND HgCl<sub>2</sub>), peptide metalloproteinase inhibitors (TIMP-1 and TIMP-2), and commercial small molecule MMP inhibitors) are used to characterize the proteolytic activity of polypeptides of the invention. The three synthetic MMP inhibitors used are: MMP inhibitor I, [IC<sub>50</sub> = 1.0  $\mu$ M against MMP-1 and MMP-8; IC<sub>50</sub> = 30  $\mu$ M against MMP-9; IC<sub>50</sub> = 150  $\mu$ M against MMP-3]; MMP-3 (stromelysin-1) inhibitor I [IC<sub>50</sub> = 5  $\mu$ M against MMP-3], and MMP-3 inhibitor II [K<sub>i</sub> = 130 nM against MMP-3]; inhibitors available through Calbiochem, catalog # 444250, 444218, and 444225, respectively). Briefly, different concentrations of the small molecule MMP inhibitors are mixed with purified polypeptides of the invention (50 $\mu$ g/ml) in 22.9  $\mu$ l of 1x HEPES buffer (50 mM HEPES, pH 7.5, 0.2 M NaCl, 10 mM CaCl<sub>2</sub>, 25  $\mu$ M ZnCl<sub>2</sub> and 0.05%Brij-35) and incubated at room temperature (24 °C) for 2-hr, then 7.1  $\mu$ l of substrate alpha-2-macroglobulin (0.2 unit/ml) is added and incubated at 37°C for 20-hr. The reactions are stopped by adding 4x sample buffer and boiled immediately for 5 minutes. After SDS-PAGE, the protein bands are visualized by silver stain.

#### *Synthetic Fluorogenic Peptide Substrates Cleavage Assay*

[001287] The substrate specificity for polypeptides of the invention with demonstrated

metalloproteinase activity can be determined using synthetic fluorogenic peptide substrates (purchased from BACHEM Bioscience Inc). Test substrates include, M-1985, M-2225, M-2105, M-2110, and M-2255. The first four are MMP substrates and the last one is a substrate of tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) converting enzyme (TACE). All the substrates are prepared in 1:1 dimethyl sulfoxide (DMSO) and water. The stock solutions are 50-500  $\mu$ M. Fluorescent assays are performed by using a Perkin Elmer LS 50B luminescence spectrometer equipped with a constant temperature water bath. The excitation  $\lambda$  is 328 nm and the emission  $\lambda$  is 393 nm. Briefly, the assay is carried out by incubating 176  $\mu$ l 1x HEPES buffer (0.2 M NaCl, 10 mM  $\text{CaCl}_2$ , 0.05% Brij-35 and 50 mM HEPES, pH 7.5) with 4  $\mu$ l of substrate solution (50  $\mu$ M) at 25  $^\circ\text{C}$  for 15 minutes, and then adding 20  $\mu$ l of a purified polypeptide of the invention into the assay cuvet. The final concentration of substrate is 1  $\mu$ M. Initial hydrolysis rates are monitored for 30-min.

***Example 66: Characterization of the cDNA contained in a deposited plasmid***

[001288] The size of the cDNA insert contained in a deposited plasmid may be routinely determined using techniques known in the art, such as PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the cDNA sequence. For example, two primers of 17-30 nucleotides derived from each end of the cDNA (i.e., hybridizable to the absolute 5' nucleotide or the 3' nucleotide end of the sequence of SEQ ID NO:X, respectively) are synthesized and used to amplify the cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25  $\mu$ l of reaction mixture with 0.5  $\mu$ g of the above cDNA template. A convenient reaction mixture is 1.5-5 mM  $\text{MgCl}_2$ , 0.01% (w/v) gelatin, 20  $\mu$ M each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of Taq polymerase. Thirty five cycles of PCR (denaturation at 94 degree C for 1 min; annealing at 55 degree C for 1 min; elongation at 72 degree C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified



product is analyzed by agarose gel electrophoresis. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA product.

[001289] Use of the above methodologies and/or other methodologies known in the art generates fragments from the clone corresponding to the approximate fragments described in Table 8, below. Accordingly, Table 8 provides a physical characterization of certain clones encompassed by the invention. The first column provides the unique clone identifier, "Clone ID NO:Z," for cDNA clones of the invention, as described in Table 1A. The second column provides the approximate size of the cDNA insert contained in the corresponding cDNA clone.

**TABLE 8**

Clone ID NO:Z	cDNA Insert Size:
HELGH89	300
HEMEF34	1200
HELGW31	1700
HELFN75	700
HELDK22	1000

[001290] It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

[001291] The entire disclosure of each document cited (including patents; patent applications, journal articles, abstracts, laboratory manuals, books, or other disclosures) in the Background of the Invention, Detailed Description, and Examples is hereby incorporated herein by reference. In addition, the CD-R copy of the

sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties. The specification and Sequence Listing of each of the following U.S. applications are herein incorporated by reference in their entirety: Application No. 60/179,065, filed on 31-Jan-2000; Application No. 60/180,628, filed on 04-Feb-2000; Application No. 60/214,886, filed on 28-Jun-2000; Application No. 60/217,487, filed on 11-Jul-2000; Application No. 60/225,758, filed on 14-Aug-2000; Application No. 60/220,963, filed on 26-Jul-2000; Application No. 60/217,496, filed on 11-Jul-2000; Application No. 60/225,447, filed on 14-Aug-2000; Application No. 60/218,290, filed on 14-Jul-2000; Application No. 60/225,757, filed on 14-Aug-2000; Application No. 60/226,868, filed on 22-Aug-2000; Application No. 60/216,647, filed on 07-Jul-2000; Application No. 60/225,267, filed on 14-Aug-2000; Application No. 60/216,880, filed on 07-Jul-2000; Application No. 60/225,270, filed on 14-Aug-2000; Application No. 60/251,869, filed on 08-Dec-2000; Application No. 60/235,834, filed on 27-Sep-2000; Application No. 60/234,274, filed on 21-Sep-2000; Application No. 60/234,223, filed on 21-Sep-2000; Application No. 60/228,924, filed on 30-Aug-2000; Application No. 60/224,518, filed on 14-Aug-2000; Application No. 60/236,369, filed on 29-Sep-2000; Application No. 60/224,519, filed on 14-Aug-2000; Application No. 60/220,964, filed on 26-Jul-2000; Application No. 60/241,809, filed on 20-Oct-2000; Application No. 60/249,299, filed on 17-Nov-2000; Application No. 60/236,327, filed on 29-Sep-2000; Application No. 60/241,785, filed on 20-Oct-2000; Application No. 60/244,617, filed on 01-Nov-2000; Application No. 60/225,268, filed on 14-Aug-2000; Application No. 60/236,368, filed on 29-Sep-2000; Application No. 60/251,856, filed on 08-Dec-2000; Application No. 60/251,868, filed on 08-Dec-2000; Application No. 60/229,344, filed on 01-Sep-2000; Application No. 60/234,997, filed on 25-Sep-2000; Application No. 60/229,343, filed on 01-Sep-2000; Application No. 60/229,345, filed on 01-Sep-2000; Application No. 60/229,287, filed on 01-Sep-2000; Application No. 60/229,513, filed on 05-Sep-2000; Application No. 60/231,413, filed on 08-Sep-2000; Application No. 60/229,509, filed on 05-Sep-2000; Application No. 60/236,367, filed on 29-Sep-2000; Application No. 60/237,039, filed on 02-Oct-2000; Application No. 60/237,038, filed on 02-Oct-2000; Application No. 60/236,370, filed on 29-Sep-2000; Application No. 60/236,802, filed on 02-Oct-2000; Application No. 60/237,037, filed

20070604 1051507

on 02-Oct-2000; Application No. 60/237,040, filed on 02-Oct-2000; Application No. 60/240,960, filed on 20-Oct-2000; Application No. 60/239,935, filed on 13-Oct-2000; Application No. 60/239,937, filed on 13-Oct-2000; Application No. 60/241,787, filed on 20-Oct-2000; Application No. 60/246,474, filed on 08-Nov-2000; Application No. 60/246,532, filed on 08-Nov-2000; Application No. 60/249,216, filed on 17-Nov-2000; Application No. 60/249,210, filed on 17-Nov-2000; Application No. 60/226,681, filed on 22-Aug-2000; Application No. 60/225,759, filed on 14-Aug-2000; Application No. 60/225,213, filed on 14-Aug-2000; Application No. 60/227,182, filed on 22-Aug-2000; Application No. 60/225,214, filed on 14-Aug-2000; Application No. 60/235,836, filed on 27-Sep-2000; Application No. 60/230,438, filed on 06-Sep-2000; Application No. 60/215,135, filed on 30-Jun-2000; Application No. 60/225,266, filed on 14-Aug-2000; Application No. 60/249,218, filed on 17-Nov-2000; Application No. 60/249,208, filed on 17-Nov-2000; Application No. 60/249,213, filed on 17-Nov-2000; Application No. 60/249,212, filed on 17-Nov-2000; Application No. 60/249,207, filed on 17-Nov-2000; Application No. 60/249,245, filed on 17-Nov-2000; Application No. 60/249,244, filed on 17-Nov-2000; Application No. 60/249,217, filed on 17-Nov-2000; Application No. 60/249,211, filed on 17-Nov-2000; Application No. 60/249,215, filed on 17-Nov-2000; Application No. 60/249,264, filed on 17-Nov-2000; Application No. 60/249,214, filed on 17-Nov-2000; Application No. 60/249,297, filed on 17-Nov-2000; Application No. 60/232,400, filed on 14-Sep-2000; Application No. 60/231,242, filed on 08-Sep-2000; Application No. 60/232,081, filed on 08-Sep-2000; Application No. 60/232,080, filed on 08-Sep-2000; Application No. 60/231,414, filed on 08-Sep-2000; Application No. 60/231,244, filed on 08-Sep-2000; Application No. 60/233,064, filed on 14-Sep-2000; Application No. 60/233,063, filed on 14-Sep-2000; Application No. 60/232,397, filed on 14-Sep-2000; Application No. 60/232,399, filed on 14-Sep-2000; Application No. 60/232,401, filed on 14-Sep-2000; Application No. 60/241,808, filed on 20-Oct-2000; Application No. 60/241,826, filed on 20-Oct-2000; Application No. 60/241,786, filed on 20-Oct-2000; Application No. 60/241,221, filed on 20-Oct-2000; Application No. 60/246,475, filed on 08-Nov-2000; Application No. 60/231,243, filed on 08-Sep-2000; Application No. 60/233,065, filed on 14-Sep-2000; Application No. 60/232,398, filed on 14-Sep-2000; Application No. 60/234,998, filed

on 25-Sep-2000; Application No. 60/246,477, filed on 08-Nov-2000; Application No. 60/246,528, filed on 08-Nov-2000; Application No. 60/246,525, filed on 08-Nov-2000; Application No. 60/246,476, filed on 08-Nov-2000; Application No. 60/246,526, filed on 08-Nov-2000; Application No. 60/249,209, filed on 17-Nov-2000; Application No. 60/246,527, filed on 08-Nov-2000; Application No. 60/246,523, filed on 08-Nov-2000; Application No. 60/246,524, filed on 08-Nov-2000; Application No. 60/246,478, filed on 08-Nov-2000; Application No. 60/246,609, filed on 08-Nov-2000; Application No. 60/246,613, filed on 08-Nov-2000; Application No. 60/249,300, filed on 17-Nov-2000; Application No. 60/249,265, filed on 17-Nov-2000; Application No. 60/246,610, filed on 08-Nov-2000; Application No. 60/246,611, filed on 08-Nov-2000; Application No. 60/230,437, filed on 06-Sep-2000; Application No. 60/251,990, filed on 08-Dec-2000; Application No. 60/251,988, filed on 05-Dec-2000; Application No. 60/251,030, filed on 05-Dec-2000; Application No. 60/251,479, filed on 06-Dec-2000; Application No. 60/256,719, filed on 05-Dec-2000; Application No. 60/250,160, filed on 01-Dec-2000; Application No. 60/251,989, filed on 08-Dec-2000; Application No. 60/250,391, filed on 01-Dec-2000; Application No. 60/254,097, filed on 11-Dec-2000; and Application No. 09/764,869, filed 17-Jan-2001.

**[01292]** Moreover, the microfiche copy and the corresponding computer readable form of the Sequence Listing of U.S. Application Serial No. 60/179,065, and the hard copy of and the corresponding computer readable form of the Sequence Listing of U.S. Application Serial No. 60/180,628 are also incorporated herein by reference in their entireties.